

FOURTH SEMESTER B.C.A. DEGREE EXAMINATION, APRIL/MAY 2005

(Vocational Course)

Optional Subject : Statistics

Paper VII—ESTIMATION

Time : Three Hours

Maximum : 90 Marks

*Each unit carries 50 marks.**Not more than 30 marks will be awarded from from each unit.**Statistical tables will be provided on request.***Unit I**

1. Let X_1, X_2, \dots, X_n be a random sample from a population with mean μ and finite variance σ^2 . Show that $\bar{X} = (X_1 + X_2 + \dots + X_n)/n$ is unbiased for μ but $S^2 = \sum_{i=1}^n (X_i - \bar{X})^2/n$ is not unbiased for σ^2 . Suggest an unbiased estimator for σ^2 .

(8 marks)

2. Define consistency of an estimator. State and prove an sufficient condition for consistency.

(8 marks)

3. If there exist two unbiased estimators for a parameters θ , show that there exist infinitely many unbiased estimators for θ .

(6 marks)

4. State Fisher-Neymann factorization criterion. Show that sample mean is a sufficient estimator for θ in the case of a Poisson population with mean θ .

(6 marks)

5. Give an example of a sufficient estimator which is neither unbiased nor consistent. (6 marks)

6. If X_1, X_2 is a random sample from a normal population with mean μ and variance 1, show that

$$T_1 = (X_1 + X_2)/2 \text{ and } T_2 = 2X_1 - X_2 \text{ are unbiased estimators of } \mu. \text{ Compare their efficiencies.}$$

(8 marks)

7. Let X_1, X_2, \dots, X_n be a random sample from a population with p.d.f. $f(x; \theta) = \begin{cases} \frac{1}{\theta}, & 0 < x < \theta \\ 0, & \text{elsewhere} \end{cases}$ where $\theta > 0$, Show that $T = \max(X_1, X_2, \dots, X_n)$ is sufficient for θ .

(8 marks)

Turn over

Unit II

8. State and prove Cramer-Rao inequality. (7 marks)
9. Distinguish between minimum variance bound estimator and uniformly minimum variance unbiased estimator. Give an example in which these are equal. (7 marks)
10. State and prove Rao-Blackwell theorem. (7 marks)
11. Let X_1, X_2, \dots, X_n be a random sample from $f(x, \theta) = \begin{cases} \theta(1-\theta)^x, & x = 0, 1, 2, \dots \\ 0, & \text{elsewhere} \end{cases}$ where $0 < \theta < 1$. Obtain Cramer-Rao lower bound to the variance of an unbiased estimator of θ^2 . (6 marks)
12. Explain minimum-chisquare method of estimation. (4 marks)
13. Let $f(x, \theta) = \begin{cases} \frac{2x}{\theta^2}, & 0 < x < \theta \\ 0, & \text{elsewhere} \end{cases}$ where $\theta > 0$. Obtain MLE of θ based on a sample of size n . (6 marks)
14. Let X_1, X_2, \dots, X_n be a random sample from $f(x) = \begin{cases} \frac{m^p}{\Gamma(p)} e^{-mx} x^{p-1}, & 0 < x < \infty \\ 0, & \text{elsewhere} \end{cases}$ where $m > 0$, $p > 0$. Obtain moment estimators of m and p . (6 marks)
15. Let X_1, X_2, \dots, X_n be a random sample from $f(x, \theta) = \begin{cases} \theta x^{\theta-1} & 0 < x < 1 \\ 0, & \text{elsewhere} \end{cases}$ where $\theta > 0$. Obtain MLE of θ . Check whether the MLE is unbiased and consistent. (7 marks)

Unit III

16. Distinguish between Point estimation and Interval estimation. (6 marks)
17. Stating the assumptions clearly, obtain confidence interval with confidence coefficient 0.95 for the parameter μ in $f(x; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\}$, $-\infty < x < \infty$. (8 marks)
18. A population follows normal distribution with mean μ and variance 25. A sample of size 10 from the population has mean 58. For the confidence interval (52, 64) for μ , compute the confidence coefficient. (6 marks)

19. Let X_1, X_2, \dots, X_n be a random sample from a normal population with mean μ and variance σ^2 . Obtain 95 % confidence interval for σ^2 , when (a) μ is known ; (b) μ is unknown. (10 marks)
20. Obtain a 90 % confidence interval for the correlation P. Given that the sample correlation based on a sample of size 18 is 0.58. (8 marks)
21. In an exit poll out of 300 voters 170 favour candidate A. Obtain 95 % confidence interval for the proportion of votes that the candidate A will be getting in the actual election. (6 marks)
22. Stating the assumptions clearly explain how do you construct large sample confidence intervals. (6 marks)