



## Problems and solutions: INMO 2013

**Problem 1.** Let  $\Gamma_1$  and  $\Gamma_2$  be two circles touching each other externally at  $R$ . Let  $l_1$  be a line which is tangent to  $\Gamma_2$  at  $P$  and passing through the center  $O_1$  of  $\Gamma_1$ . Similarly, let  $l_2$  be a line which is tangent to  $\Gamma_2$  at  $Q$  and passing through the center  $O_2$  of  $\Gamma_2$ . Suppose  $l_1$  and  $l_2$  are not parallel and intersect at  $K$ . If  $KP = KQ$ , prove that the triangle  $PQR$  is equilateral.

**Solution.** Suppose that  $P$  and  $Q$  lie on the opposite sides of line joining  $O_1$  and  $O_2$ . By symmetry we may assume that the configuration is as shown in the figure below. Then we have  $KP > KO_1 > KQ$  since  $KO_1$  is the hypotenuse of triangle  $KQO_1$ . This is a contradiction to the given assumption, and therefore  $P$  and  $Q$  lie on the same side of the line joining  $O_1$  and  $O_2$ .



Since  $KP = KQ$  it follows that  $K$  lies on the radical axis of the given circles, which is the common tangent at  $R$ . Therefore  $KP = KQ = KR$  and hence  $K$  is the circumcenter of  $\triangle PQR$ .



On the other hand,  $\triangle KQO_1$  and  $\triangle KRO_1$  are both right-angled triangles with  $KQ = KR$  and  $QO_1 = RO_1$ , and hence the two triangles are congruent. Therefore  $\widehat{QKO_1} = \widehat{RKO_1}$ , so  $KO_1$ , and hence  $PK$  is perpendicular to  $QR$ . Similarly,  $QK$  is perpendicular to  $PR$ , so it follows that  $K$  is the orthocenter of  $\triangle PQR$ . Hence we have that  $\triangle PQR$  is equilateral.  $\square$

**Alternate solution.** We again rule out the possibility that  $P$  and  $Q$  are on the opposite side of the line joining  $O_1O_2$ , and assume that they are on the same side.

Observe that  $\triangle KPO_2$  is congruent to  $\triangle KQO_1$  (since  $KP = KQ$ ). Therefore  $O_1P = O_2Q = r$  (say). In  $\triangle O_1O_2Q$ , we have  $\widehat{O_1QO_2} = \pi/2$  and  $R$  is the midpoint of the hypotenuse, so  $RQ = RO_1 = r$ . Therefore  $\triangle O_1RQ$  is equilateral, so  $\widehat{QRO_1} = \pi/3$ . Similarly,  $PR = r$  and  $\widehat{PRO_2} = \pi/3$ , hence  $\widehat{PRQ} = \pi/3$ . Since  $PR = QR$  it follows that  $\triangle PQR$  is equilateral.  $\square$

**Problem 2.** Find all positive integers  $m$ ,  $n$ , and primes  $p \geq 5$  such that

$$m(4m^2 + m + 12) = 3(p^n - 1).$$

**Solution.** Rewriting the given equation we have

$$4m^3 + m^2 + 12m + 3 = 3p^n.$$

The left hand side equals  $(4m + 1)(m^2 + 3)$ .

Suppose that  $(4m + 1, m^2 + 3) = 1$ . Then  $(4m + 1, m^2 + 3) = (3p^n, 1), (3, p^n), (p^n, 3)$  or  $(1, 3p^n)$ , a contradiction since  $4m + 1, m^2 + 3 \geq 4$ . Therefore  $(4m + 1, m^2 + 3) > 1$ .

Since  $4m + 1$  is odd we have  $(4m + 1, m^2 + 3) = (4m + 1, 16m^2 + 48) = (4m + 1, 49) = 7$  or  $49$ . This proves that  $p = 7$ , and  $4m + 1 = 3 \cdot 7^k$  or  $7^k$  for some natural number  $k$ . If  $(4m + 1, 49) = 7$  then we have  $k = 1$  and  $4m + 1 = 21$  which does not lead to a solution. Therefore  $(4m + 1, m^2 + 3) = 49$ . If  $7^3$  divides  $4m + 1$  then it does not divide  $m^2 + 3$ , so we get  $m^2 + 3 \leq 3 \cdot 7^2 < 7^3 \leq 4m + 1$ . This implies  $(m - 2)^2 < 2$ , so  $m \leq 3$ , which does not lead to a solution. Therefore we have  $4m + 1 = 49$  which implies  $m = 12$  and  $n = 4$ . Thus  $(m, n, p) = (12, 4, 7)$  is the only solution.  $\square$

**Problem 3.** Let  $a, b, c, d$  be positive integers such that  $a \geq b \geq c \geq d$ . Prove that the equation  $x^4 - ax^3 - bx^2 - cx - d = 0$  has no integer solution.

**Solution.** Suppose that  $m$  is an integer root of  $x^4 - ax^3 - bx^2 - cx - d = 0$ . As  $d \neq 0$ , we have  $m \neq 0$ . Suppose now that  $m > 0$ . Then  $m^4 - am^3 = bm^2 + cm + d > 0$  and hence  $m > a \geq d$ . On the other hand  $d = m(m^3 - am^2 - bm - c)$  and hence  $m$  divides  $d$ , so  $m \leq d$ , a contradiction. If  $m < 0$ , then writing  $n = -m > 0$  we have  $n^4 + an^3 - bn^2 + cn - d = n^4 + n^2(an - b) + (cn - d) > 0$ , a contradiction. This proves that the given polynomial has no integer roots.  $\square$

**Problem 4.** Let  $n$  be a positive integer. Call a nonempty subset  $S$  of  $\{1, 2, \dots, n\}$  good if the arithmetic mean of the elements of  $S$  is also an integer. Further let  $t_n$  denote the number of good subsets of  $\{1, 2, \dots, n\}$ . Prove that  $t_n$  and  $n$  are both odd or both even.

**Solution.** We show that  $T_n - n$  is even. Note that the subsets  $\{1\}, \{2\}, \dots, \{n\}$  are good. Among the other good subsets, let  $A$  be the collection of subsets with an integer average which belongs to the subset, and let  $B$  be the collection of subsets with an integer average which is not a member of the subset. Then there is a bijection between  $A$  and  $B$ , because removing the average takes a member of  $A$  to a member of  $B$ ; and including the average in a member of  $B$  takes it to its inverse. So  $T_n - n = |A| + |B|$  is even.  $\square$

**Alternate solution.** Let  $S = \{1, 2, \dots, n\}$ . For a subset  $A$  of  $S$ , let  $\bar{A} = \{n + 1 - a | a \in A\}$ . We call a subset  $A$  symmetric if  $\bar{A} = A$ . Note that the arithmetic mean of a symmetric subset is  $(n + 1)/2$ . Therefore, if  $n$  is even, then there are no symmetric good subsets, while if  $n$  is odd then every symmetric subset is good.

If  $A$  is a proper good subset of  $S$ , then so is  $\bar{A}$ . Therefore, all the good subsets that are not symmetric can be paired. If  $n$  is even then this proves that  $t_n$  is even. If  $n$  is odd, we have to show that there are odd number of symmetric subsets. For this, we note that a symmetric subset contains the element  $(n + 1)/2$  if and only if it has odd number of elements. Therefore, for any natural number  $k$ , the number of symmetric subsets of size  $2k$  equals the number of symmetric subsets of size  $2k + 1$ . The result now follows since there is exactly one symmetric subset with only one element.  $\square$

**Problem 5.** In an acute triangle  $ABC$ ,  $O$  is the circumcenter,  $H$  is the orthocenter and  $G$  is the centroid. Let  $OD$  be perpendicular to  $BC$  and  $HE$  be perpendicular to  $CA$ , with  $D$  on  $BC$  and  $E$  on  $CA$ . Let  $F$  be the midpoint of  $AB$ . Suppose the areas of triangles  $ODC$ ,  $HEA$  and  $GFB$  are equal. Find all the possible values of  $\hat{C}$ .

**Solution.** Let  $R$  be the circumradius of  $\triangle ABC$  and  $\Delta$  its area. We have  $OD = R \cos A$  and  $DC = \frac{a}{2}$ , so

$$[ODC] = \frac{1}{2} \cdot OD \cdot DC = \frac{1}{2} \cdot R \cos A \cdot R \sin A = \frac{1}{2} R^2 \sin A \cos A. \quad (1)$$

Again  $HE = 2R \cos C \cos A$  and  $EA = c \cos A$ . Hence

$$[HEA] = \frac{1}{2} \cdot HE \cdot EA = \frac{1}{2} \cdot 2R \cos C \cos A \cdot c \cos A = 2R^2 \sin C \cos C \cos^2 A. \quad (2)$$

Further

$$[GFB] = \frac{\Delta}{6} = \frac{1}{6} \cdot 2R^2 \sin A \sin B \sin C = \frac{1}{3} R^2 \sin A \sin B \sin C. \quad (3)$$

Equating (1) and (2) we get  $\tan A = 4 \sin C \cos C$ . And equating (1) and (3), and using this relation we get

$$\begin{aligned} 3 \cos A &= 2 \sin B \sin C = 2 \sin(C + A) \sin C \\ &= 2(\sin C + \cos C \tan A) \sin C \cos A \\ &= 2 \sin^2 C (1 + 4 \cos^2 C) \cos A. \end{aligned}$$

Since  $\cos A \neq 0$  we get  $3 = 2t(-4t + 5)$  where  $t = \sin^2 C$ . This implies  $(4t - 3)(2t - 1) = 0$  and therefore, since  $\sin C > 0$ , we get  $\sin C = \sqrt{3}/2$  or  $\sin C = 1/\sqrt{2}$ . Because  $\triangle ABC$  is acute, it follows that  $\widehat{C} = \pi/3$  or  $\pi/4$ .

We observe that the given conditions are satisfied in an equilateral triangle, so  $\widehat{C} = \pi/3$  is a possibility. Also, the conditions are satisfied in a triangle where  $\widehat{C} = \pi/4$ ,  $\widehat{A} = \tan^{-1} 2$  and  $\widehat{B} = \tan^{-1} 3$ . Therefore  $\widehat{C} = \pi/4$  is also a possibility.

Thus the two possible values of  $\widehat{C}$  are  $\pi/3$  and  $\pi/4$ . □

**Problem 6.** Let  $a, b, c, x, y, z$  be positive real numbers such that  $a + b + c = x + y + z$  and  $abc = xyz$ . Further, suppose that  $a \leq x < y < z \leq c$  and  $a < b < c$ . Prove that  $a = x, b = y$  and  $c = z$ .

**Solution.** Let

$$f(t) = (t - x)(t - y)(t - z) - (t - a)(t - b)(t - c).$$

Then  $f(t) = kt$  for some constant  $k$ . Note that  $ka = f(a) = (a - x)(a - y)(a - z) \leq 0$  and hence  $k \leq 0$ . Similarly,  $kc = f(c) = (c - x)(c - y)(c - z) \geq 0$  and hence  $k \geq 0$ . Combining the two, it follows that  $k = 0$  and that  $f(a) = f(c) = 0$ . These equalities imply that  $a = x$  and  $c = z$ , and then it also follows that  $b = y$ . □