



## Problems and Solutions: INMO-2012

1. Let  $ABCD$  be a quadrilateral inscribed in a circle. Suppose  $AB = \sqrt{2 + \sqrt{2}}$  and  $AB$  subtends  $135^\circ$  at the centre of the circle. Find the maximum possible area of  $ABCD$ .



**Solution:** Let  $O$  be the centre of the circle in which  $ABCD$  is inscribed and let  $R$  be its radius. Using cosine rule in triangle  $AOB$ , we have

$$2 + \sqrt{2} = 2R^2(1 - \cos 135^\circ) = R^2(2 + \sqrt{2}).$$

Hence  $R = 1$ .

Consider quadrilateral  $ABCD$  as in the second figure above. Join  $AC$ . For  $[ADC]$  to be maximum, it is clear that  $D$  should be the mid-point of the arc  $AC$  so that its distance from the segment  $AC$  is maximum. Hence  $AD = DC$  for  $[ABCD]$  to be maximum. Similarly, we conclude that  $BC = CD$ . Thus  $BC = CD = DA$  which fixes the quadrilateral  $ABCD$ . Therefore each of the sides  $BC$ ,  $CD$ ,  $DA$  subtends equal angles at the centre  $O$ .

Let  $\angle BOC = \alpha$ ,  $\angle COD = \beta$  and  $\angle DOA = \gamma$ . Observe that

$$[ABCD] = [AOB] + [BOC] + [COD] + [DOA] = \frac{1}{2} \sin 135^\circ + \frac{1}{2} (\sin \alpha + \sin \beta + \sin \gamma).$$

Now  $[ABCD]$  has maximum area if and only if  $\alpha = \beta = \gamma = (360^\circ - 135^\circ)/3 = 75^\circ$ . Thus

$$[ABCD] = \frac{1}{2} \sin 135^\circ + \frac{3}{2} \sin 75^\circ = \frac{1}{2} \left( \frac{1}{\sqrt{2}} + 3 \frac{\sqrt{3} + 1}{2\sqrt{2}} \right) = \frac{5 + 3\sqrt{3}}{4\sqrt{2}}.$$

Alternatively, we can use Jensen's inequality. Observe that  $\alpha, \beta, \gamma$  are all less than  $180^\circ$ . Since  $\sin x$  is concave on  $(0, \pi)$ , Jensen's inequality gives

$$\frac{\sin \alpha + \sin \beta + \sin \gamma}{3} \leq \sin \left( \frac{\alpha + \beta + \gamma}{3} \right) = \sin 75^\circ.$$

Hence

$$[ABCD] \leq \frac{1}{2\sqrt{2}} + \frac{3}{2} \sin 75^\circ = \frac{5 + 3\sqrt{3}}{4\sqrt{2}},$$

with equality if and only if  $\alpha = \beta = \gamma = 75^\circ$ .

2. Let  $p_1 < p_2 < p_3 < p_4$  and  $q_1 < q_2 < q_3 < q_4$  be two sets of prime numbers such that  $p_4 - p_1 = 8$  and  $q_4 - q_1 = 8$ . Suppose  $p_1 > 5$  and  $q_1 > 5$ . Prove that 30 divides  $p_1 - q_1$ .

**Solution:** Since  $p_4 - p_1 = 8$ , and no prime is even, we observe that  $\{p_1, p_2, p_3, p_4\}$  is a subset of  $\{p_1, p_1 + 2, p_1 + 4, p_1 + 6, p_1 + 8\}$ . Moreover  $p_1$  is larger than 3. If  $p_1 \equiv 1 \pmod{3}$ , then  $p_1 + 2$  and  $p_1 + 8$  are divisible by 3. Hence we do not get 4 primes in the set  $\{p_1, p_1 + 2, p_1 + 4, p_1 + 6, p_1 + 8\}$ . Thus  $p_1 \equiv 2 \pmod{3}$  and  $p_1 + 4$  is not a prime. We get  $p_2 = p_1 + 2, p_3 = p_1 + 6, p_4 = p_1 + 8$ .

Consider the remainders of  $p_1, p_1 + 2, p_1 + 6, p_1 + 8$  when divided by 5. If  $p_1 \equiv 2 \pmod{5}$ , then  $p_1 + 8$  is divisible by 5 and hence is not a prime. If  $p_1 \equiv 3 \pmod{5}$ , then  $p_1 + 2$  is divisible by 5. If  $p_1 \equiv 4 \pmod{5}$ , then  $p_1 + 6$  is divisible by 5. Hence the only possibility is  $p_1 \equiv 1 \pmod{5}$ .

Thus we see that  $p_1 \equiv 1 \pmod{2}$ ,  $p_1 \equiv 2 \pmod{3}$  and  $p_1 \equiv 1 \pmod{5}$ . We conclude that  $p_1 \equiv 11 \pmod{30}$ .

Similarly  $q_1 \equiv 11 \pmod{30}$ . It follows that 30 divides  $p_1 - q_1$ .

3. Define a sequence  $\langle f_0(x), f_1(x), f_2(x), \dots \rangle$  of functions by

$$f_0(x) = 1, \quad f_1(x) = x, \quad (f_n(x))^2 - 1 = f_{n+1}(x)f_{n-1}(x), \quad \text{for } n \geq 1.$$

Prove that each  $f_n(x)$  is a polynomial with integer coefficients.

**Solution:** Observe that

$$f_n^2(x) - f_{n-1}(x)f_{n+1}(x) = 1 = f_{n-1}^2(x) - f_{n-2}(x)f_n(x).$$

This gives

$$f_n(x) \left( f_n(x) + f_{n-2}(x) \right) = f_{n-1} \left( f_{n-1}(x) + f_{n+1}(x) \right).$$

We write this as

$$\frac{f_{n-1}(x) + f_{n+1}(x)}{f_n(x)} = \frac{f_{n-2}(x) + f_n(x)}{f_{n-1}(x)}.$$

Using induction, we get

$$\frac{f_{n-1}(x) + f_{n+1}(x)}{f_n(x)} = \frac{f_0(x) + f_2(x)}{f_1(x)}.$$

Observe that

$$f_2(x) = \frac{f_1^2(x) - 1}{f_0(x)} = x^2 - 1.$$

Hence

$$\frac{f_{n-1}(x) + f_{n+1}(x)}{f_n(x)} = \frac{1 + (x^2 - 1)}{x} = x.$$

Thus we obtain

$$f_{n+1}(x) = xf_n(x) - f_{n-1}(x).$$

Since  $f_0(x)$ ,  $f_1(x)$  and  $f_2(x)$  are polynomials with integer coefficients, induction again shows that  $f_n(x)$  is a polynomial with integer coefficients.

**Note:** We can get  $f_n(x)$  explicitly:

$$f_n(x) = x^n - \binom{n-1}{1}x^{n-2} + \binom{n-2}{2}x^{n-4} - \binom{n-3}{3}x^{n-6} + \dots$$

4. Let  $ABC$  be a triangle. An interior point  $P$  of  $ABC$  is said to be **good** if we can find exactly 27 rays emanating from  $P$  intersecting the sides of the triangle  $ABC$  such that the triangle is divided by these rays into 27 smaller triangles of equal area. Determine the number of **good** points for a given triangle  $ABC$ .

**Solution:** Let  $P$  be a good point. Let  $l, m, n$  be respectively the number of parts the sides  $BC, CA, AB$  are divided by the rays starting from  $P$ . Note that a ray must pass through each of the vertices the triangle  $ABC$ ; otherwise we get some quadrilaterals.

Let  $h_1$  be the distance of  $P$  from  $BC$ . Then  $h_1$  is the height for all the triangles with their bases on  $BC$ . Equality of areas implies that all these bases have equal length. If we denote this by  $x$ , we get  $lx = a$ . Similarly, taking  $y$  and  $z$  as the lengths of the bases of triangles on  $CA$  and  $AB$  respectively, we get  $my = b$  and  $nz = c$ . Let  $h_2$  and  $h_3$  be the distances of  $P$  from  $CA$  and  $AB$  respectively. Then

$$h_1x = h_2y = h_3z = \frac{2\Delta}{27},$$

where  $\Delta$  denotes the area of the triangle  $ABC$ . These lead to

$$h_1 = \frac{2\Delta}{27} \frac{l}{a}, \quad h_1 = \frac{2\Delta}{27} \frac{m}{b}, \quad h_1 = \frac{2\Delta}{27} \frac{n}{c}.$$

But

$$\frac{2\Delta}{a} = h_a, \quad \frac{2\Delta}{b} = h_b, \quad \frac{2\Delta}{c} = h_c.$$

Thus we get

$$\frac{h_1}{h_a} = \frac{l}{27}, \quad \frac{h_2}{h_b} = \frac{m}{27}, \quad \frac{h_3}{h_c} = \frac{n}{27}.$$

However, we also have

$$\frac{h_1}{h_a} = \frac{[PBC]}{\Delta}, \quad \frac{h_2}{h_b} = \frac{[PCA]}{\Delta}, \quad \frac{h_3}{h_c} = \frac{[PAB]}{\Delta}.$$

Adding these three relations,

$$\frac{h_1}{h_a} + \frac{h_2}{h_b} + \frac{h_3}{h_c} = 1.$$

Thus

$$\frac{l}{27} + \frac{m}{27} + \frac{n}{27} = \frac{h_1}{h_a} + \frac{h_2}{h_b} + \frac{h_3}{h_c} = 1.$$

We conclude that  $l + m + n = 27$ . Thus every **good** point  $P$  determines a partition  $(l, m, n)$  of 27 such that there are  $l, m, n$  equal segments respectively on  $BC, CA, AB$ .

Conversely, take any partition  $(l, m, n)$  of 27. Divide  $BC, CA, AB$  respectively in to  $l, m, n$  equal parts. Define

$$h_1 = \frac{2l\Delta}{27a}, \quad h_2 = \frac{2m\Delta}{27b}.$$

Draw a line parallel to  $BC$  at a distance  $h_1$  from  $BC$ ; draw another line parallel to  $CA$  at a distance  $h_2$  from  $CA$ . Both lines are drawn such that they intersect at a point  $P$  inside the triangle  $ABC$ . Then

$$[PBC] = \frac{1}{2}ah_1 = \frac{l\Delta}{27}, \quad [PCA] = \frac{m\Delta}{27}.$$

Hence

$$[PAB] = \frac{n\Delta}{27}.$$

This shows that the distance of  $P$  from  $AB$  is


$$h_3 = \frac{2n\Delta}{27c}.$$

Therefore each triangle with base on  $CA$  has area  $\frac{\Delta}{27}$ . We conclude that all the triangles which partitions  $ABC$  have equal areas. Hence  $P$  is a **good** point.

Thus the number of **good** points is equal to the number of positive integral solutions of the equation  $l + m + n = 27$ . This is equal to

$$\binom{26}{2} = 325.$$

5. Let  $ABC$  be an acute-angled triangle, and let  $D, E, F$  be points on  $BC, CA, AB$  respectively such that  $AD$  is the median,  $BE$  is the internal angle bisector and  $CF$  is the altitude. Suppose  $\angle FDE = \angle C$ ,  $\angle DEF = \angle A$  and  $\angle EFD = \angle B$ . Prove that  $ABC$  is equilateral.



**Solution:** Since  $\triangle BFC$  is right-angled at  $F$ , we have  $FD = BD = CD = a/2$ . Hence  $\angle BFD = \angle B$ . Since  $\angle EFD = \angle B$ , we have  $\angle AFE = \pi - 2\angle B$ . Since  $\angle DEF = \angle A$ , we also get  $\angle CED = \pi - 2\angle B$ . Applying sine rule in  $\triangle DEF$ , we have

$$\frac{DF}{\sin A} = \frac{FE}{\sin C} = \frac{DE}{\sin B}.$$

Thus we get  $FE = c/2$  and  $DE = b/2$ . Sine rule in  $\triangle CED$  gives

$$\frac{DE}{\sin C} = \frac{CD}{\sin(\pi - 2B)}.$$

Thus  $(b/\sin C) = (a/2 \sin B \cos B)$ . Solving for  $\cos B$ , we have

$$\cos B = \frac{a \sin c}{2b \sin B} = \frac{ac}{2b^2}.$$

Similarly, sine rule in  $\triangle AEF$  gives

$$\frac{EF}{\sin A} = \frac{AE}{\sin(\pi - 2B)}.$$

This gives (since  $AE = bc/(a + c)$ ), as earlier,

$$\cos B = \frac{a}{a + c}.$$

Comparing the two values of  $\cos B$ , we get  $2b^2 = c(a + c)$ . We also have

$$c^2 + a^2 - b^2 = 2ca \cos B = \frac{2a^2c}{a + c}.$$

Thus

$$4a^2c = (a + c)(2c^2 + 2a^2 - 2b^2) = (a + c)(2c^2 + 2a^2 - c(a + c)).$$

This reduces to  $2a^3 - 3a^2c + c^3 = 0$ . Thus  $(a - c)^2(2a + c) = 0$ . We conclude that  $a = c$ . Finally

$$2b^2 = c(a + c) = 2c^2.$$

We thus get  $b = c$  and hence  $a = c = b$ . This shows that  $\triangle ABC$  is equilateral.

6. Let  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  be a function satisfying  $f(0) \neq 0$ ,  $f(1) = 0$  and

(i)  $f(xy) + f(x)f(y) = f(x) + f(y)$ ;

(ii)  $(f(x - y) - f(0))f(x)f(y) = 0$ ,

for all  $x, y \in \mathbb{Z}$ , simultaneously.

(a) Find the set of all possible values of the function  $f$ .

(b) If  $f(10) \neq 0$  and  $f(2) = 0$ , find the set of all integers  $n$  such that  $f(n) \neq 0$ .

**Solution:** Setting  $y = 0$  in the condition (ii), we get

$$(f(x) - f(0))f(x) = 0,$$

for all  $x$  (since  $f(0) \neq 0$ ). Thus either  $f(x) = 0$  or  $f(x) = f(0)$ , for all  $x \in \mathbb{Z}$ . Now taking  $x = y = 0$  in (i), we see that  $f(0) + f(0)^2 = 2f(0)$ . This shows

that  $f(0) = 0$  or  $f(0) = 1$ . Since  $f(0) \neq 0$ , we must have  $f(0) = 1$ . We conclude that

either  $f(x) = 0$  or  $f(x) = 1$  for each  $x \in \mathbb{Z}$ .

This shows that the set of all possible value of  $f(x)$  is  $\{0, 1\}$ . This completes (a).

Let  $S = \{n \in \mathbb{Z} \mid f(n) \neq 0\}$ . Hence we must have  $S = \{n \in \mathbb{Z} \mid f(n) = 1\}$  by (a). Since  $f(1) = 0$ , 1 is not in  $S$ . And  $f(0) = 1$  implies that  $0 \in S$ . Take any  $x \in \mathbb{Z}$  and  $y \in S$ . Using (ii), we get

$$f(xy) + f(x) = f(x) + 1.$$

This shows that  $xy \in S$ . If  $x \in \mathbb{Z}$  and  $y \in \mathbb{Z}$  are such that  $xy \in S$ , then (ii) gives

$$1 + f(x)f(y) = f(x) + f(y).$$

Thus  $(f(x) - 1)(f(y) - 1) = 0$ . It follows that  $f(x) = 1$  or  $f(y) = 1$ ; i.e., either  $x \in S$  or  $y \in S$ . We also observe from (ii) that  $x \in S$  and  $y \in S$  implies that  $f(x - y) = 1$  so that  $x - y \in S$ . Thus  $S$  has the properties:

(A)  $x \in \mathbb{Z}$  and  $y \in S$  implies  $xy \in S$ ;

(B)  $x, y \in \mathbb{Z}$  and  $xy \in S$  implies  $x \in S$  or  $y \in S$ ;

(C)  $x, y \in S$  implies  $x - y \in S$ .

Now we know that  $f(10) \neq 0$  and  $f(2) = 0$ . Hence  $f(10) = 1$  and  $10 \in S$ ; and  $2 \notin S$ . Writing  $10 = 2 \times 5$  and using (B), we conclude that  $5 \in S$  and  $f(5) = 1$ . Hence  $f(5k) = 1$  for all  $k \in \mathbb{Z}$  by (A).

Suppose  $f(5k + l) = 1$  for some  $l$ ,  $1 \leq l \leq 4$ . Then  $5k + l \in S$ . Choose  $u \in \mathbb{Z}$  such that  $lu \equiv 1 \pmod{5}$ . We have  $(5k + l)u \in S$  by (A). Moreover,  $lu = 1 + 5m$  for some  $m \in \mathbb{Z}$  and

$$(5k + l)u = 5ku + lu = 5ku + 5m + 1 = 5(ku + m) + 1.$$

This shows that  $5(ku + m) + 1 \in S$ . However, we know that  $5(ku + m) \in S$ . By (C),  $1 \in S$  which is a contradiction. We conclude that  $5k + l \notin S$  for any  $l$ ,  $1 \leq l \leq 4$ . Thus

$$S = \{5k \mid k \in \mathbb{Z}\}.$$

—————00000—————