



ENGINEERING & MANAGEMENT EXAMINATIONS, JUNE - 2009
MATHEMATICS
SEMESTER - 2

Time : 3 Hours]

[Full Marks : 70

GROUP - A**(Multiple Choice Type Questions)**1. Choose the correct alternatives for any *ten* of the following :

10 × 1 = 10

- i) If $A = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$, then A^{100} is

- a) $\begin{bmatrix} 1 & 0 \\ -150 & 1 \end{bmatrix}$ b) $\begin{bmatrix} 1 & 0 \\ -50 & 1 \end{bmatrix}$
 c) $\begin{bmatrix} 1 & 0 \\ -100 & 1 \end{bmatrix}$ d) None of these.

- ii) The set of vectors $\{(2, 1, 1), (1, 2, 2), (1, 1, 1)\}$ in R^3 is

- a) linearly dependent b) linearly independent
 c) basis of R^3 d) none of these.

- iii) The matrix $A = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$ is

- a) an orthogonal matrix b) a symmetric matrix
 c) an idempotent matrix d) a null matrix.

v) The value of the determinant

$$\begin{vmatrix} 1 & 4 & 16 \\ 1^2 & 2^2 & 4^2 \\ 0 & 1 & 6 \end{vmatrix}$$

is

- a) 0
c) 4

- b) 1
d) 22.

v) The solution of a system of n linear equations with n unknowns is unique if and only if

- a) $\det A = 0$
c) $\det A < 0$

- b) $\det A > 0$
d) $\det A \neq 0$.

where A is the matrix of the coefficients of the unknowns in the linear equations

vi) The eigenvalues of the matrix $\begin{bmatrix} 1 & 4 \\ 4 & 1 \end{bmatrix}$ are

- a) -5, -3
c) 3, -5

- b) -5, 3
d) 5, 3.

vii) The general solution of $p = \log (px - y)$ where $p = \frac{dy}{dx}$ is

- a) $y = cx - c$
c) $y = c^2x - e^{-c}$

- b) $y = cx - e^c$
d) none of these.

viii) Which of the following is not true (the notations have their usual meaning)?

- a) $\Delta = E - I$
c) $\frac{\Delta}{\nabla} = \Delta + \nabla$

- b) $\Delta \cdot \nabla = \Delta - \nabla$
d) $\nabla = I - E^{-1}$

ix) $\Delta^2 e^x$ is equal to ($h = 1$)

- a) $(e - 1)^2 e^x$ b) $(e - 1) e^x$
 c) $e^{2x} (e - 1)$ d) e^{2x} .

x) The value of $\int_0^\infty \frac{\sin t}{t} dt$ is equal to

- a) $\frac{\pi}{3}$ b) $\frac{\pi}{6}$
 c) $\frac{\pi}{4}$ d) $\frac{\pi}{2}$.

xi) If S and T are two subspaces of a vector space V , then which one of the following is a subspace of V also?

- a) $S \cup T$ b) $S \cap T$
 c) $S - T$ d) $T - S$.

xii) If $\lambda^3 - 6\lambda^2 + 9\lambda - 4$ is the characteristic equation of a square matrix A , then A^{-1} is equal to

- a) $A^2 - 6A + 9I$ b) $\frac{1}{4} A^2 - \frac{3}{2} A + \frac{9}{4} I$
 c) $\frac{1}{4} A^2 - \frac{3}{2} A + \frac{9}{4}$ d) $A^2 - 6A + 9$.

xiii) Co-factor of -3 in the determinant

$$\begin{vmatrix} -2 & -3 & 4 \\ 1 & 0 & 1 \\ 0 & -1 & 4 \end{vmatrix}$$

- a) 4 b) -4
 c) 0 d) none of these.

**GROUP - B****(Short Answer Type Questions)**

Answer any three of the following.

3 × 5 = 15

2. If A be a skew symmetric and $(I + A)$ be a non-singular matrix, then show that $B = (I - A)(I + A)^{-1}$ is orthogonal.
3. Evaluate $L^{-1} \left\{ \frac{1}{(s-1)^2(s-2)^3} \right\}$.
4. Solve the differential equation

$$\frac{dy}{dx} + y = y^3 (\cos x - \sin x).$$

5. Evaluate the definite integral $\int_1^4 (x + x^3) dx$ by using Trapezoidal rule, taking five ordinates and calculate the error.

6. If $A(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$, then show that

$$A(\theta)A(\phi) = A(\phi), A(\theta) = A(\theta + \phi).$$

GROUP - C**(Long Answer Type Questions)**

Answer any three of the following.

3 × 15 = 45

7. a) If $A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & -4 & 1 \\ 3 & 0 & -3 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 0 \\ 2 & 1 & -1 \end{bmatrix}$, show that $AB = 5I$.

Utilise this result to solve the following system of equations :

$$2x + y + z = 5$$

$$x - y = 0$$

$$2x + y - z = 1$$

2241 (03/06)



- b) Solve : $(y - px)(p - 1) = p$ and obtain the singular solution. Here $p = \frac{dy}{dx}$.
- c) Construct the interpolation polynomial for the function $y = \sin \pi x$, taking the points $x_0 = 0, x_1 = \frac{1}{6}, x_2 = \frac{1}{2}$.

Hence find $f\left(\frac{1}{3}\right)$ where $y = f(x)$.

8. a) Solve the differential equation

$$\frac{d^2y}{dx^2} - 5 \frac{dy}{dx} + 6y = x^2 e^{3x}.$$

- b) Apply suitable interpolation formula to calculate $f(9)$ correct up to two significant figures from the following data :

$x:$	2	4	6	8	10
$f(x):$	5	10	17	29	49

- c) Determine the conditions under which the system of equations

$$x + y + z = 1$$

$$x + 2y - z = b$$

$$5x + 7y + az = b^2$$

admits of

- i) only one solution
- ii) no solution
- iii) many solutions.



9. a) Prove that $P^T A P$ is a symmetric or a skew-symmetric matrix according as A is symmetric or skew-symmetric.

- b) Find the eigenvalues and the eigenvectors of the matrix $\begin{bmatrix} 4 & 6 \\ 2 & 9 \end{bmatrix}$.

- c) Solve by Cramer's rule the following system of equations :

$$3x + y + z = 4$$

$$x - y + 2z = 6$$

$$x + 2y - z = -3.$$

10. a) What is meant by linear independence of a set of n -vectors ?

- b) Solve by the method of variation of parameters the equation

$$\frac{d^2y}{dx^2} + 9y = \sec 3x.$$

c) Prove that $\Delta = \begin{vmatrix} (b+c)^2 & a^2 & a^2 \\ b^2 & (c+a)^2 & b^2 \\ c^2 & c^2 & (a+b)^2 \end{vmatrix} = 2abc(a+b+c)^3$

END