Roll No.

Total No. of Questions: 10]

[Total No. of Pages: 03

28/200

B.Pharmacy (Sem. - 2nd) **ADVANCED MATHEMATICS**

SUBJECT CODE: PHM-1.2.2

Paper ID: [D0108]

[Note: Please fill subject code and paper ID on OMR]

Time: 03 Hours

Maximum Marks: 80

Instruction to Candidates:

- 1) Section - A is Compulsory.
- Attempt any Four questions from Section B. 2)
- Attempt any Three questions from Section C. 3)

Section - A

Q1)

 $(15 \times 2 = 30)$

- Solve $x\sqrt{(1+y^2)} dx + y\sqrt{(1+x^2)} dy = 0$. a)
- Solve $3e^{x}\cos^{2}y \, dx + (1 e^{x})\cot y \, dy = 0$.
- Solve $dy/dx + \frac{1}{x}y = x^n$.
- Solve $\frac{dy}{dx} = \frac{y}{x} + \tan \frac{y}{x}$. d)
- Define the equations reducible to homogeneous equations. e)
- Find $L\left\{\frac{e^{-at}t^{n-1}}{(n-1)!}\right\}$. f)
- Find L $\{(\sin at at \cos at)\}$. g)
- Find L⁻¹ $\left\{ \frac{p}{p^2 + 2} + \frac{6p}{p^2 16} + \frac{3}{p 3} \right\}$. h)
- State second translation theorem for inverse Laplace transforms. **i**)
- Find L⁻¹ $\left\{ \frac{e^{-5p}}{(p-2)^4} \right\}$.

- k) Compare collection of data by census method and sample method in four points.
- The following data relate to the plan outlay of a country for three plans

 Five year plan:

 I IV VII

 Outlay (Rs.'000 crore): 196 2060 8820

 Represent the data by circle diagram.
- m) Draw a pie chart to represent the following data

Age of drivers: <u>under 20 20 - 40 40 - 60 over 60 Total</u>

Percent of totals: 15 60 20 5 100

- n) Write student and paired t test.
- o) Define Binomial distribution.

Section - B

 $(4 \times 5 = 20)$

- **Q2)** Solve $(1 + y^2) dx = (\tan^{-1} y x) dy$.
- Q3) Use convolution theorem to find $L^{-1} \left\{ \frac{p}{(p^2+4)^3} \right\}$.
- Q4) Find the median of the following data:

Income (in Rs.): 100 150 80 200 250 180 No. of Persons: 24 26 16 20 6 30

Q5) The scores of 8 students in an examination in mathematics and statistics are given below.

5 6 7 8 2 3 4 Student No.: 1 52 54 50 60 70 48 58 55 Marks in Maths: 68 51 48 60 55 · 47 53 62 Marks in Stat.:

Find rank correlation coefficient and compare the two values.

Q6) How many tosses of a coin are needed so that the probability of getting at least one head is 0.875.

Section - C

$$(3 \times 10 = 30)$$

Q7) From the following data given below calculate a coefficient of skewness based on percentile.

Marks : less than 10 less than 20 less than 30 less than 40 less than 50

No. of students : 4 10 30 40 47

Q8) Determine the relationship between the semi-inter quartile range and standard distribution in a standard normal probability curve.

Q9) Solve
$$[tD^2 + (1-2t)D - 2]y = 0$$
 if $y(0) = 1$, $y'(0) = 2$.

Q10) Solve
$$(D-2)x - (D+1)y = 6e^{3t}$$

 $(2D-3)x + (D-3)y = 6e^{3t}$
if $x = 3$, $y = 0$ when $t = 0$.

