



One Stop for Colleges  
Education  
Career

- Covers complete CAT syllabus
- Lessons, tests by IIT/IIM toppers
- QA,VA and DI Sections
- 1000+ Questions and Sample tests
- Last minute preparation and help
- Its Free 😊

## Minglebox e-CAT Prep

Free Online Preparation for CAT 2011 with Minglebox e-CAT Prep. Cover basic concepts of "Topic" under "Section" for MBA Entrance Exam Preparation with Study material, solved examples and tests prepared by CAT coaching experts.

### Refresher Material

#### Natural Numbers

1 is the smallest Natural number, 0 is the smallest Whole number, and there is no largest or smallest Integer.

Even numbers are multiples of 2. Any even number can be written as  $2n$ , where  $n$  is an integer. ***0 is an even number***.

Odd numbers are numbers which when divided by 2 leave a remainder of 1. Any odd number can be written as  $2n + 1$ , where  $n$  is an integer.

Factors of a given natural number say  $n$ , is another natural number say  $f$  if  $n$  is completely divisible by  $f$ . Ex factors of 18 are 1, 2, 3, 6, 9, 18 and number of factors is 6.

Highest factor of any natural number is number itself and lowest positive factor is 1. 1 is the factor of every natural number.

The number of factors of any natural number is finite.

## Prime Number

A natural number which has exactly 2 factors is a prime number. Ex number 2 has factors 1 and 2 only. Similarly 3,5,7,11,13,17,19,23,29,31,37,41 etc . **1 is not a prime number.**

A number  $n$  is a prime number if it is not divisible by any prime less than  $[\sqrt{n}]$  where  $[\sqrt{n}]$  is the largest natural number less than or equal to  $\sqrt{n}$ .

Any natural number  $n$  can be written in the form and in one unique way only  $n = p^\alpha \times q^\beta \times r^\gamma \dots$ , where  $p, q, r, \dots$  are different primes and  $\alpha, \beta, \gamma, \dots$  are the powers of the prime number respectively.

### Example

$18 = 2^1 \times 3^2$ . Here 2 and 3 are the prime numbers and 1 and 2 are the respective powers of the primes.

The number of factors of any natural number  $n$ , which can be factored as above is  $= (1+\alpha) \times (1+\beta) \times (1+\gamma) \dots$  etc. Thus number of factors of 18 is  $(1+1) \times (1+2) = 6$ .

The number of odd factors will be given as follows. If the given number does not have any power of 2 then number of factors is same as number of odd factors. But if the number has any term like  $2^\alpha$  in its factorization, as product of prime number powers, where  $\alpha \geq 1$ , then exclude the  $(1+\alpha)$  term and calculate the number of odd factors as  $(1+\beta) \times (1+\gamma) \dots$  Etc. Thus number of odd factors of 18 is  $(1+2) = 3$ , viz... 1, 3 and 9.

**The sum of all the factors is given by the expression  $(p^{\alpha-1} - 1) \times (q^{\beta-1} - 1) \times (r^{\gamma-1} - 1) / ((p-1) \times (q-1) \times (r-1)) \dots$**

Composite number is a number which has more than 2 factors. For example number 18 has 6 factors viz. 1, 2, 3, 6, 9, 18.

## Remainder

Any whole number say  $m$  is divided by another natural number say  $n$  then there exists numbers  $q$  and  $r$  such that  $m = nxq + r$ . Where  $q$  is known as the **quotient** and  $r$  is known as the **remainder**. For any  $m, n \in \mathbb{N}$  (the set of natural numbers)  $q$  and  $r \in \mathbb{W}$  (the set of whole numbers). Thus  $0 \leq r$

## Class of integers

As discussed above the remainder obtained when any number is divided by say 5 then remainder is either 0, 1, 2, 3, 4 only. Therefore any number can be written as either as  $5k, 5k+1, 5k+2, 5k+3, 5k+4$ . Thus entire set of numbers has been split into 5 non overlapping sets.

## HCF

HCF of any given set of numbers is a number which completely divides each number in the given set and the number is highest such number possible.

## LCM

LCM of any given set of numbers is the smallest such number which is divisible by each number of the given set.

For any 2 given numbers  **$\text{HCF} \times \text{LCM} = \text{Product of the 2 numbers}$** .

**HCF of fractions = HCF of numerators of all the given fractions / LCM of the denominators of all the fractions.**

**LCM of fractions = LCM of numerators of all the fractions / HCF of the denominators of all the fractions.**

## Divisibility Rules

If the last digit of a number is even then number is divisible by 2

If the sum of the digits of a number is divisible by 3 then 9 then number is divisible by 3 and 9 respectively.

If the last 2 digits of the number are divisible by 4 then number is divisible by 4 and if last 3 digits is divisible by 8 then number is divisible by 8.

If the last digit of the number is 0 or 5 then number is divisible by 5

If the sum of digits of the number is divisible by 3 and the last digit is even then number is divisible by 6.

A given number is divisible by 7 if the number of tens in the original number - twice the units digits is divisible by 7 ex . to check whether 343 is divisible by 7 or not , Thus twice the units digit is  $2 \times 3 = 6$  and number of tens in the number is 34 .Therefore 343 is divisible by 7 if  $34 - 6$  is divisible by 7 i.e. 28 is divisible by 7.

A number is divisible by 11 if the difference of the sum of digits occurring the even numbered places and the sum of digits occurring in the odd number of places is divisible by 11.

## Surds and Indices

### Rules of Indices

$$b^l \times b^m \times b^n \dots = b^{l+m+p\dots}$$

$$(b^m)^n = b^{mn}$$

If bases are same then powers are also same. i.e . if  $a^m = a^p$  and if  $a \neq 1$  or 0 then it implies  $m = p$ .

The last digit of the square of any number cannot be 2,3,7 or 8.

Any perfect square is of the form either  $4k$  or  $4k-1$  or  $4k+1$ .

Product of any number of even numbers is even and any number of odd numbers is odd.

The product of any  $n$  consecutive natural numbers is divisible by  $n!$ .

### Useful Algebraic identities

$(a^n + b^n)$  is divisible by  $(a+b)$  for all odd values of  $n$ .

$(a^n - b^n)$  is divisible by both  $(a+b)$  and  $(a-b)$  for even values of  $n$ .

$(a^n - b^n)$  is divisible by  $(a-b)$  for all values of  $n$  (both odd and even).

Sum of first  $n$  natural numbers  $= n(n+1)/2$ .

Sum of squares of first  $n$  natural numbers  $= n(n+1)(2n+1)/6$

Sum of the cubes of first  $n$  natural numbers is given by  $\{n \times (n+1)/2\}^2$ .

### Units digit of any power of a number (Cyclicity)

If we consider the units digit of the powers of 2 i.e.  $2^x$  for different values of  $x$  then we find the unit's digit is 2,4,6,8,2,4,6,8,2,4,6,8.... for  $x= 1,2,3,4,5,6,7,8,9,10,11,12,...$ . Similar patterns exist for the unit's digit of other numbers. The results are summarized in the table below.

## Solved Examples

	Unit's digit of $a^x$ , $k$ is an natural number			
Number "a"	$x=4k+1$	$x=4k+2$	$x=4k+3$	$x=4k$
1	1	1	1	1
2	2	4	6	8
3	3	9	7	1
4	4	6	4	6
5	5	5	5	5
6	6	6	6	6
7	7	9	3	1
8	8	4	2	6
9	9	1	9	1

### QUESTION

1) The Rightmost non zero digit of  $5670^{5670}$  is

- (a) 7      (b) 1      (c) 3      (d) 9

### SOLUTION

Here we use rule of cyclicity to solve this problem. The first non zero digit to the left of 0 is 7 hence the rightmost non zero digit will be same as the units digit of  $7^{5670}$ . Now next we need to find out the form of 5670. Dividing 5670 by 4 we get 2 as remainder. Hence 5670 is of the form  $4k + 2$ . Therefore the unit's digit is 9. Therefore answer is d

2) The numbers 13409 and 16760 when divided by a 4 digit integer n leave the same remainder then the value of n is

- (a) 1127      (b) 1117      (c) 1357      (d) 1547

### SOLUTION

Here 13409 and 16760 on division by n leave the same remainder hence can be written as  $nxm + r$  and can be written  $nxp + r$ . Therefore subtracting the 2 equations we get  $nx(m-n) = 3351 = 1117 \times 3$  But n is 4 digit number hence n is 1117. Hence answer is b.

### Refresher Material

#### Percentages:

A student gets 20 marks out of 30 in an examination, and then we express this information as follows. If the student got 20 marks when the total marks were 30 what would he have got if the total marks were 100?

The amount to this question is = 66.66%.

The person of finding the answer to the question what is the value if total in 100 is what leads is to percentage. This if we say student scored 66.66 percent it means if total is 100 then student got 66.66 marks.

#### Equivalence between fraction and percentage

Fraction	Equivalence percentage
1	100%
1/2	50%
1/3	33.33%
1/4	25%
1/5	20%
1/6	16.67%
1/7	14.28%
1/8	12.5%
1/9	11.11%
1/11	9.09%

## Percentage Change

If a quantity A is increased by 'x' then percentage increase =

$$\frac{\text{change}}{\text{original}} \times 100 = \frac{(a-x)-a}{a} \times 100$$

$$= \frac{100x}{A} \%$$

If a quantity is increased by a% then new value is given by  $(1 + \frac{a}{100}) \times \text{original quantity}$  similar if a quantity is decreased by a% then new

Value is given by  $(1 - \frac{a}{100}) \times \text{original quantity}$ .

If a quantity is increased by a% and then further by b% these percent change given are equivalent to a single percent change given by

$$(a+b+\frac{ab}{100}) \%$$

Similarly if there is a successive decrease of a% followed by b% then effective percentage decrease is

$$(a+b-\frac{ab}{100})$$

## Selling Price, Marked Price and Cost Price

### Selling price

(SP) is the price at which an article is sold cost price is the price at which an article is bought. If **SP > CP** then the difference **SP – CP** is known as the profit, **P P = SP – CP**

and if **SP < CP** then the difference **CP – SP** is known as loss, **L L = CP – SP**

$$\text{Profit Percentage } P\% = \frac{\text{profit}}{\text{CP}} \times 100$$

$$\text{Loss Percentage } L\% = \frac{\text{loss}}{\text{CP}} \times 100$$

### Marked price.

The price which is displayed on the tag of the article is known as marked price.

Generally the SP is less than the marked price (MP) the difference **MP – SP** is known as discount D

$$\text{Discount \%, } D\% = \frac{\text{Discount}}{\text{MP}} \times 100$$

## Solved Example

### Question

A shopkeeper sells sugar for Rs 25 per kg and makes a profit of 20%. If he sells the sugar for Rs 22.50 per Kg find his profit?

### Solution

Let the cost price CP of sugar be Rs x per Kg then **SP = CP + Profit**

$$\text{But Profit} = \frac{\text{Profit} \times \text{CP}}{100} = \frac{20}{100} \times \text{CP}$$

$$\text{SP} = \text{CP} \left(1 + \frac{20}{100}\right)$$

$$25 = \text{CP} (1.2)$$

$$\text{CP} = \frac{25}{1.2} = 20.83 \text{ Rs/kg}$$

Hence profit when SP is 22.50 Rs/ Kg

$$= 1.67 \text{ Rs/Kg}$$

$$\text{Profit \%} = \frac{1.67}{20.83} \times 100 = 8 \%$$

### Question

A man sells 2 pens Rs 100 each at a profit of 10% and another at a loss of 10% find his overall profit or loss percentage.

### Solution

If the cost price of the 2 pens be x and y respectively then using profit %  $\frac{SP-CP}{CP} \times 100$  we get

$$CP = \frac{100 \times P}{100 + P} \text{ in case of profit}$$

$$\text{And } CP = \frac{100 \times L}{100 - L} \text{ in case of loss}$$

$$\text{Hence } x = \frac{100 \times 10}{100 + 10} \text{ and } y = \frac{100 \times 10}{100 - 10}$$

$$x = \frac{100}{11} \text{ and } y = \frac{100}{9}$$

$$\text{Total CP of both pens} = \frac{100}{11} + \frac{100}{9} = \frac{2000}{99}$$

$$\text{Total loss} = \frac{2000}{99} - 200 = \frac{2000 - 1980}{99}$$

$$= \frac{20}{99}$$

$$\text{Loss \%} = \frac{\frac{20}{99}}{\frac{2000}{99}} \times 100$$

$$= 1\%$$

**Note:** In case 2 article are sold at same selling price one at a profit of a% another are at loss of a% then there is overall loss on the whole out lay and loss percentage is given by  $\frac{a^2}{100} \%$

### Refresher Material

#### Averages

In arithmetic mean or the average of  $n$  quantities  $x_1 + x_2 + x_3 + \dots + x_n$  is given by

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n}$$

The average is  $\geq$  the smallest of  $x_1 + x_2 + \dots + x_n$

The average is  $\leq$  the greater of  $x_1 + x_2 + \dots + x_n$

The Sum of deviation of each element  $\beta$  with respect to AM is equal to zero i.e

$$(x_1 - \bar{x}) + (x_2 - \bar{x}) + \dots + (x_n - \bar{x}) = 0$$

#### Rules of Allegation

A shopkeeper mixes two varieties of sugar costing Rs 25/Kg and Rs 20/Kg in a certain ratio such that the cost of the mixture is Rs 23/Kg then find the ratio in which the 2 types of sugar were mixed?

Let the ratio in which sugar of Rs 25/Kg and Rs 20/Kg be  $p:q$  then total cost of mixture =  $\frac{25p + 20q}{p + q} = 23$

Solving for p and q we get  $\frac{p}{q} = \frac{23-20}{25-23}$

This is the allegation rule.

**Formally it states the ratio of dear quantity and the cheap quantity is equal to the ratio of the difference of the mean price and the cheaper price and the difference of the dear price and the mean**

Price symbolically  $\frac{p}{q} = \frac{Mp - Cp}{Dp - Mp}$  where cp- cheaper price

Mp – mean price, Dp – Dear price

## Mixtures

When we mix two or more than 2 pure substance, we get what is known as a mixture.

**Mixture of Mixtures:** - If we mix two mixtures which have components say A and B in the final mixture is given by

$$\frac{a}{b} = A$$

The weight of first and second mixture respectively.

A special problem on mixtures

If a contains 'x' liter of liquid 'x' and if 'y' liters is taken out and replaced by 'y' liters of liquid 'z' and if the above step is repeated with the mixture so obtained that is x liter of mixture replaced with y liter of liquid 'z' then if the above operation is repeated 'n' times then

$$\frac{\text{liquid X left the container after } n^{\text{th}} \text{ operation}}{\text{Initial quantity of liquid X the container contains}} = \left(\frac{x-y}{x}\right)^n$$

$$\frac{\text{Amount Liquid Z the solution after } n^{\text{th}} \text{ operation}}{\text{initial quantity of liquid X the container contains}} = 1 - \left(\frac{x-y}{x}\right)^n$$

**Note:** The final volume of mixture remains contents and is same as initial volume same as initial of liquid X

## Solved Examples

---

### Question

The average age of 10 students in a class is increased by 2 year when two students aged 12 year and 14 year are replaced by 2 girls. Find the average age of the two girls.

### Solution

Let the sum of the ages of the 2 girls be  $y$  years, also sum of the ages of the 2 students who have replaced = 26 years average age of group has gone up by 2 years that means the increase in total age of the 10 students is 20 years this increase is due to the age of the 2 girls hence  $y - 26 = 20$ ,  $y = 46$  therefore average age of the 2 girls.

### Question

There are 60 students in a class. These students are divided into three groups A, B and C of 15, 20 and 25 students each. These groups A and C are combined to form group D. What is the average weight of the students in group D?

- (1). More than the average weight of A
- (2). More than the average weight of C
- (3) Less than the average weight of C

(4) Cannot be determined.

### **Solution**

Number of students in group D is more than number of students in group A or group C. But there is no information about the weight of students in group A or group C. Hence answer is (4)

**Note:** As a group D has students from group C whichever group has higher average weight, the average weight of group D will be that group's average weight.

### Refresher Material

#### Work

If a person A can do a piece of work in 'a' days, then working at the same uniform speed A will do  $\frac{1}{a}$  fraction of the work in one day.

For example,

Days taken to complete 'a' work	25	12.5	1/2
Fraction of work done in a day	$1/25^{\text{th}}$	$1/12.5^{\text{th}}$	2

Working together – If working alone can do some work in 'a' days and if another person B working alone can do it in 'b' days, then if A and B

**Start working together the amount of work done by A & B in one day =  $\frac{1}{a} + \frac{1}{b}$**

**Therefore total work is done by A and B working together in  $\frac{ab}{a+b}$  days**

**Note:** Remember it is work done that is additive and not number of days.

## LCM technique

If A completes a work in 'a' days working alone and B completes it in 'b' days working alone, then instead of assuming the work as 1 unit we can assume total work to be LCM of (a,b) this will avoid the fraction in the calculation.

## Proportionality (concept of efficiency)

If A is thrice as efficient as B then if A takes 12 days to complete a work then B will take  $12 \times 3 = 36$  days to complete the work. Whenever relative efficiency of team member is given the most suitable approach is to consider the capacities of each team member in terms of any given team member.

Amount of Work done is directly proportional number of person employed to do the work and days worked.

If amount of work is constant then number of days taken to complete the work is inversely proportional to days taken.

Men, Women and children working in a given project.

### Question

If work can be done in 10 days by 4 men or 6 women or 10 children. How many days are required for 3 men, 8 women and 6 days to complete it?

### Solution

1 Man will take 40 days; hence 3 men will take  $40/3$  days to complete it

1 woman will take 60 days; hence 8 women will take  $60/8$  days

1 child will take 100 days; hence 6 days will take  $100/6$  days

then total time taken if they all worked together is =  $\frac{1}{\frac{3}{40} + \frac{8}{60} + \frac{6}{100}} = \frac{600}{45+80+36} = \frac{600}{161}$  days.

### Question

A house can be built by A in 50 days and B can demolish it in 60 days. If A and B work on alternative days, in how many days will the house be built, assuming A starts first and once house is built completely B does not demolish it?

### Solution

In 2 days cycle amount of work done =  $\frac{1}{50} - \frac{1}{60} = \frac{1}{300}$  but does that mean work gets done in 600 days. Think again as a finisher  $\frac{1}{50}$  th of

the work in one days , hence if work done is less than  $\frac{1}{50}$  then A finishes the work and B does not demolish it .

In two days amount of work done is  $\frac{1}{300}$  th therefore  $\frac{49}{50}$  th fraction of the work is finished in

588 days work left to be done on 589<sup>th</sup> days is  $\frac{1}{50}$  which A will finish. Hence work gets finished in 589 days.

### Refresher Material

#### Speed

Speed is the rate at which distance is covered by a moving body these speed

$$s = \frac{\text{distance}(d)}{\text{time taken}(t)} \text{ or}$$

$$s = \frac{d}{t}$$

if speed is constant then  $d \propto t$

if distance is constant then  $s \propto \frac{1}{t}$

if time is constant then  $d \propto s$

#### Average speed

If a body cover half of the distance 'd' it a speed of  $s_1$  and the half at speed  $s_2$  then average speed is given by

$$\frac{\text{total distance covered}}{\text{total time taken}} = \frac{\frac{2d}{\frac{s_1}{d} + \frac{s_2}{d}}}{\frac{2d}{\frac{s_1}{d} + \frac{s_2}{d}}} = \frac{2s_1 s_2}{s_1 + s_2}$$

### Circular races

If 2 persons say a,b start running a race on a circular track, then the faster runner (say A) will be together with the slower runner for the first time when faster runner has gained one complete length of the track or round over the slower runner.

### Relative speed

The time taken by a train of length 'l' moving with speed 's' to pass a pole is =  $\frac{l}{s}$

Time taken by train of length 'l' moving with speed 's' to completely pass a platform of length 'd' =  $\frac{l+d}{s}$

Time taken by train of length 'l' moving with speed  $s_1$  to pass another train of length 'd' moving with speed  $s_2$  in the direction (provided  $s_1 > s_2$ ) =  $\frac{l+d}{s_1 - s_2}$

Time taken by a train, of length l, and moving with speed  $s_1$  to pass another train of length 'd' moving with  $s_2$  in the opposite direction  $\frac{l+d}{s_1 + s_2}$

If person covers a certain distance 'd' moving speed  $s_1$  and taking time  $t_1$ , then the person covers the same distance moving at speed  $s_2$  and taking time  $t_2$  then  $\frac{s_1}{s_2} = \frac{t_1}{t_2}$

If two persons X and Y start at the same time in opposite direction from two places and arrive at their respective destination in x and y hours later after having met on their way then  $\frac{\text{speed of x}}{\text{speed of y}} = \frac{\sqrt{y}}{\sqrt{x}}$

If speed of boat, in still water is 'a' (m/s) then speed of boat in going down stream in a river which is following at 'b' (m/s) is (a+b) m/s. Hence if a boat cover a certain distance 'd' downstream and then reruns it to its starting point covering the same distance 'd' then  $\frac{\text{upstream journey time}}{\text{downstream journey time}} = \frac{a+b}{a-b}$

## Solved Examples

### Question

2 boats A and B start from point P at the same constant speed of 15 km/hr in still water. Boat A goes to Q, where boat B goes till R, which is equidistant from P and Q and returns to P. If the time taken by boat A is 1.5 times that of B, what is the speed of the current?

1. 10 k mph
2. 7.5 k mph,
3. 5k mph
4. Data insufficient

### Solution

If the total distance between P and Q be '2D' km. Boat A has taken more time, the direction of the current must be from Q to P if the speed of the current must be X k mph then

$$\text{Time taken by A} = \frac{2d}{15-x}$$

$$\text{Time taken by B} = \frac{d}{15+x} + \frac{d}{15-x}$$

Given  $\frac{2d}{15-x} = \frac{3}{2} \left( \frac{d}{15+x} + \frac{d}{15-x} \right)$

$$\frac{4}{3} = \frac{30}{15+x}$$

Solving we get  $x = 7.5$  k mph have option 2 is the right option.

### Question

A car covers half the half distance at speed 60kmph and another half of the distance at speed 30kmph. Find the average speed of the car for the whole journey?

### Solution

If total distance be '2d' then time taken to cover the first half of the journey, =

$$t_1 = \frac{d}{60} \left( \frac{\text{distance}}{\text{speed}} \right)$$

and time taken to cover the second half of the journey =  $\frac{d}{30}$

Therefore total time taken =  $t_1 + t_2 = \frac{d}{60} + \frac{d}{30}$

$$\text{Avg .Speed} = \frac{\text{total distance}}{\text{total time taken}} = \frac{2d}{\frac{d}{60} + \frac{d}{30}} = \frac{2}{\frac{1}{60} + \frac{1}{30}} = 40 \text{ kmph}$$

### Refresher Material

Linear equation in one variable A linear equation in one variable is respected as  $ax + b = 0$ , where  $a, b$  are real constants and  $a \neq 0$ .  $a$  is known as the coefficient of 'x' variable 'x' solution of linear equation in one variable is  $x = -\frac{b}{a}$  any linear equation in one variable always has a solution.

### Linear equation in 2 variables

Linear equation in 2 variable – A linear equation in 2 variable is represented as  $ax + by = c$  where  $a, b, c$  are real constants, where either  $a$  or  $b \neq 0$  but not both simultaneously equation  $ax + by = c$  has infinite number of solution for real  $x$  and  $y$  but for suitable restrictions equation may have finite number of solutions.

### Simultaneous linear equations

Simultaneously linear equations in two variables

A simultaneously linear equation in 2 variables is represented as  $a_1x + b_1y = c_1$

and  $a_2x + b_2y = c_2$  (2) where  $a_1, b_1, a_2, b_2, c_1, c_2$  are real constants the above pair of equations may have none, one or infinite number of solutions depending on the relationship between the constants.

If  $\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{c_1}{c_2}$  then we have infinite number of solutions.

If  $\frac{a_1}{b_1} \neq \frac{a_2}{b_2} \neq \frac{c_1}{c_2}$  then there is no solution

If  $\frac{a_1}{b_1} = \frac{a_2}{b_2}$  then there is one unique solution

### Equation in one variable

Equation in one variable but of higher degree. The degree of equation is the highest power of variable that exists in the equation for example

$x^3 - 7x + 2 = 0$  is an equation of degree 3 in one variable .

$x^2 - 3x + 2 = 0$  is an equation of degree 2 in one variable .

An equation of second degree in one variable is known as quadratic equation.

### Quadratic equation

A equation of the form  $ax^2 + bx + c = 0$  where a,b,c are real numbers and ,  $a \neq 0$  is known as a quadratic equation .

Roots of a quadratic equation A value of 'x' say which makes the left hand side expression equal to right hand side that is '0' .

Therefore if there exists a number  $\alpha$  such that  $a\alpha^2 + b\alpha + c = 0$  , then ' $\alpha$ ' is said to be a root of the equation . In general, any quadratic will have roots either real or imaginary.

Relationship between roots of the equation and co-efficients of the equation.

If  $\alpha, \beta$  are the roots of the equation and then  $\alpha + \beta$  (sum of the roots) =  $\frac{-b}{a} \left( \frac{-\text{coefficient of } x}{\text{coefficient of } x^2} \right)$   
 and  $\alpha\beta$  (product of roots) =  $\frac{c}{a} \left( \frac{\text{constant of } x}{\text{coefficient of } x^2} \right)$  method to solve a quadratic equation.

Completing the square method at  $ax^2 + bx + c = 0$  be the given equation.

Step1: Add and subtract the quantity  $\left(\frac{b}{2a}\right)^2$  that is  $\left(\frac{\frac{1}{2} \times \text{coeff of } x}{\text{coeff of } x^2}\right)^2$

Step 2: Rearrange the terms as follows

## Sample Examples

$$a\left(x^2 + 2 \cdot \frac{b}{a} \cdot x + \left(\frac{b}{2a}\right)^2\right) + c \frac{-b^2}{4a^2} = 0$$

$$\left(x + \frac{b}{2a}\right) + \frac{4ac - b^2}{4a^2} = 0$$

$$x + \frac{b}{2a} = \frac{\pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Discriminant the expression  $b^2 - 4ac$  is known as the discrimination of the quadratic. Just by finding the discriminant we can decide the nature of the root

$D > 0$	Roots real and distinct
$D = 0$	Roots real and equal
$D < 0$	Roots are imaginary
$D > 0$ and a perfect square	Roots are distinct real and Rational

Factorization method before attempting this method, check the discriminant of the equation. If the discriminant D is zero or a perfect square, then this method will easily give the roots of the equation. We illustrated this by an example.

### Question

Find the roots of  $x^2 - 5x + 6 = 0$

### Solution

#### Step i :-

Find the discriminant of the equation here by comparison we get  $a = 1$ ,  $b = -5$ , and  $c = 6$  the discriminant  $D = b^2 - 4ac = (-5)^2 - 4 \cdot 1 \cdot 6 = 25 - 24 = 1$  which is a perfect square

#### Step ii :-

Consider the constant term C, and factorize it into 2 factors whose product is C. here

$$C = 6$$

Therefore 2 factor products of 6 are  $1 \times 6, -1 \times -6, 2 \times 3, -2 \times -3$

For each pair try to see in which case the sum of factors is equal to b (the coefficient of x) here  $b = -5$ , we see that  $-2 \times -3$  satisfies this as  $-5 = (-2) + (-3)$

#### Step iii :-

Write the coefficient of x as sum of the 2 factors obtained in step

So  $x^2 - 5x + 6 = x^2 - 2x - 3x + 6 = 0$

take out the common terms for first 2 terms and do the same for last 2 terms

$x(x-2) - 3(x-2)$  (x-2) is common in both the terms, so pull it out we get  $(x-2)(x-3) = 0$

**Step iv :-**

Equation each term equal to '0' to get the root

$$x-2=0 \text{ or } x-3=0$$

$$x=2 \text{ or } x=3$$

Hence roots of  $x^2 - 5x + 6 = 0$  are  $x=2$  and  $x=3$  respectively

### Refresher Material

#### Ratio

Ratio is a tool for comparing 2 quantities. For example in a clear if there are 20 girls and 30 boys then we can compare the number of boys and girls as follow:

$$\frac{\text{Number of girls}}{\text{Number of boys}} = \frac{20}{30} = \frac{2}{3}$$

which is offer stated as

Number of girls: Number of boys = 2 : 3

$$\text{Number of girls} = \frac{2}{3} \times \text{Number of boys}$$

$$\text{Number of boys} = \frac{3}{2} \times \text{number of girls}$$

Ratio is a pure number it does not have a unit. The 2 numbers used in ratio are known as the '**terms**' of the ratio. The first term is known as '**antecedent**' and the second term is known as '**consequent**'.

If the ratio of 2 or more quantities is given by a:b:c

then actual quantity is given by ak, bk, ck

where k is constant.

## Proportion

Equality 2 ratio is known as proportionality that is if  $\frac{a}{b} = \frac{c}{d}$  then a,b,c,d are in proportion similarly if  $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = \dots\dots\dots$  etc .a, b, c, d, e, f etc are in proportion.

If  $\frac{a}{b} = \frac{c}{d}$  then

$$ad = bc$$

$$\frac{b}{a} = \frac{d}{c}$$

$$\frac{a}{c} = \frac{b}{d}$$

$$\frac{a+b}{b} = \frac{c+d}{d}$$

**Question**

If  $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = \frac{2}{3}$  then the value of  $\frac{2a^2 - 3c^2 + 4e^2}{2b^2 - 3d^2 + 4f^2}$  is

**Solution**

$$\text{As } \frac{a}{b} = \frac{c}{d} = \frac{e}{f} = \frac{2}{3}$$

$$a = \frac{2}{3}b$$

$$c = \frac{2}{3}d$$

$$e = \frac{2}{3}f \text{ putting values of } a, c, e \text{ in the expression}$$

$$\frac{2a^2 - 3c^2 + 4e^2}{2b^2 - 3d^2 + 4f^2} = \frac{2\left(\frac{2}{3}b\right)^2 - 3\left(\frac{2}{3}d\right)^2 + 4\left(\frac{2}{3}f\right)^2}{2b^2 - 3d^2 + 4f^2} = \left(\frac{2}{3}\right)^2 = \frac{4}{9}$$

### Question

The ratio of the current ages of Anita and Sunita is 4:3. 8 years hence, the ratio of their ages will be 6:5. then their current ages are

### Solution

Let this current age of Anita and Sunita be  $4k$  and  $3k$  then their ages 8 years hence will be  $4k+8$ ,  $3k+8$  hence of their ages 8 years

hence will be  $\frac{4k+8}{3k+8} = \frac{6}{5}$

Solving we get  $20k + 40 = 18k + 48$

$$2k = 8$$

$$k = 4$$

Hence their current ages are 16 and 12 years respectively.

### Refresher Material

A comparison relationship between two algebraic expression or quantities is known as an Inequalities.

#### For example

$$3x+7 > x^2+1$$

$$x^2+3x+2 \geq 13x+9 \text{ etc.}$$

In any inequality problem variable  $x$  is assumed to be real number unless otherwise. Any inequality is generally either one of the following

'greater then' ( $>$ )

'greater than or equal to' ( $\geq$ )

'less than' ( $<$ )

'less than or equal to' ( $\leq$ )

### Solution of an inequality

By solution of an inequalities we seek to find a set of value for the variable involve in the problem so that inequalities holds true for all the values lying in the solution set.

## An Example

### Question

$$3x+7 > x+1$$

### Solution

We find that any value of  $x > -3$  is a solution of the above inequalities. For instance

let us take  $x = -2, -1, 0, .5$  etc. We find

$$3(-2)+7 > -2+1 \Rightarrow 1 > -1 \text{ hence true}$$

$$3(-1)+7 > -1+1 \Rightarrow 4 > 0 \text{ hence true}$$

$$3(0)+7 > 0+1 \Rightarrow 7 > 0 \text{ hence true}$$

## Law of inequalities

If  $x > y$  then for all  $c$   $x+c > y+c$  and  $x-c > y-c$

If  $x > y$  and  $c > 0$  then  $cx > cy$  and  $\frac{x}{c} > \frac{y}{c}$

If  $x > y$  then for  $c < 0$ ,  $xc < yc$  and  $\frac{x}{c} < \frac{y}{c}$  that is sign of inequalities gets reversed when both sides of

the inequalities are multiplied or divided by the same negative quantity. If  $x > y > 0$  then

$$\frac{1}{x} < \frac{1}{y}$$

$$8 > 2 > 0$$

$$\frac{1}{8} < \frac{1}{2}$$

also if  $0 > x > y$  then  $\frac{1}{x} < \frac{1}{y}$

but  $x > 0 > y$  then  $\frac{1}{x} > \frac{1}{y}$

## Relationship between Powers of a Number

If  $x > 1$  then .....  $x^3 > x^2 > x > \frac{1}{x} > \frac{1}{x^2}$  .....

but if  $0 < 1$  then .....  $x^3 < x^2 < x < \frac{1}{x} < \frac{1}{x^2} < \frac{1}{x^3}$  .....

In general if  $x > 1$  then  $x^m > x^n$  if  $m > n$  and if  $0 < x < 1$  then  $x^m < x^n$  if  $m < n$

## Solving quadratic inequalities

Let us study how the sign of a quadratic expression changes as we vary the value of  $x$ .

let us consider  $x^2 - 3x + 2$  that is

$(x-2)(x-1)$  for  $x > 2$  we find  $x-2 > 0$  and  $x-1 > 0$  therefore for all  $x > 2$   $(x-2)(x-1) > 0$

for  $1 < x < 2$   $x-2 < 0$  but  $x-1 > 0 \Rightarrow (x-2)(x-1) < 0$

where for  $x < 1$  then  $x-2 < 0$  and  $x-1 < 0 \Rightarrow (x-2)(x-1) < 0$

thus if we have to solve  $x^2 - 3x + 2 > 0$  we proceed as follows.

- factorize the given expression linear factors.
- Equate each linear term equal to zero and find values of  $x$ .
- Arrange the value of  $x$  in directly order, so if values are  $\alpha_1$  and  $\alpha_2$  and if  $\alpha_1 < \alpha_2$  then for all  $x > \alpha_2$  expression is  $> 0$  for all  $\alpha_1 < x < \alpha_2$  then expression is  $< 0$  and for all  $x < \alpha_1$  expression  $> 0$ .

While factorizing the expression in linear factor make sure to make the sign of coefficient of  $x^2$  as positive.

$$\left(x + \frac{b}{a}\right) + \frac{4ac - b^2}{4a^2} = 0$$

$$x + \frac{b}{2a} = \frac{\pm \sqrt{b^2 - 4ac}}{2a}$$

### Question

**Solve**  $(x-1)^3(x+1)^2(x-3) < 0$

### Solution

Observe for all  $x \in \mathbb{R}$  and for  $x+1 \neq 0$ ,  $(x+1)^2 > 0$ . Therefore  $(x+1)^2$  is the positive factor of the above expression hence we can divided both sides by  $(x+1)^2$  we get  $(x-1)^3(x-3) < 0$  also  $(x-1)^3 = (x-1)^2 \cdot (x-1)$  we again observe for all  $x \in \mathbb{R}$  and for  $(x-1) \neq 0$   $(x-1)^2 > 0$

Hence we just need to consider  $(x-1)(x-3)$  for solving inequalities now let's equal each linear term equal to zero we get  $x = 1$  and  $x = 3$  therefore for all  $x > 3$   $(x-1) > 0$  and  $(x-3) > 0$  hence  $(x-1)(x-3) > 0$  also for  $x < 1$ ,  $(x-1) < 0$  and  $(x-3) < 0$  hence again  $(x-1)(x-3) > 0$  therefore solution set is given by  $x \in (-\infty, 1) \cup (3, \infty)$

### Question

**Solve**  $x^2 - |x| + 2 < 0$

### Solution

We know that  $x^2 > 0$  for all  $x > 0$

$$x^2 = |x|^2$$

Hence  $x^2 - 3|x| + 2 < 0$  become  $|x|^2 - 3|x| + 2 < 0$  factorizing into linear factor we get

$$(|x| - 2)(|x| - 1) < 0$$

Equality each linear factor equal to zero, we get

$$|x| = 2 \text{ and } |x| = 1$$

Hence if  $1 < |x| < 2$  then  $(|x| - 2)(|x| - 1) < 0$

but  $|x| = x$  for  $x > 0$  and

$= -x$  for  $-x < 0$

above solution is  $1 < x < 2$  or  $1 < -x < -2$  or  $-1 > x > -2$  hence following we get solution is

$$x \in (-2, -1) \cup (1, 2)$$

### Refresher Material

#### Parallel Lines

Two lines lying in a plane are said to be parallel if they never meet.

#### Polygons


A closed figure, bounded by finite number of line segments, all lying in the same plane, is known as polygon.

Sum of interior angles of a polygon having 'n' sides, =  $(2n-4)90^\circ$

Sum of exterior angles of a polygon =  $360^\circ$  (irrespective of number of sides)

## Triangles

A triangle is a polygon formed by 3 line segments joined end to end.



Sum of interior angles of a triangle = 180

In any given triangle sum of 2 interior angles is equal to the third remote angle.


AD is the median of triangle, then  $AB^2 + AC^2 = 2(AD^2 + DC^2)$

G is the centroid of the triangle. Centroid of a triangle divides the median AD in the ratio 2:1

## Angle Bisector Theorem


If AD is the bisector of the angle ABC then AD divided the side BC in the ratio

$$\frac{AB}{AC} \text{ ie } \frac{AB}{AC} = \frac{BD}{CD}$$



## Circle Theorem


If AC is a chord of circle then angle ABC subtended by the chord at a point B on the circle is equal to



to angle subtended at point O lying in the same segment ie  $\angle ABC = \angle ADC$ . Also angle ABC is half of angle AOC.

## Alternate Segment Theorem

If XT is a tangent to the circle then  $\angle ATB = \angle ACT$ .



$$\begin{aligned}\text{Area of a triangle} &= \frac{1}{2} \times \text{base} \times \text{Height} \\ &= \sqrt{s(s-a)(s-b)(s-c)} \\ \text{where } s &= \frac{a+b+c}{2}\end{aligned}$$

$$\text{Area of a circle of radius } r = \pi r^2$$

$$\text{Area of a square of side } a = a^2$$

$$\text{Length of diagonal of square} = \sqrt{2a}$$

**Area of rectangle of length l and breath b= lb**

**Volume of cube of edge length  $a=a^3$**

**Length of cube =  $\sqrt[3]{a}$**

**Volume of sphere of radius r =  $\frac{4}{3}\pi r^3$**

**Volume of hemisphere of radius r =  $\frac{2}{3}\pi r^3$**

**Surface area of cube =  $6a^2$**

**Surface area of sphere =  $4\pi r^2$**

**Surface area of hemisphere =  $2\pi r^2$**

### QUESTION

The sum of the areas of two circles, which touch each other externally, is  $153\pi$ . If the sum of their radii is 15, find the ratio of the larger to the smaller radius.

- (1) 4
- (2) 2
- (3) 3
- (4) none of these.

### SOLUTION

Let the radii of the 2 circles be  $r_1$  and  $r_2$ , then  $r_1 + r_2 = 15$  (given) and  $\pi r_1^2 + \pi r_2^2 = 153\pi$  (given)

$$r_1^2 + r_2^2 = 153$$


$$r_1^2 + (15 - r_1)^2 = 153$$

solving we get,  $r_1 = 12$  and  $r_2 = 3$

ratio of the larger radius to the smaller one is  $12 : 3 = 4 : 1$  hence option (1) is the answer.

## QUESTION

In the adjoining figure, points A, B, C and D lie on the circle.  $AD = 24$  and  $BC = 12$ . what is the ratio of the area of  $\triangle CBE$  to that of the triangle  $\triangle ADE$  ?



- (1) 1:4
- (2) 1:2
- (3) 1:3
- (4) Data insufficient.

## SOLUTION

In  $\triangle CBE$  and  $\triangle ADE$ ,  $\angle CBA = \angle CDA$ .

(a chord of a circle subtends equal angle at all points on the circumference, lying in the same segment)

similarly  $\angle BCD = \angle BAD$  and  $\angle BEC = \angle AED$

Therefore  $\triangle CBE = \triangle ADE$  ( AAA similarity rule)


Now  $\frac{BC}{DA} = \frac{12}{24} = \frac{1}{2}$

Hence  $\frac{BE}{AE} = \frac{CE}{AE} = \frac{1}{2}$  hence option b).

### Refresher Material


#### Coordinate Geometry

In two dimensional Coordinate Geometry, location of any point lying in the plane, is given by specifying the perpendicular distances of the point, from a set of fixed mutually perpendicular lines. The fixed mutually perpendicular lines are known as X-axes and Y-axes respectively. The point of intersection is known as the origin 'O'.



These 2 lines divide the given plane in 4 parts known as quadrants. Distances measured to the right hand side of origin O are treated as positive and distances measured towards the left of origin are treated as negative. In a similar way distances measured along the Y-axis and above the X-axis are treated as positive distance measured below the X-axis are treated as positive distances measured below the X-axis are treated as negative.

The Coordinates of a point are specified as an ordered pair, Comprising of its distance measured along the X and Y axis. Distance measured along one X-axis from the origin is known as abscissa, and usually denoted X, the distance measured along the Y-axis from the origin is called the ordinate of the point and is denoted by y. Thus coordinate of a point are specified as (x,y) i.e as (abscissa, ordinate). The 4 quadrants are known as 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> & 4<sup>th</sup> quadrant. The below diagram summarizes the sign of the abscissa and the ordinate of points lying in 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup> quadrant.




### Distance Formula

The distance d between 2 points  $P_1$  and  $P_2$  having coordinates

$(X_1, Y_1)$  and  $(X_2, Y_2)$  is given by

$$D = \sqrt{(X_2 - X_1)^2 + (Y_2 - Y_1)^2}$$

## Section formula and mid point formula



The coordinates  $P(x, y)$  of a point which divides the join of points  $P_1(x_1, y_1)$  and  $P_2(x_2, y_2)$  in the

ratio  $m:n$  is given by  $X = \frac{mx_2 + nx_1}{m+n}$  ;  $Y = \frac{my_2 + ny_1}{m+n}$

### Mid point Formula


The Coordinates of the mid point of the line joining  $P_1(x_1, y_1)$  and  $P_2(x_2, y_2)$  is given by  $X = \frac{x_1 + x_2}{2}$  ;  $Y = \frac{y_1 + y_2}{2}$

### Coordinates of the centroid of a triangle

Let  $P_1(x_1, y_1)$  ,  $P_2(x_2, y_2)$  and  $P_3(x_3, y_3)$  be the coordinates

of the vertices of triangle  $P_1P_2P_3$  , then coordinates of it's centroid  $(x, y)$  is given by  $X =$

$$\frac{x_1 + x_2 + x_3}{3} ; Y = \frac{y_1 + y_2 + y_3}{3}$$



The coordinates of the incentre I (x,y) of the triangle  $P_1P_2P_3$  with vertices  $P_1(x_1, y_1)$  ;  $P_2(x_2, y_2)$

$P_3(x_3, y_3)$  and side length a,b,c is given by  $X = \frac{ax_1+bx_2+cx_3}{a+b+c}$   $\frac{ay_1+by_2+cy_3}{a+b+c}$

### Area of a triangle

The area of triangle having vertices  $P_1(x_1, y_1)$  ;  $P_2(x_2, y_2)$  and  $P_3(x_3, y_3)$  is given by  $= \frac{1}{2} [ (x_1y_2 - x_2y_1) + (x_2y_3 - x_3y_2) + (x_3y_1 - x_1y_3) ]$

*Note* – If area of is Zero  $\Rightarrow$  points  $P_1, P_2, P_3$  are collinear.

Condition for 4 points, no three of which are collinear to be a parallelogram. Let  $P_1(x_1, y_1)$

$P_2(x_2, y_2)$  ,  $P_3(x_3, y_3)$  ,  $P_4(x_4, y_4)$  be 4 points lying in a plane then  $P_1$  ,  $P_2$  ,  $P_3$  ,  $P_4$  are the vertices of

a parallelogram if  $x_1+x_3=x_2+x_4$  and  $y_1+y_3=y_2+y_4$




$p_1, p_2, p_3, p_4$  are the vertices of a square if in addition to above or  $x_4 - x_1 = y_3 - y_2, y_4 - y_1 = x_3 - x_2$

## Equation of a Line

### *Slope of a line*

The slope of a line is defined as tangent of the angle  $\alpha$  which the line makes in the positive direction of the x-axis in the anti-clockwise direction.



The slope is generally represented by 'm' thus in the notion used above  $m = \tan \alpha$

The equation of a line is given by  $y = mx + c$  where 'c' is the intercept made on the y-axis by the line.

Equation of x-axis is  $y = 0$

Equation of y-axis is  $x = 0$

2 lines are said to be parallel if they have the same slope that is if where  $m_1 = m_2$  where  $m_1$  and  $m_2$  are the slopes of the 2 lines.

2 lines said to be perpendicular if their product of their slope is -1. that is if  $m_1, m_2 = -1$  then lines are perpendicular to each other.  
Different forms of the equation of a line.

Line passing through point  $P(x_1, y_1)$  and having slope 'm'.  $(y - y_1) = m(x - x_1)$

Line passing through points  $P_1(x_1, y_1)$  and  $P_2(x_2, y_2)$ .  $y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$

If a line makes intercept a, b on x and y-axis respectively then equation of line is given by  $\frac{x}{a} + \frac{y}{b} = 1$

$$\theta = \tan^{-1} \left( \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| \right)$$

**Angle between 2 lines having slopes  $m_1$  and  $m_2$  is given by** where  $\theta$  is the actual

angle between the 2 lines.

**Perpendicular distance between parallel lines,** Let the equation of 2 lines be  $y = mx + c_1$  and line is given by

$$\left| \frac{C_1 - C_2}{\sqrt{1 + m^2}} \right|$$

*Perpendicular distance of a point from a line* . Let  $p(x_1, y_1)$  be any point and let  $y = mx + c$  be any line, then the perpendicular distance of point p from line  $y = mx + c$  is given by  $\left| \frac{y_1 - mx_1 - c}{\sqrt{1 + m^2}} \right|$

### Equation of a Circle

The equation of a circle having center at point  $(h, x)$  and radius 'r' given by  $(x - h)^2 + (y - x)^2 = r^2$

## Solved Examples

### Question

Find the equation of the line with slope 2 and intercept on the y-axis as -7?

### Solution

We know that equation of the line having slope 'm' and intercept on y-axis is 'c' the equation is  $y = mx + c$ , hence equation of line is  $y = 2x - 7$

The equation of a line which makes an intercept of 3 on x-axis and -3 on y-axis is

1.  $x - 2y = 5$
2.  $x - 2y = 3$
3.  $4x + 13y = 7$
4. none of these

### Solution

The line with the x-axis at point (3,0) and y-axis at the point (0,-3) hence equation of line is using

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1} (x - x_1)$$

$$y - 0 = \frac{-3 - 0}{0 - 3} (x - 3)$$

$y = x - 3$  or  $x - y = 3$  hence answer is option (2)

## Refresher Material

### Permutations

The number of arrangements of 'n' distinct object into 'n' distinct position is given by  $n, n-1, n-2, \dots, 3, 2, 1$ , which is often denoted as  $n!$  (read as n 'factorial'). The number of arrangements is also known as "Permutations". Thus number of permutation of n distinct object into n distinct places is denoted as  $P_n^n$ , read as permutations of n distinct objects taken n at a time.

The number of Permutations of n distinct objects taken from a group of 'r' distinct objects, in n distinct place is given by  $P_r^n = \frac{n!}{(n-r)!}$

The number of Permutations of n object out of which m are of one type, q is of second type etc. Such that  $n \neq m \neq q \dots$  is given by  $\frac{n!}{m! \cdot q! \dots}$

The number of Permutation of n distinct objects taken all at times around a circle is given by  $(n-1)!$  And  $\frac{(n-1)!}{2}$  accordingly as clockwise and anticlockwise arrangements are treated as different or same. This formula is valid for a necklace.

Number of arrangements of n objects, such that any p out of those occur together is given by  $(n-p+1)! \cdot p!$ .

Number of arrangements of n distinct things, when any object can be repeated any number of times is given by  $n^n$ .

### Combinations

Number of combinations of  $r$  distinct things taken out of  $n$  distinct things is given by

$$C_r^n = \frac{n!}{(n-r)!r!}$$

Please note in case of combinations also known as selections the order in which things are chosen is not important. Thus if we consider 4 distinct alphabets viz. A, B, C & D then no of permutations of any 3 alphabets out of is given by

ABC      ABD      ACB      ADB      ACD

ADC, etc whereas no. of selections is ABC, ABD, ACD and BCD i.e. 4 only.

Number of ways of selecting  $r$  things out of  $n$  distinct things is same as number of ways of selecting  $n-r$  things out of  $n$  things. i.e.

$$C_r^n = C_{n-r}^n$$

$$C_r^n = C_{r-1}^n + C_{r+1}^n$$

No of ways in which none or some objects can be chosen out of  $n$  distinct of objects is given by

$$2^n = C_0^n + C_1^n + C_2^n + \dots C_n^n$$

## Distribution

The total number of ways in which  $n$  identical things can be distributed into  $r$  distinct boxes, such that one or more than one box may remain empty, but not all the boxes can be empty is given by

$$C_{r-1}^{n+r-1}$$

If blank boxes are not allowed i.e each box has at least one object then no of ways is given by

Number of non-negative solutions of is given by

$$C_{r-1}^{n-1}$$

Number of Non – negative solution of  $x_1 + x_2 + \dots + x_r = n$  is given by

$$C_{r-1}^{n+r-1}$$

## Solved Example

### Question

A 3 digit number is formed using the digits 2, 3 and 4 without repeating any one of them what is the sum of all such possible numbers?

### Solution

Let us first find the total number of 3 digit numbers which be formed using the digits 2,3 and 4. The total number of such numbers is  $3! = 6$  now out of these 6 numbers we notice that number 2 appears in the hundreds place as well as in tens place and units place. So let us see how many numbers are there where 2 appears in the hundreds place we find there are exactly  $2!$  numbers. Similarly numbers 3 and 4 appear  $2!$  times in hundreds place. Hence if split each of the 6 numbers as say for example 234 as

$2 \times 100 + 3 \times 10 + 4$  then if we add all the numbers we see that numbers 2,3,4 appear in hundreds place

$2!$  times each. Sum of hundreds digits of all the numbers – Similarly sum of all the

$(2+3+4) \times 2! \times 100$  numbers in the tens and units digit is given by  $(2+3+4) \times 2! \times 10$  and

$(2+3+4) \times 2! \times 1$  therefore sum of all the numbers is

$$(2+3+4) \times 2! \times (100+10+1)$$

$$= 9 \times 2! \times 111$$

$$= 1998.$$

### Question

Atul has 9 friends; 4 males and 5 females. In how many ways can he invite them, if he wants to have exactly 3 females in the invites?

### Solution

The 3 girls which are to be invited can be selected in  ${}^5C_3$  ways. Also the remaining 4 males none, all or some can be invited in  $2^4$  ways. Hence Total no of ways Atul can invite his friends is

$${}^5C_3 \times 2^4 = 10 \cdot 16 = 160$$

### Refresher Material

#### Progression

A progression or sequence is defined as a succession of terms arranged in a definite according to some rule.

For example the sequence of Odd numbers 1, 3, 5, 7, 9, 11, etc.....

The numbers in the sequence are called the terms of the sequence. A sequence having a finite number of terms is known as finite sequence.

The first terms in a sequence is generally denoted by  $a_1$ , second term by  $a_2, a_3, \dots$  or  $T_2, T_3, \dots$  etc. The  $n^{\text{th}}$  term of the sequence is denoted by  $a_n$  or  $T_n$  and is also known as the general term of the sequence, as by assigning value to 'n',  $T_n$  can be made to represent any of the terms in the sequence.

#### Example

If the general term of a sequence is given by  $2n$ , find the First, sixth and  $8^{\text{th}}$  term.

Given  $T_n = 2n$

Therefore

First Term  $\Rightarrow n=1$

Therefore  $T_1 = 2 \cdot 1 = 2$

Similarly  $T_6 = 2 \cdot 6 = 12$  and

$T_8 = 2 \cdot 8 = 16$

## Sequences

### Some Standard type of sequences:

#### Arithmetic Progression (AP)

In this type of sequence, the difference between any two consecutive terms of the sequence is a constant. The constant is known as the 'Common difference.' Thus if the first term in AP is  $a$ , and Common difference is ' $d$ ' then second term is given by

$$T_2 = a+d, \text{ third terms } T_3 = a+2d \text{ etc.....}$$

in general  $T_n = a+(n-1)d$ .

Sum of  $n$  terms of an AP is given by  $S_n = \frac{n}{2}[2a+(n-1)d]$

Also  $S_n = \frac{n}{2}[T_1 + T_n]$

#### Geometric Progression (GP)

In this type of sequence the ratio of 2 consecutive terms is a constant. The constant ratio is known as common ratio and is usually denoted by ' $r$ '. Thus if ' $a$ ' is the first term of a GP

then  $T_2 = ar$ ,  $T_3 = ar^2$  etc.

Sum of n terms of a GP,  $S_n = \frac{a(r^n - 1)}{r - 1}$  if  $r > 1$

$$= \frac{a(1 - r^n)}{1 - r} \text{ if } r < 1$$

### Sum up to infinite terms of a GP

If  $r < 1$  then  $r^n \rightarrow 0$  as  $n \rightarrow \infty$  i.e. if let's say  $r = \frac{1}{2}$  then if we consider higher and higher values of  $n$  then  $r^n$  becomes smaller and smaller. Hence if we consider the sum up to very large number of terms, we say

the sum up to infinite terms,  $S_\infty = \frac{a}{1 - r}$

### Harmonic Progression (HP)

A sequence  $a_1, a_2, a_3, \dots$  is said to be a Harmonic Progression

if  $\frac{1}{a_1}, \frac{1}{a_2}, \dots$  are in AP.

The  $n^{\text{th}}$  term of HP is given by  $T_n = \frac{1}{a + (n - 1)d}$

There is no formula for finding the sum up to  $n$  terms of a HP.

Arithmetic mean (AM) of 2 positive numbers a,b is defined as  $AM = \frac{a+b}{2}$

Geometric mean (GM) of 2 positive numbers a,b is defined as  $GM = \sqrt{ab}$

Harmonic mean (HM) of 2 positive numbers a,b is defined as  $HM = \frac{2ab}{a+b}$  .

For any given a,b

$$AM \geq GM \geq HM$$

## Solved Example

### Question

Find the 10<sup>th</sup> terms of the series:  $5, \frac{7}{2}, \frac{7}{4}, \frac{9}{8}, \dots$

- (1)  $\frac{-9}{512}$   
(2)  $\frac{-15}{512}$   
(3)  $\frac{-1023}{512}$   
(4)  $\frac{-2559}{512}$

### Solution

$$4 + 1, 3 + \frac{1}{2}, 2 + \frac{1}{4}, \dots$$

Term of the series can be split as

We find 4, 3, 2, ..... are in AP with common difference of -1 hence 10<sup>th</sup> term is

given by  $T_{10} = 4 + (10-1)(-1) = -5$

[as  $T_n = a + (n-1)d$ ]

Also  $1, \frac{1}{2}, \frac{1}{4}$  are in GP

Hence  $T_{10} = 1 \cdot \left(\frac{1}{2}\right)^{10-1} = \frac{1}{2^9}$

[as  $T_n = a \cdot r^{n-1}$ ]

Hence  $T_{10} = -5 + \frac{1}{2^9} = \frac{-2559}{512}$

### Question

If the 8<sup>th</sup> term of an AP is 88 and the 88<sup>th</sup> term is 8, then 100<sup>th</sup> term is :

- (1) 1
- (2) 2
- (3) -4
- (4) 8

### Solution

Given 8<sup>th</sup> = 88 and 88<sup>th</sup> term = 8, let the first term of AP be 'a' and common difference be 'd'.

then  $T_8 = a + 7d$  and  $T_{88} = a + 87d$

$$\Rightarrow 88 = a + 7d \quad (1)$$

$$\text{and } 8 = a + 87d \quad (2)$$

Subtracting we get  $80 = -80d$

$$\Rightarrow d = -1$$

and putting (1) we get  $a = 95$

Hence,

$$T_{100} = a + 99d$$

$$= 95 + 99(-1)$$

$$= -4$$

Hence option (3)

- For More Sample questions –
  - <http://www.minglebox.com/online-test-preparation/cat-mba-exam>
- Practice Number Systems Refresher Tests –
  - <http://www.minglebox.com/online-test-preparation/cat-mba-exam>
- Discuss your doubts on CAT –
  - <http://www.minglebox.com/forum/mba>
- Minglebox e-CAT Prep test series –
  - <http://www.minglebox.com/online-test-preparation/cat-mba-exam>