

# AIPMT 2008 Examination Paper

1. Which two of the following five physical parameters have the same dimensions?

- (a) energy density
- (b) refractive index
- (c) dielectric constant
- (d) Young's modulus
- (e) magnetic field

- (1) (a) and (e)
- (2) (b) and (d)
- (3) (c) and (e)
- (4) (a) and (d)

2. If the error in the measurement of radius of a sphere is 2%, then the error in the determination of volume of the sphere will be

- (1) 2%
- (2) 4%
- (3) 6%
- (4) 8%


3. The distance travelled by a particle starting from rest and moving with an acceleration  $\frac{4}{3} \text{ ms}^{-2}$ , in the third second is

- (1)  $\frac{19}{3} \text{ m}$
- (2) 6 m
- (3) 4 m
- (4)  $\frac{10}{3} \text{ m}$

4. A particle moves in a straight line with a constant acceleration. It changes its velocity from  $10 \text{ ms}^{-1}$  to  $20 \text{ ms}^{-1}$  while passing through a distance 135 m in t second. The value of t is

- (1) 9
- (2) 10
- (3) 1.8
- (4) 12

5. A particle shows distance-time curve as given in this figure. The maximum instantaneous velocity of the particle is around the point



- (1) A
- (2) B
- (3) C
- (4) D


6. A particle of mass  $m$  is projected with velocity  $v$  making an angle of  $45^\circ$  with the horizontal. When the particle lands on the level ground the magnitude of the change in its momentum will be

- (1) zero
- (2)  $2mv$
- (3)  $mv/\sqrt{2}$
- (4)  $mv\sqrt{2}$

7. Sand is being dropped on a conveyor belt at the rate of  $M$  kg/s. The force necessary to keep the belt moving with a constant velocity of  $v$  m/s will be

- (1) Zero
- (2)  $Mv$  newton
- (3)  $2Mv$  newton
- (4)  $\frac{Mv}{2}$  newton

8. Three forces acting on a body are shown in the figure. To have the resultant force only along the  $y$ -direction, the magnitude of the minimum additional force needed is



- (1)  $\sqrt{3}$  N
- (2) 0.5 N
- (3) 1.5 N

(4)  $\frac{\sqrt{3}}{4} \text{ N}$

9. Water falls from a height of 60 m at the rate of 15 kg/s to operate a turbine. The losses due to frictional forces are 10% of energy. How much power is generated by the turbine? ( $g = 10 \text{ m/s}^2$ )

- (1) 7.0 kW
- (2) 8.1 kW
- (3) 10.2 kW
- (4) 12.3 kW

10. A shell of mass 200 gm is ejected from a gun of mass 4 kg by an explosion that generates 1.05 kJ of energy. The initial velocity of the shell is

- (1)  $120 \text{ ms}^{-1}$
- (2)  $100 \text{ ms}^{-1}$
- (3)  $80 \text{ ms}^{-1}$
- (4)  $40 \text{ ms}^{-1}$

11. The ratio of the radii of gyration of a circular disc to that of a circular ring, each of same mass and radius, around their respective axes is

- (1)  $\sqrt{2} : \sqrt{3}$
- (2)  $\sqrt{3} : \sqrt{2}$
- (3)  $1 : \sqrt{2}$
- (4)  $\sqrt{2} : 1$

12. A thin rod of length  $L$  and mass  $M$  is bent at its midpoint into two halves so that the angle between them is  $90^\circ$ . The moment of inertia of the bent rod about an axis passing through the bending point and perpendicular to the plane defined by the two halves of the rod is

- (1)  $\frac{\sqrt{2} ML^2}{24}$
- (2)  $\frac{ML^2}{24}$
- (3)  $\frac{12}{ML^2}$
- (4)  $\frac{ML^2}{6}$

13. A roller coaster is designed such that riders experience "weightlessness" as they go round the top of a hill whose radius of curvature is 20 m. The speed of the car at the top of the hill is between

- (1) 13 m/s and 14 m/s
- (2) 14 m/s and 15 m/s

- (3) 15 ms/ and 16 m/s
- (4) 16 m/s and 17 m/s

14. If Q, E and W denote respectively the heat added, change in internal energy and the work done in a closed cycle process, then

- (1)  $Q = 0$
- (2)  $W = 0$
- (3)  $Q = W = 0$
- (4)  $E = 0$

15. On a new scale of temperature (which is linear) and called the W scale, the freezing and boiling points of water are  $39^\circ \text{W}$  and  $239^\circ \text{W}$  respectively. What will be the temperature on the new scale, corresponding to a temperature of  $39^\circ \text{C}$  on the Celsius scale?

- (1)  $139^\circ \text{W}$
- (2)  $78^\circ \text{W}$
- (3)  $117^\circ \text{W}$
- (4)  $200^\circ \text{W}$

16. At  $10^\circ \text{C}$  the value of the density of a fixed mass of an ideal gas divided by its pressure is x. At  $110^\circ \text{C}$  this ratio is

- (1)  $\frac{283}{383}x$
- (2) x
- (3)  $\frac{383}{283}x$
- (4)  $\frac{10}{110}x$

17. Two Simple Harmonic Motions of angular frequency 100 and  $1000 \text{ rad s}^{-1}$  have the same displacement amplitude. The ratio of their maximum accelerations is

- (1)  $1 : 10^4$
- (2)  $1 : 10$
- (3)  $1 : 10^2$
- (4)  $1 : 10^3$

18. The wave described by  $y = 0.25 \sin(10\pi x - 2\pi t)$ , where x and y are in meters and t in seconds, is a wave travelling along the

- (1) -ve x direction with amplitude 0.25 m and wavelength  $\lambda = 0.2 \text{ m}$
- (2) -ve x direction with frequency 1 Hz
- (3) +ve x direction with frequency  $\pi \text{ Hz}$  and wavelength  $\lambda = 0.2 \text{ m}$
- (4) +ve x direction with frequency 1 Hz and wavelength  $\lambda = 0.2 \text{ m}$

19. A point performs simple harmonic oscillation of period T and the equation of motion is given by  $x = a \sin(\omega t + \pi/6)$ . After the elapse of what fraction of the time

period the velocity of the point will be equal to half of its maximum velocity?

- (1)  $\frac{T}{12}$
- (2)  $\frac{T}{8}$
- (3)  $\frac{T}{6}$
- (4)  $\frac{T}{3}$

20. Two points are located at a distance of 10 m and 15 m from the source of oscillation. The period of oscillation is 0.05 sec and the velocity of the wave is 300 m/sec. What is the phase difference between the oscillations of two points?

- (1)  $\frac{\pi}{6}$
- (2)  $\frac{\pi}{3}$
- (3)  $\frac{2\pi}{3}$
- (4)  $\pi$

21. The velocity of electromagnetic radiation in a medium of permittivity  $\epsilon_0$  and permeability  $\mu_0$  is given by

- (1)  $\sqrt{\frac{\mu_0}{\epsilon_0}}$
- (2)  $\sqrt{\frac{\epsilon_0}{\mu_0}}$
- (3)  $\sqrt{\mu_0 \epsilon_0}$
- (4)  $\frac{1}{\sqrt{\mu_0 \epsilon_0}}$

22. Two periodic waves of intensities  $I_1$  and  $I_2$  pass through a region at the same time in the same direction. The sum of the maximum and minimum intensities is

- (1)  $2(I_1 + I_2)$
- (2)  $I_1 + I_2$
- (3)  $(\sqrt{I_1} + \sqrt{I_2})^2$
- (4)  $(\sqrt{I_1} - \sqrt{I_2})^2$

23. Two thin lenses of focal lengths  $f_1$  and  $f_2$  are in contact and coaxial. The power of the combinations is

- (1)  $\frac{f_1 + f_2}{f_1 f_2}$
- (2)  $\sqrt{\frac{f_1}{f_2}}$
- (3)  $\sqrt{\frac{f_2}{f_1}}$
- (4)  $\frac{f_1 + f_2}{2}$


24. A boy is trying to start a fire by focusing Sunlight on a piece of paper using an equiconvex lens of focal length 10 cm. The diameter of the Sun is  $1.39 \times 10^9$  m and its mean distance from the earth is  $1.5 \times 10^{11}$  m. What is the diameter of the Sun's image on the paper?

- (1)  $12.4 \times 10^{-4}$  m
- (2)  $9.2 \times 10^{-4}$  m
- (3)  $6.5 \times 10^{-4}$  m
- (4)  $6.5 \times 10^{-5}$  m

25. The energy required to charge a parallel plate condenser of plate separation  $d$  and plate area of cross-section  $A$  such that the uniform electric field between the plates is  $E$ , is

- (1)  $\frac{1}{2} \epsilon_0 E^2 A d$
- (2)  $\frac{1}{2} \epsilon_0 E^2 / A.d$
- (3)  $\epsilon_0 E^2 / A d$
- (4)  $\epsilon_0 E^2 A d$

26. A thin conducting ring of radius  $R$  is given a charge  $+Q$ . The electric field at the centre  $O$  of the ring due to the charge on the part  $AKB$  of the ring is  $E$ . The electric field at the centre due to the charge on the part  $ACDB$  of the ring is



- (1)  $3 E$  along  $OK$
- (2)  $3 E$  along  $KO$
- (3)  $E$  along  $OK$

(4) E along KO

27. The electric potential at a point in free space due to a charge  $Q$  coulomb is  $Q \times 10^{11}$  volts. The electric field at that point is

- (1)  $12\pi\epsilon_0 Q \times 10^{22}$  volt/m
- (2)  $4\pi\epsilon_0 Q \times 10^{22}$  volt/m
- (3)  $12\pi\epsilon_0 Q \times 10^{20}$  volt/m
- (4)  $4\pi\epsilon_0 Q \times 10^{20}$  volt/m


28. A cell can be balanced against 110 cm and 100 cm of potentiometer wire, respectively with and without being short circuited through a resistance of  $10 \Omega$ . Its internal resistance is

- (1) Zero
- (2) 1.0 ohm
- (3) 0.5 ohm
- (4) 2.0 ohm

29. A wire of a certain material is stretched slowly by ten per cent. Its new resistance and specific resistance become respectively

- (1) 1.1 times, 1.1 times
- (2) 1.2 times, 1.1 times
- (3) 1.21 times, same
- (4) Both remain the same

30. In the circuit shown, the current through the  $4 \Omega$  resistor is 1 amp when the points P and M are connected to a d.c. voltage source. The potential difference between the points M and N is




- (1) 3.2 volt
- (2) 1.5 volt
- (3) 1.0 volt
- (4) 0.5 volt

31. An electric kettle takes 4A current at 220 V. How much time will it take to boil 1 kg of water from temperature 20°C? The temperature of boiling water is 100°C.

- (1) 4.2 min
- (2) 6.3 min
- (3) 8.4 min
- (4) 12.6 min

32. A current of 3 amp. flows through the  $2\ \Omega$  resistor shown in the circuit. The power dissipated in the  $5\ \Omega$  resistor is



- (1) 5 watt
- (2) 4 watt
- (3) 2 watt
- (4) 1 watt

33. A particle of mass  $m$ , charge  $Q$  and kinetic energy  $T$  enters a transverse uniform magnetic field of induction  $\vec{B}$ . After 3 seconds the kinetic energy of the particle will be

- (1) 4 T
- (2) 3 T
- (3) 2 T
- (4) T

34. A closed loop PQRS carrying a current is placed in a uniform magnetic field. If the magnetic forces on segments PS, SR and RQ are  $F_1$ ,  $F_2$  and  $F_3$  respectively and are in the plane of the paper and along the directions shown, the force on the segment QP is

- (1)  $F_3 - F_1 + F_2$
- (2)  $F_3 - F_1 - F_2$
- (3)  $\sqrt{(F_3 - F_1)^2 + F_2^2}$
- (4)  $\sqrt{(F_3 - F_1)^2 - F_2^2}$

35. A circular disc of radius 0.2 meter is placed in a uniform magnetic field of induction  $\frac{1}{\pi} \left( \frac{\omega b}{m^2} \right)$  in such a way that its axis makes an angle of  $60^\circ$  with  $\vec{B}$ . The magnetic flux linked with the disc is



- (1) 0.01 wb
- (2) 0.02 wb
- (3) 0.06 wb
- (4) 0.08 wb

36. A galvanometer of resistance  $50\ \Omega$  is connected to a battery of 3 V along with a resistance of  $2950\ \Omega$  in series. A full scale deflection of 30 divisions is obtained in the galvanometer. In order to reduce this deflection to 20 divisions, the resistance in series should be

- (1) 4450  $\Omega$
- (2) 5050  $\Omega$
- (3) 5550  $\Omega$
- (4) 6050  $\Omega$

37. Curie temperature is the temperature above which

- (1) Ferromagnetic material becomes diamagnetic material
- (2) Ferromagnetic material becomes paramagnetic material
- (3) Paramagnetic material becomes diamagnetic material
- (4) Paramagnetic material becomes ferromagnetic material

38. A long solenoid has 500 turns. When a current of 2 ampere is passed through it, the resulting magnetic flux linked with each turn of the solenoid is  $4 \times 10^{-3}$  wb. The self-inductance of the solenoid is

- (1) 4.0 henry
- (2) 2.5 henry
- (3) 2.0 henry
- (4) 1.0 henry

39. In an a.c. circuit the e.m.f. ( $e$ ) and the current ( $i$ ) at any instant are given respectively by

$$e = E_0 \sin \omega t$$

$$i = I_0 \sin(\omega t - \phi)$$

The average power in the circuit over one cycle of a.c. is

- (1)  $E_0 I_0$
- (2)  $\frac{E_0 I_0}{2}$
- (3)  $\frac{E_0 I_0}{2} \sin \phi$
- (4)  $\frac{E_0 I_0}{2} \cos \phi$

40. In the phenomenon of electric discharge through gases at low pressure, the coloured

glow in the tube appears as a result of

- (1) Collision between different electrons of the atoms of the gas
- (2) Excitation of electrons in the atoms
- (3) Collision between the atoms of the gas
- (4) Collisions between the charged particles emitted from the cathode and the atoms of the gas

41. The work function of a surface of a photosensitive material is 6.2 eV. The wavelength of the incident radiation for which the stopping potential is 5 V lies in the

- (1) X-ray region
- (2) Ultraviolet region
- (3) Visible region
- (4) Infrared region

42. A particle of mass 1 mg has the same wavelength as an electron moving with a velocity of  $3 \times 10^6 \text{ ms}^{-1}$ . The velocity of the particle is (mass of electron =  $9.1 \times 10^{-31} \text{ kg}$ )

- (1)  $2.7 \times 10^{-21} \text{ ms}^{-1}$
- (2)  $2.7 \times 10^{-18} \text{ ms}^{-1}$
- (3)  $9 \times 10^{-2} \text{ ms}^{-1}$
- (4)  $3 \times 10^{-31} \text{ ms}^{-1}$

43. The ground state energy of hydrogen atom is -13.6 eV. When its electron is in the first excited state, its excitation energy is

- (1) 0
- (2) 3.4 eV
- (3) 6.8 eV
- (4) 10.2 eV

44. Two radioactive materials  $X_1$  and  $X_2$  have decay constants  $5\lambda$  and  $\lambda$  respectively. If initially they have the same number of nuclei, then the ratio of the number of nuclei of  $X_1$

to that of  $X_2$  will be  $\frac{1}{e}$  after a time

- (1)  $\frac{e}{\lambda}$
- (2)  $\lambda$
- (3)  $\frac{1}{2}\lambda$
- (4)  $\frac{1}{4\lambda}$

45. Two nuclei have their mass numbers in the ratio of 1:3. The ratio of their nuclear densities would be

- (1) 1 : 1
- (2) 1 : 3
- (3) 3 : 1

(4)  $(3)^{1/3} : 1$

46. If  $M(A, Z)$ ,  $M_p$  and  $M_n$  denote the masses of the nucleus  ${}^A_ZX$ , proton and neutron respectively in units of  $u$  ( $1u = 931.5 \text{ MeV} / c^2$ ) and  $BE$  represents its bonding energy in  $\text{MeV}$ , then

(1)  $M(A, Z) = ZM_p + (A - Z) M_n + BE / c^2$

(2)  $M(A, Z) = ZM_p + (A - Z) M_n - BE / c^2$

(3)  $M(A, Z) = ZM_p + (A - Z) M_n + BE$

(4)  $M(A, Z) = ZM_p + (A - Z) M_n - BE$

47. The voltage gain of an amplifier with 9% negative feedback is 10. The voltage gain without feedback will be

(1) 100

(2) 90

(3) 10

(4) 1.25

48. If the lattice parameter for a crystalline structure is  $3.6 \text{ \AA}$ , then the atomic radius in fcc crystal is


(1)  $1.27 \text{ \AA}$

(2)  $1.81 \text{ \AA}$

(3)  $2.10 \text{ \AA}$

(4)  $2.92 \text{ \AA}$

49. The circuit



is equivalent to

(1) OR gate

(2) AND gate

(3) NAND gate

(4) NOR gate

50. A p-n photodiode is made of a material with a band gap of  $2.0 \text{ eV}$ . The minimum frequency of the radiation that can be absorbed by the material is nearly

(1)  $20 \times 10^{14} \text{ Hz}$

(2)  $10 \times 10^{14} \text{ Hz}$

(3)  $5 \times 10^{14} \text{ Hz}$



55. Kohlrausch's law states that at

- (1) Infinite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte, whatever be the nature of the other ion of the electrolyte
- (2) Finite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte, whatever be the nature of the other ion of the electrolyte
- (3) Infinite dilution each ion makes definite contribution to equivalent conductance of an electrolyte depending on the nature of the other ion of the electrolyte
- (4) Infinite dilution, each ion makes definite contribution to conductance of an electrolyte whatever be the nature of the other ion of the electrolyte


56. The measurement of the electron position is associated with an uncertainty in momentum, which is equal to  $1 \times 10^{-18} \text{ g cm s}^{-1}$ . The uncertainty in electron velocity is, (mass of an electron is  $9 \times 10^{-28} \text{ g}$ )

- (1)  $1 \times 10^{11} \text{ cm s}^{-1}$
- (2)  $1 \times 10^9 \text{ cm s}^{-1}$
- (3)  $1 \times 10^6 \text{ cm s}^{-1}$
- (4)  $1 \times 10^5 \text{ cm s}^{-1}$

57. Which of the following are not state functions?

- (I)  $q + w$
  - (II)  $q$
  - (III)  $w$
  - (IV)  $H-TS$
- (1) (II) and (III)
  - (2) (I) and (IV)
  - (3) (II), (III) and (IV)
  - (4) (I), (II) and (III)

58. The bromination of acetone that occurs in acid solution is represented by this equation



These kinetic data were obtained for given reaction concentrations

Initial concentrations, M		
[CH <sub>3</sub> COCH <sub>3</sub> ]	[Br <sub>2</sub> ]	[H <sup>+</sup> ]
0.30	0.05	0.05
0.30	0.10	0.05
0.30	0.10	0.10
0.40	0.05	0.20

Initial rate, disappearance of Br<sub>2</sub>, M s<sup>-1</sup>

$$5.7 \times 10^{-5}$$

$$5.7 \times 10^{-5}$$

$$1.2 \times 10^{-4}$$

$$3.1 \times 10^{-4}$$

Based on these data, the rate equation is

(1) Rate =  $k[\text{CH}_3\text{COCH}_3][\text{Br}_2][\text{H}^+]$

(2) Rate =  $k[\text{CH}_3\text{COCH}_3][\text{H}^+]$

(3) Rate =  $k[\text{CH}_3\text{COCH}_3][\text{Br}_2]$

(4) Rate =  $k[\text{CH}_3\text{COCH}_3][\text{Br}_2][\text{H}^+]^2$

59. What volume of oxygen gas (O<sub>2</sub>) measured at 0°C and 1 atm, is needed to burn completely 1 L of propane gas (C<sub>3</sub>H<sub>8</sub>) measured under the same conditions?

(1) 10 L

(2) 7 L

(3) 6 L

(4) 5 L

60. Bond dissociation enthalpy of H<sub>2</sub>, Cl<sub>2</sub> and HCl are 434, 242 and 431 kJmol<sup>-1</sup> respectively. Enthalpy of formation of HCl is

(1) 245 kJmol<sup>-1</sup>

(2) 93 kJmol<sup>-1</sup>

(3) -245 kJmol<sup>-1</sup>

(4) -93 kJmol<sup>-1</sup>

61. Which of the following statements is not correct?

(1) The number of Bravais lattices in which a crystal can be categorized is 14

(2) The fraction of the total volume occupied by the atoms in a primitive cell is 0.48

(3) Molecular solids are generally volatile

(4) The number of carbon atoms in a unit cell of diamond is 4

62. Equal volumes of three acid solutions of pH 3, 4 and 5 are mixed in a vessel. What will be the H<sup>+</sup> ion concentration in the mixture?

(1)  $1.11 \times 10^{-3}$  M

(2)  $1.11 \times 10^{-4}$  M

- (3)  $3.7 \times 10^{-4}$  M  
 (4)  $3.7 \times 10^{-3}$  M

63. The values of  $K_{p1}$  and  $K_{p2}$  for the reactions  
 $X \rightleftharpoons Y+Z$ .....(1) and



are in ratio of 9:1. If degree of dissociation of X and A be equal, then total pressure at equilibrium (1) and (2) are in the ratio

- (1) 1 : 1  
 (2) 3 : 1  
 (3) 1 : 9  
 (4) 36 : 1

64. If the concentration of  $OH^-$  ions in the reaction

$Fe(OH)_3(s) \rightleftharpoons Fe^{3+}(aq) + 3OH^-(aq)$  is decreased by  $\frac{1}{4}$  times, then equilibrium concentration of  $Fe^{3+}$  will increase by

- (1) 4 times  
 (2) 8 times  
 (3) 16 times  
 (4) 64 times

65. For the gas phase reaction,  $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$  Which of the following conditions is correct?

- (1)  $\Delta H > 0$  and  $\Delta S < 0$   
 (2)  $\Delta H = 0$  and  $\Delta S < 0$   
 (3)  $\Delta H > 0$  and  $\Delta S > 0$   
 (4)  $\Delta H < 0$  and  $\Delta S < 0$

66. The rate constants  $k_1$  and  $k_2$  for two different reactions are

$10^{16} \cdot e^{-2000/T}$  and  $10^{15} \cdot e^{-1000/T}$  respectively. The temperature at which  $k_1 = k_2$  is

- (1)  $\frac{1000}{2.303}$  K  
 (2) 1000 K  
 (3)  $\frac{2000}{2.303}$  K  
 (4) 2000 K

67. Standard free energies of formation (in kJ/mol) at 298 K are -237.2, -394.4 and -8.2 for  $H_2O(l)$ ,  $CO_2(g)$  and pentane (g) respectively. The value of  $E^\circ_{cell}$  for the pentane-oxygen fuel cell is

- (1) 0.0968 V

- (2) 1.968 V
- (3) 2.0968 V
- (4) 1.0968 V

68. The dissociation equilibrium of a gas  $AB_2$  can be represented as,  
 $2AB_2(g) \rightleftharpoons 2AB(g) + B_2(g)$ . The degree of dissociation is 'x' and is small compared to 1. The expression relating the degree of dissociation (x) with equilibrium constant  $K_p$  and total pressure P is

- (1)  $\left(\frac{K_p}{P}\right)$
- (2)  $\left(\frac{2K_p}{P}\right)$
- (3)  $\left(\frac{2K_p}{P}\right)^{1/3}$
- (4)  $\left(\frac{2K_p}{P}\right)^{1/2}$

69. The sequence of ionic mobility in aqueous solution is

- (1)  $Na^+ > K^+ > Rb^+ > Cs^+$
- (2)  $K^+ > Na^+ > Rb^+ > Cs^+$
- (3)  $Cs^+ > Rb^+ > K^+ > Na^+$
- (4)  $Rb^+ > K^+ > Cs^+ > Na^+$

70. Percentage of free space in a body centred cubic unit cell is

- (1) 28%
- (2) 30%
- (3) 32%
- (4) 34%

71. The correct order of decreasing second ionisation enthalpy of Ti (22), V (23), Cr (24) and Mn (25) is

- (1)  $Ti > V > Cr > Mn$
- (2)  $Cr > Mn > V > Ti$
- (3)  $V > Mn > Cr > Ti$
- (4)  $Mn > Cr > Ti > V$

72. How many moles of lead (II) chloride will be formed from a reaction between 6.5 g of PbO and 3.2 g of HCl?

- (1) 0.029
- (2) 0.044
- (3) 0.333
- (4) 0.011



73. Which of the following complexes exhibits the highest paramagnetic behaviour?  
Where gly = glycine, en = ethylenediamine and bpy = bipyridyl moities). (At. number Ti = 22, V = 23, Fe = 26, Co = 27)

- (1)  $[\text{Ti}(\text{NH}_3)_6]^{3+}$
- (2)  $[\text{V}(\text{gly})_2(\text{OH})_2(\text{NH}_3)_2]^+$
- (3)  $[\text{Fe}(\text{en})(\text{bpy})(\text{NH}_3)_2]^{2+}$
- (4)  $[\text{Co}(\text{OX})_2(\text{OH})_2]^-$

74. Volume occupied by one molecule of water (density =  $1 \text{ g cm}^{-3}$ ) is

- (1)  $5.5 \times 10^{-23} \text{ cm}^3$
- (2)  $9.0 \times 10^{-23} \text{ cm}^3$
- (3)  $6.023 \times 10^{-23} \text{ cm}^3$
- (4)  $3.0 \times 10^{-23} \text{ cm}^3$

75. Number of moles of  $\text{MnO}_4^-$  required to oxidize one mole of ferrous oxalate completely in acidic medium will be

- (1) 0.2 moles
- (2) 0.6 moles
- (3) 0.4 moles
- (4) 7.5 moles

76. On the basis of the following  $E^\circ$  values, the strongest oxidizing agent is



$$E^\circ = -0.35 \text{ V}$$



$$E^\circ = -0.77 \text{ V}$$

- (1)  $[\text{Fe}(\text{CN})_6]^{3-}$
- (2)  $[\text{Fe}(\text{CN})_6]^{4-}$
- (3)  $\text{Fe}^{2+}$
- (4)  $\text{Fe}^{3+}$

77. The alkali metals form salt-like hydrides by the direct synthesis at elevated temperature. The thermal stability of these hydrides decreases in which of the following orders?

- (1)  $\text{LiH} > \text{NaH} > \text{KH} > \text{RbH} > \text{CsH}$
- (2)  $\text{CsH} > \text{RbH} > \text{KH} > \text{NaH} > \text{LiH}$
- (3)  $\text{KH} > \text{NaH} > \text{LiH} > \text{CsH} > \text{RbH}$
- (4)  $\text{NaH} > \text{LiH} > \text{KH} > \text{RbH} > \text{CsH}$

78. Which one of the following arrangements does not give the correct picture of the trends indicated against it?

- (1)  $\text{F}_2 > \text{Cl}_2 > \text{Br}_2 > \text{I}_2$  : Electronegativity
- (2)  $\text{F}_2 > \text{Cl}_2 > \text{Br}_2 > \text{I}_2$  : Oxidizing power
- (3)  $\text{F}_2 > \text{Cl}_2 > \text{Br}_2 > \text{I}_2$  : Electron gain enthalpy
- (4)  $\text{F}_2 > \text{Cl}_2 > \text{Br}_2 > \text{I}_2$  : Bond dissociation energy

79. With which one of the following elements silicon should be doped so as to give p-type of semiconductor?

- (1) Boron
- (2) Germanium
- (3) Arsenic
- (4) Selenium

80. In which of the following coordination entities the magnitude of  $\Delta O$  (CFSE in octahedral field) will be maximum (at. no. Co = 27)?

- (1)  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$
- (2)  $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$
- (3)  $[\text{Co}(\text{NH}_3)_6]^{3+}$
- (4)  $[\text{Co}(\text{CN})_6]^{3-}$

81. The angular shape of molecule ( $\text{O}_3$ ) consists of

- (1) 2 sigma and 1 pi bond
- (2) 1 sigma and 2 pi bonds
- (3) 2 sigma and 2 pi bonds
- (4) 1 sigma and 1 pi bond

82. The correct order of increasing bond angles in the following triatomic species is

- (1)  $\text{NO}_2^+ < \text{NO}_2^- < \text{NO}_2$
- (2)  $\text{NO}_2^- < \text{NO}_2^+ < \text{NO}_2$
- (3)  $\text{NO}_2^- < \text{NO}_2 < \text{NO}_2^+$
- (4)  $\text{NO}_2^+ < \text{NO}_2 < \text{NO}_2^-$


83. Four diatomic species are listed below the different sequences. Which of these presents the correct order of their increasing bond order?

- (1)  $\text{He}_2^+ < \text{O}_2^- < \text{NO} < \text{C}_2^{2-}$
- (2)  $\text{O}_2^- < \text{NO} < \text{C}_2^{2-} < \text{He}_2^+$
- (3)  $\text{NO} < \text{C}_2^{2-} < \text{O}_2^- < \text{He}_2^+$
- (4)  $\text{C}_2^{2-} < \text{He}_2^+ < \text{NO} < \text{O}_2^-$

84. Equimolar solutions of the following were prepared in water separately. Which one of the solutions will record the highest pH?

- (1)  $\text{CaCl}_2$
- (2)  $\text{SrCl}_2$
- (3)  $\text{BaCl}_2$
- (4)  $\text{MgCl}_2$

85. In the hydrocarbon



The state of hybridization of carbons 1, 3 and 5 are in the following sequence

- (1)  $\text{sp}^3$ ,  $\text{sp}^2$ ,  $\text{sp}$
- (2)  $\text{sp}^2$ ,  $\text{sp}$ ,  $\text{sp}^3$
- (3)  $\text{sp}$ ,  $\text{sp}^3$ ,  $\text{sp}^2$
- (4)  $\text{sp}$ ,  $\text{sp}^2$ ,  $\text{sp}^3$

86. Green chemistry means such reactions which

- (1) Study the reactions in plants
- (2) Produce colour during reactions
- (3) Reduce the use and production of hazardous chemicals
- (4) Are related to the depletion of ozone layer

87. A strong base can abstract an  $\alpha$ -hydrogen from




- (1) Alkane
- (2) Alkene
- (3) Amine
- (4) Ketone

88. How many stereoisomers does the molecule have?



- (1) 2
- (2) 4
- (3) 6
- (4) 8

89. The stability of carbanions in the following



is in the order of




- (1)  $a > c > b > d$
- (2)  $a > b > c > d$
- (3)  $b > c > d > a$
- (4)  $d > b > c > a$

90. The relative reactivities of acyl compounds towards nucleophilic substitution are in the order of

- (1) Acyl chloride > Ester > Acid anhydride > Amide

- (2) Acyl chloride > Acid anhydride > Ester > Amide  
 (3) Ester > Acyl chloride > Amide > Acid anhydride  
 (4) Acid anhydride > Amide > Ester > Acyl chloride


91. Base strength of







is in the order of

- (1)  $a > b > c$   
 (2)  $b > a > c$   
 (3)  $c > b > a$   
 (4)  $a > c > b$

92.







A (predominantly) is



93. In DNA, the complimentary bases are

- (1) Uracil and adenine; cytosine and guanine
- (2) Adenine and thymine; guanine and cytosine
- (3) Adenine and thymine; guanine and uracil
- (4) Adenine and guanine; thymine and cytosine


94. Which one the following is most reactive towards electrophilic attack?

- (1) 
- (2) 
- (3) 
- (4) 




95. An organic compound contains carbon, hydrogen and oxygen. Its elemental analysis gave C, 38.71% and H, 9.67%. The empirical formula of the compound would be

- (1) CH<sub>4</sub>O
- (2) CH<sub>3</sub>O
- (3) CH<sub>2</sub>O
- (4) CHO

96. In a S<sub>N</sub>2 substitution reaction of the type




Which one of the following has the highest relative rate?

- (1) CH<sub>3</sub>CH<sub>2</sub>Br
- (2) 
- (3) 
- (4) 

97. Acetophenone when reacted with a base, C<sub>2</sub>H<sub>5</sub>ONa, yields a stable compound which has the structure?

- (1)
- (2)
- (3)
- (4)

98. In a reaction of aniline a coloured product C was obtained.



The structure of C would be

- (1)
- (2)
- (3)
- (4)

99. Which of the following statements is not true?

- (1) Natural rubber is a 1, 4 - polymer of isoprene
- (2) In vulcanization, the formation of sulphur bridges between different chains make rubber harder and stronger
- (3) Natural rubber has the trans-configuration at every double bond
- (4) Buna-S is a copolymer of butadiene and styrene

100. Which one of the following is an amine hormone?

- (1) Progesterone
- (2) Thyroxine
- (3) Oxyurin

(4) Insulin

101. Select one of the following of important features distinguishing Gnetum from Cycas and Pinus and showing affinities with angiosperms

- (1) Embryo development and apical meristem
- (2) Absence of resin duct and leaf venation
- (3) Presence of vessel elements and absence of archegonia
- (4) Perianth and two integuments

102. Thermococcus, Methanococcus and Methanobacterium exemplify

- (1) Bacteria that contain a cytoskeleton and ribosomes
- (2) Archaeobacteria that contain protein homologous to eukaryotic core histones
- (3) Archaeobacteria that lack any histones resembling those found in eukaryotes but whose DNA is negatively supercoiled
- (4) Bacteria whose DNA is relaxed or positively supercoiled but which have a cytoskeleton as well as mitochondria

103. Which one of the following is heterosporous?

- (1) Equisetum
- (2) Dryopteris
- (3) Salvinia
- (4) Adiantum

104. In which one of the following male and female gametophytes do not have free living independent existence?

- (1) Cedrus
- (2) Pteris
- (3) Funaria
- (4) Polytrichum

105. Which one of the following groups of three animals each is correctly matched with their one characteristic morphological feature?

Animals	Morphological feature
(1) Cockroach, Locust, Taenia	- Metameric segmentation
(2) Liver fluke, Sea anemone, Sea cucumber	- Bilateral symmetry
(3) Centipede, Prawn, Sea urchin	- Jointed appendages
(4) Scorpion, Spider, Cockroach	- Ventral solid central nervous system

106. Which one of the following phyla is correctly matched with its two general characteristics?

- (1) Mollusca - Normally oviparous and development through a trochophore or veliger larva
- (2) Arthropoda - Body divided into head, thorax and abdomen and respiration by tracheae
- (3) Chordata - Notochord at some stage and separate anal and urinary openings to the outside
- (4) Echinodermata - Pentamerous radial symmetry and mostly internal fertilization

107. Which one of the following in birds, indicates their reptilian ancestry?

- (1) Eggs with a calcareous shell
- (2) Scales on their hind limbs
- (3) Four-chambered heart
- (4) Two special chambers crop and gizzard in their digestive tract

108. Ascaris is characterized by

- (1) Presence of true coelom and metamerism (metamerisation)
- (2) Absence of true coelom but presence of metamerism
- (3) Presence of neither true coelom nor metamerism
- (4) Presence of true coelom but absence of metamerism

109. Which one of the following is not a characteristic of phylum Annelida?

- (1) Ventral nerve cord
- (2) Closed circulatory system
- (3) Segmentation
- (4) Pseudocoelom

110. Cellulose is the major component of cell walls of

- (1) Saccharomyces
- (2) Pythium
- (3) Xanthomonas
- (4) Pseudomonas

111. Vacuole in a plant cell

- (1) Lacks membrane and contains water and excretory substances
- (2) Is membrane-bound and contains storage proteins and lipids
- (3) Is membrane-bound and contains water and excretory substances
- (4) Lacks membrane and contains air

112. A competitive inhibitor of succinic dehydrogenase is



- (1) Malate
- (2) Malonate
- (3) Oxaloacetate
- (4)  $\alpha$ -ketoglutarate

113. Polysome is formed by

- (1) Ribosomes attached to each other in a linear arrangement
- (2) Several ribosomes attached to a single mRNA
- (3) Many ribosomes attached to a strand of endoplasmic reticulum
- (4) A ribosome with several subunits

114. Carbohydrates are commonly found as starch in plant storage organs. Which of the following five properties of starch (a - e) make it useful as a storage material?

- a. Easily translocated
- b. Chemical non-reactive
- c. Easily digested by animals
- d. Osmotically inactive
- e. Synthesized during photosynthesis

The useful properties are

- (1) Both a & e
- (2) Both b & c
- (3) Both b & d
- (4) a, c & e

115. In the light of recent classification of living organisms into three domains of life (bacteria, archaea and eukarya), which one of the following statements is true about archaea?

- (1) Archaea completely differ from prokaryotes
- (2) Archaea resemble eukarya in all respects
- (3) Archaea have some novel features that are absent in other prokaryotes and eukaryotes
- (4) Archaea completely differ from both prokaryotes and eukaryotes

116. Keeping in view the 'fluid mosaic model' for the structure of cell membrane, which one of the following statements is correct w.r.t. the movement of lipids and proteins from one lipid monolayer to the other (described as flip-flop movement)?

- (1) Neither lipids nor proteins can flip-flop
- (2) Both lipids and proteins can flip-flop
- (3) While lipids can rarely flip-flop, proteins cannot
- (4) While proteins can flip-flop, lipids cannot

117. In germinating seeds fatty acids are degraded exclusively in the

- (1) Mitochondria

- (2) Proplastids
- (3) Glyoxysomes
- (4) Peroxisomes

118. The two sub-units of ribosome remain united at a critical ion level of

- (1) Calcium
- (2) Copper
- (3) Manganese
- (4) Magnesium

119. Thorn of Bougainvillea and tendril of cucurbita are example of

- (1) Retrogressive evolution
- (2) Analogous organs
- (3) Homologous organs
- (4) Vestigial organs

120. Haploids are more suitable for mutation studies than the diploids. This is because

- (1) All mutations, whether dominant or recessive are expressed in haploids
- (2) Haploids are reproductively more stable than diploids
- (3) Mutagens penetrate in haploids more effectively than diploids
- (4) Haploids are more abundant in nature than diploids

121. Which one of the following pairs of nitrogenous bases of nucleic acids, is wrongly matched with the category mentioned against it?

- (1) Adenine, Thymine - Purines
- (2) Thymine, Uracil - Pyrimidines
- (3) Uracil, Cytosine - Pyrimidines
- (4) Guanine, Adenine - Purines

122. Which one of the following conditions in humans is correctly matched with its chromosomal abnormality/ linkage?

- (1) Down syndrome - 44 autosomes + XO
- (2) Klinefelter syndrome - 44 autosomes + XXY
- (3) Colour blindness - Y-linked
- (4) Erythroblastosis foetalis - X-linked

123. In the DNA molecules

- (1) There are two strands which run antiparallel-one in 5' → 3' direction and other in 3' → 5'
- (2) The total amount of purine nucleotides and pyrimidine nucleotides is not always equal
- (3) There are two strands which run parallel in the 5' → 3' direction
- (4) The proportion of Adenine in relation to thymine varies with the organism

124. What is true about the isolated small tribal populations?

- (1) There is no change in population size as they have a large gene pool
- (2) There is a decline in population as boys marry girls only from their own tribe
- (3) Hereditary diseases like colour blindness do not spread in the isolated population
- (4) Wrestlers who develop strong body muscles in their life time pass this character on to their progeny

125. Which one of the following scientist's name is correctly matched with the theory put forth by him?

- (1) Mendel - Theory of Pangenesis
- (2) Weismann - Theory of continuity of Germplasm
- (3) Pasteur - Inheritance of acquired characters
- (4) de Vries - Natural selection

126. Which one of the following is incorrect about the characteristics of protobionts (coacervates and micropheres) as envisaged in the abiogenic origin of life?

- (1) They could maintain an internal environment
- (2) They were able to reproduce
- (3) They could separate combinations of molecules from the surroundings
- (4) They were partially isolated from the surroundings

127. Darwin's Finches are an excellent example of

- (1) Connecting links
- (2) Adaptive radiation
- (3) Seasonal migration
- (4) Brood parasitism

128. Which one of the following pairs of items correctly belongs to the category of organs mentioned against it?

- (1) Wings of honey-bee and wings of crow - Homologous organs
- (2) Thorn of Bougainvillea and tendrils of Cucurbita - Analogous organs
- (3) Nictitating membrane and blind spot in human eye - Vestigial organs
- (4) Nephridia of earthworm and malpighian tubules of cockroach - Excretory organs

129. The fruit is chambered, developed from inferior ovary and has seeds with succulent testa in

- (1) Cucumber
- (2) Pomegranate
- (3) Orange
- (4) Guava

130. The  $C_4$  plants are photosynthetically more efficient than  $C_3$  plants because

- (1) They have more chloroplasts
- (2) The  $CO_2$  compensation point is more
- (3)  $CO_2$  generated during photorespiration is trapped and recycled through PEP carboxylase
- (4) The  $CO_2$  efflux is not prevented

131. The chemiosmotic coupling hypothesis of oxidative phosphorylation proposes that adenosine triphosphate (ATP) is formed because

- (1) There is a change in the permeability of the inner mitochondrial membrane toward adenosine diphosphate (ADP)
- (2) High energy bonds are formed in mitochondrial proteins
- (3) ADP is pumped out of the matrix into the intermembrane space
- (4) A proton gradient forms across the inner membrane

132. Dry indehiscent single-seeded fruit formed bicarpellary syncarpous inferior ovary is

- (1) Cremocarp
- (2) Caryopsis
- (3) Cypsela
- (4) Berry

133. The rupture and fractionation do not usually occur in the water column in vessel/tracheids during the ascent of sap because of

- (1) Transpiration pull
- (2) Lignified thick walls
- (3) Cohesion and adhesion
- (4) Weak gravitational pull

134. Senescence as an active developmental cellular process in the growth and functioning of a flowering plant, is indicated in

- (1) Floral parts
- (2) Vessels and tracheid differentiation
- (3) Leaf abscission
- (4) Annual plants

135. Vascular tissues in flowering plants develop from

- (1) Dermatogen
- (2) Phellogen
- (3) Plerome
- (4) Periblem

136. In leaves of  $C_4$  plants malic acid synthesis during  $CO_2$  fixation occurs in

- (1) Guard cells
- (2) Epidermal cells
- (3) Mesophyll cells
- (4) Bundle sheath

137. Importance of day length in flowering of plants was first shown in

- (1) Petunia
- (2) Lemna
- (3) Tobacco
- (4) Cotton

138. Endosperm is consumed by developing embryo in the seed of

- (1) Maize
- (2) Coconut
- (3) Castor
- (4) Pea

139. Nitrogen fixation in root nodules of *Alnus* is brought about by

- (1) Azorhizobium
- (2) Bradyrhizobium
- (3) Clostridium
- (4) Frankia

140. The energy releasing process in which the substrate is oxidised without an external electron acceptor is called

- (1) Glycolysis
- (2) Fermentation
- (3) Photorespiration
- (4) Aerobic respiration

141. Replum is present in the ovary of flower of

- (1) Pea
- (2) Lemon
- (3) Mustard
- (4) Sunflower

142. The fleshy receptacle of syconus of fig encloses a number of

- (1) Mericarps
- (2) Achenes

- (3) Samaras
- (4) Berries

143. Electrons from excited chlorophyll molecule of photosystem II are accepted first by

- (1) Ferredoxin
- (2) Cytochrome - b
- (3) Cytochrome - f
- (4) Quinone

144. Which type of white blood cells are concerned with the release of histamine and the natural anticoagulant heparin?

- (1) Monocytes
- (2) Neutrophils
- (3) Basophils
- (4) Eosinophils

145. Which one of the following is the true description about an animal concerned?

- (1) Cockroach - 10 pairs of spiracles (2 pairs on thorax and 8 pairs on abdomen)
- (2) Earthworm - The alimentary canal consists of a sequence of pharynx, oesophagus, stomach, gizzard and intestine
- (3) Frog - Body divisible into three regions - head, neck and trunk
- (4) Rat - Left kidney is slightly higher in position than the right one

146. Which one of the following is the correct matching of the site of action on the given substrate, the enzyme acting upon it and the end product?

- (1) *Stomach* : Fats  $\xrightarrow{\text{Lipase}}$  micelles
- (2) *Duodenum* : Triglycerides  $\xrightarrow{\text{Trypsin}}$  monoglycerides
- (3) *Small intestine* : Starch  $\xrightarrow{\alpha \text{ Amylase}}$  Disaccharide (Maltose)
- (4) *Small intestine* : Proteins  $\xrightarrow{\text{Pepsin}}$  Amino acids

147. What is vital capacity of our lungs?

- (1) Total lungs capacity minus residual volume
- (2) Inspiratory reserve volume plus tidal volume
- (3) Total lungs capacity minus expiratory reserve volume
- (4) Inspiratory reserve volume plus expiratory reserve volume

148. Which one of the following is the correct difference between Rod Cells and Cone Cells of our retina?

	Rod Cells	Cone Cells
--	-----------	------------

(1) Distribution	More concentrated in centre of retina	Evenly distributed all over retina
(2) Visual acuity	High	Low
(3) Visual pigment contained	Iodopsin	Rhodopsin
(4) Overall function	Vision in poor light	Colour vision and detailed vision in bright light

149. Which one of the following items gives its correct total number?

- (1) Cervical vertebrae in humans - 8
- (2) Floating ribs in humans - 4
- (3) Amino acids found in proteins - 16
- (4) Types of diabetes - 3

150. Given below is a diagrammatic cross section of a single loop of human cochlea



Which one of the following options correctly represents the names of three different parts?

- (1) A : Perilymph, B : Tectorial membrane, C : Endolymph
- (2) B : Tectorial membrane, C : Perilymph, D : Secretory cells
- (3) C : Endolymph, D : Sensory hair cells, A : Serum
- (4) D : Sensory hair cells, A : Endolymph, B : Tectorial membrane

151. Given below are four methods (A - D) and their modes of action (a - d) in achieving contraception. Select their correct matching from the four options that follow

Method	Mode of Action
(A) The pill	(a) Prevents sperms reaching cervix
(B) Condom	(b) Prevents implantation
(C) Vasectomy	(c) Prevents ovulation
(D) Copper T	(d) Semen contains no sperms

Matching

- (1) A-(b), B-(c), C-(a), D-(d)
- (2) A-(c), B-(a), C-(d), D-(b)
- (3) A-(d), B-(a), C-(b), D-(c)
- (4) A-(c) B-(d), C-(a), D-(b)

152. What will happen if the secretion of parietal cells of gastric glands is blocked with an inhibitor?

- (1) Enterokinase will not be released from the duodenal mucosa and so trypsinogen is not converted to trypsin
- (2) Gastric juice will be deficient in chymosin
- (3) Gastric juice will be deficient in pepsinogen
- (4) In the absence of HCl secretion, inactive pepsinogen is not converted into the active enzyme pepsin.

153. During the propagation of a nerve impulse, the action potential results from the movement of

- (1)  $\text{Na}^+$  ions from extracellular fluid to intracellular fluid
- (2)  $\text{K}^+$  ions from extracellular fluid to intracellular fluid
- (3)  $\text{Na}^+$  ions from intracellular fluid to extracellular fluid
- (4)  $\text{K}^+$  ions from intracellular fluid to extracellular fluid

154. The blood calcium level is lowered by the deficiency of

- (1) Calcitonin
- (2) Parathormone
- (3) Thyroxine
- (4) Both Calcitonin and Parathormone

155. The most active phagocytic white blood cells are

- (1) Neutrophils and monocytes
- (2) Neutrophils and eosinophils
- (3) Lymphocytes and macrophages
- (4) Eosinophils and lymphocytes

156. Earthworms have no skeleton but during burrowing, the anterior end becomes turgid and acts as a hydraulic skeleton. It is due to

- (1) Setae
- (2) Coelomic fluid
- (3) Blood
- (4) Gut peristalsis

157. In humans, blood passes from the post caval to the diastolic right atrium of heart due to

- (1) Pressure difference between the post caval and atrium
- (2) Pushing open of the venous valves
- (3) Suction pull
- (4) Stimulation of the sino auricular node



158. In humans, at the end of the first meiotic division, the male germ cells differentiate into the

- (1) Spermatozonia
- (2) Primary spermatocytes
- (3) Secondary spermatocytes
- (4) Spermatids

159. Which one of the following is resistant to enzyme action?

- (1) Leaf cuticle
- (2) Cork
- (3) Wood fibre
- (4) Pollen exine

160. The length of different internodes in a culm of sugarcane is variable because of

- (1) Intercalary meristem
- (2) Shoot apical meristem
- (3) Position of axillary buds
- (4) Size of leaf lamina at the node below each internode

161. Which one of the following pairs of plant structures has haploid number of chromosomes?

- (1) Egg nucleus and secondary nucleus
- (2) Megaspore mother cell and antipodal cells
- (3) Egg cell and antipodal cells
- (4) Nucellus and antipodal cells

162. What does the filiform apparatus do at the entrance into ovule?

- (1) It guides pollen tube from a synergid to egg
- (2) It helps in the entry of pollen tube into a synergid
- (3) It prevents entry of more than one pollen tube into the embryo sac
- (4) It brings about opening of the pollen tube

163. Unisexuality of flowers prevents

- (1) Autogamy and geitonogamy
- (2) Autogamy, but not geitonogamy
- (3) Both geitonogamy and xenogamy
- (4) Geitonogamy, but not xenogamy

164. Which extraembryonic membrane in humans prevents desiccation of the embryo inside the uterus?

- (1) Amnion
- (2) Chorion
- (3) Allantois

(4) Yolk sac

165. Which one of the following statements is incorrect about menstruation?

- (1) The beginning of the cycle of menstruation is called menarche
- (2) During normal menstruation about 40 ml blood is lost
- (3) The menstrual fluid can easily clot
- (4) At menopause in the female, there is especially abrupt increase in gonadotropic hormones

166. The haemoglobin of a human foetus

- (1) Has a higher affinity for oxygen than that of an adult
- (2) Has a lower affinity for oxygen than that of the adult
- (3) Its affinity for oxygen is the same as that of an adult
- (4) Has only 2 protein subunits instead of 4

167. Consider the statements given below regarding contraception and answer as directed thereafter

- (a) Medical Termination of Pregnancy (MTP) during first trimester is generally safe
- (b) Generally chances of conception are nil until mother breast-feeds the infant upto two years
- (c) Intrauterine devices like copper-T are effective contraceptives
- (d) Contraception pills may be taken upto one week after coitus to prevent conception

Which two of the above statements are correct?

- (1) a, b
- (2) b, c
- (3) c, d
- (4) a, c

168. In human adult females oxytocin

- (1) Causes strong uterine contractions during parturition
- (2) Is secreted by anterior pituitary
- (3) Stimulates growth of mammary glands
- (4) Stimulates pituitary to secrete vasopressin

169. Which one of the following is the correct percentage of the two (out of the total of 4) green house gases that contribute to the total global warming?

- (1) Methane 20%,  $N_2O$  18%
- (2) CFCs 14%, Methane 20%
- (3)  $CO_2$  40%, CFSs 30%
- (4)  $N_2O$  6%,  $CO_2$  86%

170. Quercus species are the dominant component in

- (1) Tropical rain forests
- (2) Temperate deciduous forests
- (3) Alpine forests
- (4) Scrub forests

171. About 70% of total global carbon is found in

- (1) Forests
- (2) Grasslands
- (3) Agroecosystems
- (4) Oceans

172. Which one of the following is not observed in biodiversity hotspots?

- (1) Species richness
- (2) Endemism
- (3) Accelerated species loss
- (4) Lesser inter-specific competition

173. World summit on Sustainable Development (2002) was held in

- (1) South Africa
- (2) Brazil
- (3) Sweden
- (4) Argentina

174. The slow rate of decomposition of fallen logs in nature is due to their

- (1) Low cellulose content
- (2) Low moisture content
- (3) Poor nitrogen content
- (4) Anaerobic environment around them

175. Consider the following statements concerning food chains

- (a) Removal of 80% tigers from an area resulted in greatly increased growth of vegetation
- (b) Removal of most of the carnivores resulted in an increased population of deers
- (c) The length of food chains is generally limited to 3-4 trophic levels due to energy loss
- (d) The length of food chains may vary from 2 to 8 trophic levels.

Which two of the above statements are correct?

- (1) a, b
- (2) b, c
- (3) c, d
- (4) a, d

176. According to Central Pollution Control Board (CPCB), which particulate size in diameter (in micrometers) of

the air pollutants is responsible for greatest harm to human health?

- (1) 5.2 - 2.5
- (2) 2.5 or less
- (3) 1.5 or less
- (4) 1.0 or less

177. The table below gives the populations (in thousands) of ten species (A - J) in four areas (a - d) consisting of the number of habitats given within brackets against each. Study the table and answer the question which follows

Area and Number of habitats	Species, and their populations (in thousands) in the areas									
	A	B	C	D	E	F	G	H	I	J
a (11)	2.3	1.2	0.52	6.0	—	3.1	1.1	9.0	—	10.3
b (11)	10.2	—	0.62	—	1.5	3.0	—	8.2	1.1	11.2
c (13)	11.3	0.9	0.48	2.4	1.4	4.2	0.8	8.4	2.2	4.1
d (12)	3.2	10.2	11.1	4.8	0.4	3.3	0.8	7.3	11.3	2.1

Which area out of a to d shows maximum species diversity?

- (1) a
- (2) b
- (3) c
- (4) d

178. A lake near a village suffered heavy mortality of fishes within a few days. Consider the following reasons for this

- (a) Lots of urea and phosphate fertilizer were used in the crops in the vicinity
- (b) The area was sprayed with DDT by an aircraft
- (c) The lake water turned green and stinky
- (d) Phytoplankton populations in the lake declined initially thereby greatly reducing photosynthesis

Which two of the above were the main causes of fish mortality in the lake?

- (1) a, b
- (2) b, c
- (3) c, d
- (4) a, c

179. Consider the following four statements (a - d) about certain desert animals such as kangaroo rat.

- (a) They have dark colour and high rate of reproduction and excrete solid urine
- (b) They do not drink water, breathe at a slow rate to conserve water and have their body covered with thick hairs
- (c) They feed on dry seeds and do not require drinking water
- (d) They excrete very concentrated urine and do not use water to regulate body

temperature

Which two of the above statements for such animals are true?

- (1) a and b
- (2) c and d
- (3) b and c
- (4) c and a

180. A transgenic food crop which may help in solving the problem of night blindness in developing countries is

- (1) Golden rice
- (2) Flavr Savr tomatoes
- (3) Starlink maize
- (4) Bt Soybean

181. Bacterial leaf blight of rice is caused by a species of

- (1) Erwinia
- (2) Xanthomonas
- (3) Pseudomonas
- (4) Alternaria

182. Which one of the following is linked to the discovery of Bordeaux mixture as a popular fungicide?

- (1) Black rust of wheat
- (2) Bacterial leaf blight of rice
- (3) Downy mildew of grapes
- (4) Loose smut of wheat

183. Which one of the following is being tried in India as a biofuel substitute for fossil fuels?

- (1) Aegilops
- (2) Jatropha
- (3) Azadirachta
- (4) Musa

184. Trichoderma harzianum has proved a useful microorganism for

- (1) Biological control of soil-borne plant pathogens
- (2) Bioremediation of contaminated soils
- (3) Reclamation of wastelands
- (4) Gene transfer in higher plants

185. Gel electrophoresis is used for

- (1) Isolation of DNA molecule
- (2) Cutting of DNA into fragments
- (3) Separation of DNA fragments according to their size
- (4) Construction of recombinant DNA by joining with cloning vectors

186. To which type of barriers under innate immunity, do the saliva in the mouth and the tears from the eyes, belong?

- (1) Physical barriers
- (2) Cytokine barriers
- (3) Cellular barriers
- (4) Physiological barriers

187. Match the disease in Column I with the appropriate items (pathogen/prevention/treatment) in Column II

Column I		Column II
(a) Amoebiasis	(i)	Treponema palladium
(b) Diphtheria	(ii)	Use only sterilized food and water
(c) Cholera	(iii)	DPT Vaccine
(d) Syphilis	(iv)	Use oral rehydration therapy

- (1) a-(ii), b-(iii), c-(iv), d-(i)
- (2) a-(i), b-(ii), c-(iii), d-(iv)
- (3) a-(ii), b-(iv), c-(i), d-(iii)
- (4) a-(ii), b-(i), c-(iii), d-(iv)

188. Consider the following statements about biomedical technologies

- (a) During open heart surgery blood is circulated in the heart-lung machine
- (b) Blockage in coronary arteries is removed by angiography
- (c) Computerised Axial Tomography (CAT) shows detailed internal structure as seen in a section of body
- (d) X-ray provides clear and detailed images of organs like prostate glands and lungs

Which two of the above statements are correct?

- (1) a and b
- (2) b and d
- (3) c and d
- (4) a and c

189. Which one of the following pairs of codons is correctly matched with their function or the signal for the particular amino acid?

- (1) UUA, UCA - Leucine

- (2) GUU, GCU - Alanine
- (3) UAG, UGA - Stop
- (4) AUG, ACG - Start / Methionine

190. Which one of the following is the correct statement regarding the particular psychotropic drug specified?

- (1) Barbiturates cause relaxation and temporary euphoria
- (2) Hashish causes after thought perceptions and hallucinations
- (3) Opium stimulates nervous system and causes hallucinations
- (4) Morphine leads to delusions and disturbed emotions

191. Cry 1 endotoxins obtained from *Bacillus Thuringiensis* are effective against

- (1) Boll worms
- (2) Mosquitoes
- (3) Flies
- (4) Nematodes

192. Modern detergents contain enzyme preparations of

- (1) Thermophiles
- (2) Acidophiles
- (3) Alkaliphiles
- (4) Thermoacidophiles

193. The linking of antibiotic resistance gene with the plasmid vector became possible with

- (1) Exonucleases
- (2) DNA ligase
- (3) Endonucleases
- (4) DNA polymerase

194. Which one of the following proved effective for biological control of nematodal disease in plants?

- (1) *Paecilomyces lilacinus*
- (2) *Pisolithus tinctorius*
- (3) *Pseudomonas cepacia*
- (4) *Gliocladium virens*

195. Main objective of production/use of herbicide resistant GM crops is to

- (1) Reduce herbicide accumulation in food articles for health safety
- (2) Eliminate weeds from the field without the use of manual labour
- (3) eliminate weeds from the field without the use of herbicides
- (4) Encourage eco-friendly herbicides

196. Consider the following four measures (a - d) that could be taken to

successfully grow chick-pea in an area where bacterial blight disease is common

- (a) Spray with Bordeaux mixture
- (b) Control of the insect vector of the disease pathogen
- (c) Use of only disease-free seeds
- (d) Use of varieties resistant to the disease

Which two of the above measures can control the disease?

- (1) (a) and (d)
- (2) (b) and (c)
- (3) (a) and (b)
- (4) (c) and (d)

197. Human insulin is being commercially produced from a transgenic species of

- (1) *Saccharomyces*
- (2) *Escherichia*
- (3) *Mycobacterium*
- (4) *Rhizobium*

198. Cornea transplant in humans is almost never rejected. This is because

- (1) It is a non-living layer
- (2) Its cells are least penetrable by bacteria
- (3) It has no blood supply
- (4) It is composed of enucleated cells

199. Which of the following pairs of organs includes only the endocrine glands?

- (1) Adrenal and Ovary
- (2) Parathyroid and Adrenal
- (3) Pancreas and Parathyroid
- (4) Thymus and Testes

200. What is antisense technology?

- (1) RNA polymerase producing DNA
- (2) A cell displaying a foreign antigen used for synthesis of antigens
- (3) Production of somaclonal variants in tissue cultures
- (4) When a piece of RNA that is complementary in sequence is used to stop expression of a specific gene