		[3861]-152					
		F. E. (Semester - I) Examination - 2010					
		APPLIED SCIENCE - I					
(CHEMISTRY)							
		(2008 Pattern)					
	Time: 2 Hours] [Max. Marks: 50						
Instru	uctions						
		(1) Answer 3 questions.					
		(2) Black figures to the right indicate full marks.					
		(3) Neat diagrams must be drawn wherever necessary.(4) Use of logarithmic tables, slide rale Mollier charts, electron	:				
		pocket calculator and steam in bles is allowed.	<i>ic</i>				
Q.1)	(A)	(a) Define Axis of Symmetry for Crystals. Draw diagrams showing two-fold, three fold and four-fold axes of symmetry for Cubic Sepstals.	41				
		(b) Give difference in Schottky and Frenkel point defects in Ionic Crysta's. [04]	-				
	(B)	Derive Bragg's Law Equation for Crystals acting as reflection grating.	5]				
	(C)	What is Atomic Packing Factor (APF) for Crystals? Calculate APF for Closest Packing Cubic Crystal System. [04]	4]				
		OR					
Q.2)	(A)	Give Bravais Lattices for Cubic, Orthorhombic and Hexagonal Crystal Systems. Draw diagrams for Bravais Lattices of Cubic Crystal Systems.	7]				
	(B)	What is Luminescence? Describe structure of ZnS shortly. How is Luminescence of ZnS increased? Give uses of ZnS as Phosphor. [0:	5]				
	(C)	Give structure and properties of Tetrathiofulvalene. How is Tetrathiofulvalene converted to stable charge-transfer compound? Give uses of such compound. [0:	5]				
[3861]-152	1 P.T.O).				

Total No. of Questions: 6] [Total No. of Printed Pages: 3

- Q.3) (A) Define a Primary Standard. What are the conditions for a chemical to be the Primary Standard? [04]
 - (b) 50 ml of a Chloride Water Sample requires 5.7 ml of M/100 AgNO₃ in Mohr's Method of Precipitation Titration. Find amount of Chloride Ions per litre in the Water. (Ionic Weight of Cl⁻ = 35.5)
 - (B) Explain weak acid strong base titration regarding pet reaction, titration curve and suitable indicators. Give formula for calculation of pH of titration mixture at the stage of equivalence point.

 (Assume strong base solution in burette.) [06]
 - (C) Calculate equivalent weight of KMnO₄ oxidising reagent in acidic medium. (Atomic Weights: K) 39, Mn = 55, O = 16) [03]

OR

- Q.4) (A) (a) Show that colour change pH interval of an acid-base titration indicator is about 2 pH-units. [04]
 - (b) Balance any one of the following redox reactions by ion-electron or nascent oxygen method: [03]
 - (i) $KMnO_4 + H_2SO_4 + FeSO_4 \rightarrow MnSO_4 + Fe_2(SO_4)_3 + H_2O$
 - (ii) $Fe^{+2} + Sn^{+4} \rightarrow Fe^{+3} + Sn^{+2}$
 - (B) 25 ml 0.1N ACL is titrated against 0.075N NaOH from burette.

 Calculate pH of titration mixture at following stages: [05]
 - (a) 10 ml NaOH added.
 - (b) 10 ml NaOH added.
 - (C) Explain role of any two indicators for Precipitation Titrations. [04]

Q.5)	(A)	Give polymerisation reaction, properties and uses of any tw of the following polymers:	o [08]
		(a) S.B.R.	
	•	(b) ABS Plastic	
		(c) Epoxy Resin	
		(d) Polystyrene	
	(B)	Give a note on compounding of polymers regarding:	[05]
		(a) Plasticizers	. ,
		(b) Fillers	
		(c) Stabilizers	
	(C)	Compare Natural Rubber and Vulcanised Natural Rubber.	[04]
Q.6)	(A)	(a) Distinguish between LOPE and HDPE.	[04]
		(b) Classify Co-polymers	[03]
	(B)	Write a short note one of the following:	[05]
		(a) Biodegradable Polymers	[oc]
		(b) Conducting Polymers	
	(C)	Give Free Radical Mechanism of Chain Polymerisation.	[05]
		• ———	
	•	7,	
	1	\	