

Chemistry Paper
2007

IMPORTANT NOTES FOR CANDIDATES


- Attempt ALL the 44 questions.
- Questions 1-30 (objective questions) carry *three* marks each and questions 31-44 (subjective questions) carry *fifteen* marks each.
- Write the answers to the objective questions in the Answer Table for Objective Questions provided on page 13 only.

1. The compound, which

- (i) reacts rapidly with acetyl chloride,
- (ii) does not react with 2,4-dinitrophenylhydrazine **and**
- (iii) does not form a yellow precipitate with excess of iodine in aqueous alkali is

- (A) acetone
- (B) diethyl ether
- (C) 2-methyl-2-propanol
- (D) ethanol


2.



The given compounds **1** and **2** are

- (A) identical
- (B) diastereomeric
- (C) enantiomeric
- (D) constitutionally isomeric

3. The correct order of dipole moments (μ) of the following compounds is



(A) $\mu_1 > \mu_2 > \mu_3$

(B) $\mu_2 > \mu_3 > \mu_1$


(C) $\mu_3 > \mu_1 > \mu_2$

(D) $\mu_2 > \mu_1 > \mu_3$

4. Which one of the following compounds gives positive test for both nitrogen and halogen with its Lassaigne's extract?



5. Which one of the following compounds is optically active?







(A) 1

(B) 2

(C) 3

(D) 4

6. The compounds that react with aqueous NaHCO_3 to release CO_2 are




- (A) 1 and 3
- (B) 2 and 4
- (C) 2 and 3
- (D) 1 and 4

7. The complementary strand of DNA for the following single stranded DNA sequence, $5' \text{-A-T-C-A-T-G-C-3}'$ is


- (A) $5' \text{-A-T-C-A-T-G-C-3}'$
- (B) $5' \text{-T-A-G-T-A-C-G-3}'$
- (C) $5' \text{-G-C-A-T-G-A-T-3}'$
- (D) $5' \text{-C-G-T-A-C-T-A-3}'$

8. The value of 'n' for the following molecule according to Hückel's rule is








- (A) 16
- (B) 4
- (C) 3
- (D) 14


9. Which one of the following compounds reacts with nitrous acid to give the product [P]?



10. The main product obtained in the following reaction is




11. For a reaction with rate equation $-dC/dt = kC^2$, C_0 and C are the concentrations of the reactant at time 0 and t respectively. If 10 minutes were required for C_0 to become $C_0/2$, the time required for C_0 to become $C_0/4$ is
- (A) 10 min
 (B) 20 min
 (C) 30 min
 (D) 40 min
12. For a cyclic process performed by an ideal gas, changes in some thermodynamic functions are zero. Indicate the set in which all the functions are zero.
- (A) $w, \Delta E, \Delta H, \Delta G$
 (B) $q, \Delta S, \Delta H, \Delta A$
 (C) $q, \Delta E, \Delta S, \Delta G$
 (D) $\Delta E, \Delta S, \Delta H, \Delta A$
13. The plot of Gibb's free energy G and the extent of a reaction ξ is given below for the reaction $A \rightleftharpoons B$. If μ_A and μ_B are the chemical potentials of A and B respectively, the **INCORRECT** statement is







- (A) at point X, $\mu_A > \mu_B$
 (B) at point Y, $\Delta G = 0$
 (C) at point Z, $\mu_A > \mu_B$
 (D) at equilibrium, the composition of the reaction mixture can be identified

14. The overlap between the atomic orbitals sketched below is



- (A) positive
 (B) negative
 (C) zero
 (D) no overlap
15. The pH of a 1.0×10^{-3} M solution of a weak acid HA is 4.0. The acid dissociation constant K_a is
- (A) 1.0×10^{-3}
 (B) 1.0×10^{-4}
 (C) 1.0×10^{-5}
 (D) 2.0×10^{-5}
16. The normalisation constant 'A' for the wavefunction $\psi(\phi) = Ae^{(im\phi)}$ where $0 \leq \phi \leq 2\pi$ is
- (A) $1/\sqrt{2\pi}$
 (B) $\sqrt{2\pi}$
 (C) 2π
 (D) $1/\sqrt{2}$

17. The standard potential of a Daniel cell is + 1.10 V and the equilibrium constant for the cell reaction is 1.5×10^{37} . It can be concluded that
- (A) zinc oxidises copper
 - (B) displacement of copper by zinc goes to near completion
 - (C) copper oxidises zinc
 - (D) displacement of zinc by copper goes to completion
18. Which one of the following figures, showing kinetic energy of the ejected electron *versus* the frequency (ν) of the incident photon, represents the Einstein's photoelectric effect?




19. An aqueous solution containing 0.01 M FeCl_3 and 0.06 M HClO_4 has the same ionic strength as a solution of
- (A) 0.09 M NaCl
 - (B) 0.04 M Na_2SO_4
 - (C) 0.06 M CuSO_4
 - (D) 0.03 M H_3PO_4

20. Which one of the following species is the conjugate base of HO^- ?

- (A) H_2O
- (B) O^{2-}
- (C) O_2^-
- (D) O_2^{2-}

21. The solid-liquid phase diagram for the Mg-Zn system is shown in the figure below where the vertical line at $x(\text{Mg}) = 0.33$ represents the formation of a congruent melting compound MgZn_2 . The figure is divided into seven regions depending upon the physical state of the system. The composition of the region #6 represents



- (A) single phase of a solution of Mg and Zn
 - (B) two phase region between the solid Zn and solid MgZn_2
 - (C) two phase region between the liquid and solid MgZn_2
 - (D) two phase region between solid Mg and solid MgZn_2
22. In the extraction of metals from their ores, which one of the following reduction methods can bring about a non-spontaneous reduction?
- (A) electrolytic reduction
 - (B) reduction by carbon
 - (C) reduction by another metal
 - (D) reduction by hydrogen

23. The correct order of the ionic radii is
- (A) $\text{In}^{3+} > \text{Sn}^{4+} > \text{Sr}^{2+} > \text{Rb}^+$
 - (B) $\text{Sn}^{4+} > \text{In}^{3+} > \text{Sr}^{2+} > \text{Rb}^+$
 - (C) $\text{Rb}^+ > \text{In}^{3+} > \text{Sr}^{2+} > \text{Sn}^{4+}$
 - (D) $\text{Rb}^+ > \text{Sr}^{2+} > \text{In}^{3+} > \text{Sn}^{4+}$
24. The correct valence shell electronic configuration of the element with atomic number 22 is
- (A) $[\text{Ar}]4s^23d^2$
 - (B) $[\text{Ar}]3d^4$
 - (C) $[\text{Ar}]3d^24s^2$
 - (D) $[\text{Ar}]4s^24p^2$
25. The ligand with only sigma (σ) bonding character is
- (A) CN^-
 - (B) CH_3^-
 - (C) CO
 - (D) NO
26. Which one of the following species is **NOT** isoelectronic with CO ?
- (A) N_2
 - (B) CN^-
 - (C) NO^+
 - (D) O_2^+
27. During Wittig reaction, a phosphorus ylide gets converted to
- (A) R_3P
 - (B) $\text{R}_3\text{P}=\text{O}$
 - (C) $\text{R}_3\text{P}^+\text{HOH}^-$
 - (D) $\text{R}_2\text{P}-\text{PR}_2$

28. Which of the following reactions does **NOT** give H_3PO_4 ?



29. The ionic radii of Ca^{2+} and F^- are 100 pm and 133 pm respectively. The coordination number of Ca^{2+} in the ionic solid will be

(A) 8

(B) 6

(C) 4

(D) 2

30. The shape of CH_3^- ion is


(A) trigonal planar

(B) tetrahedral

(C) trigonal pyramidal


(D) linear

31.



Identify the structures of the intermediate compounds **Q**, **R** and **S**. Show the transformation for each step. (15)



32. (a) For the following scheme of transformations, draw the structures of **A**, **B**, **C** and **D**.




(9)

- (b) Complete hydrolysis of a pentapeptide with 6 N HCl at 110 °C in a sealed tube gave 2 equivalents of glycine, one equivalent each of tyrosine, leucine and phenylalanine. Reaction of the pentapeptide with Sanger's reagent (2,4-dinitrofluorobenzene, DNFB) and subsequent hydrolysis gave the DNFB derivative of tyrosine. Chymotrypsin cleavage of this peptide yielded tyrosine, leucine and a tripeptide. Deduce the sequence of the pentapeptide. (6)

33. Complete the following reactions with appropriate structures for **E**, **F**, **G**, **H** and **I**.




(9)




(6)

34. (a) Account for the following transformation with an appropriate mechanism. Give the structure of the Hofmann exhaustive methylation product of 1,2-dihydro derivative of [X].



- (b) The optically pure ester [J] is hydrolysed in aqueous acetic acid to form a racemic mixture of *cis*-4,4-dimethyl-2-acetoxycyclopentanol [K]. Give a mechanistic explanation to account for the formation of [K] and the observed change in the optical activity.



35. (a) **M** is a first row transition metal. MCl_2 on treatment with aqueous ammonia gives a blue colored solution of complex **N**. A solution of MCl_2 also gives a bright red precipitate of complex **O** with ethanolic dimethylglyoxime.
- (i) Identify **M** and draw the structure of **O**.
 - (ii) Determine the hybridisation of **M** in complex **N**.
 - (iii) Identify the paramagnetic complex. (9)
- (b) $[Cr(H_2O)_6]^{3+}$ gave an absorption at 208 kJ/mol which corresponds to Δ_o . Calculate the crystal field stabilisation energy of this complex in kJ/mol. (6)

36. (a) Consider the ethers $\text{H}_3\text{SiOSiH}_3$ and H_3COCH_3 .


(i) Which ether has more Lewis base character?

(ii) Which angle [Si-O-Si and C-O-C] is greater?

Justify your answer.

(9)

(b) Starting from SiO_2 , show how the following polymer is prepared industrially?



(6)

37. (a) A solution of metal ion (M^{2+}) when treated with H_2S gas gives a black precipitate **A**. Precipitate **A** dissolves in hot concentrated nitric acid to give **B** along with elemental sulfur. The metal ion solution also gives a white precipitate **C** with an excess of KI . Write the chemical formulae of **A**, **B** and **C**. (9)
- (b) Why are potassium permanganate solutions unstable in the presence of Mn^{2+} ions? In the quantitative estimation of iron present in iron ores dissolved in dilute HCl , titrations with dichromate are preferred over titrations with permanganate. Rationalise. (6)

38. (a) Al_2Cl_6 and Al_2Me_6 are dimeric in gas phase. Draw their structures. Which compound has more Lewis acid character? Explain. (9)
- (b) Arrange the halides SnCl_2 , PbCl_2 , SiCl_2 in increasing order of their stability. Give reasons for your answer. (6)

39. (a) Acidification of an aqueous solution of yellow sodium chromate gives an orange colored compound **A**. A saturated solution of **A** on treatment with concentrated H_2SO_4 gives a bright orange solid **B**. Compound **A** in the presence of concentrated H_2SO_4 reacts with anion **C** to give a deep red colored liquid. Identify **A**, **B** and **C**. (9)

(b) ${}_{84}^{215}\text{Po}$ undergoes an α emission to give element **X** followed by a β emission to give element **Y**.

(i) Write the valence shell electronic configuration of **Y**.

(ii) Indicate the groups of the periodic table to which **X** and **Y** belong. (6)


40. (a) When an ideal monoatomic gas is expanded from 1.5 bar, 24.8 L and 298 K into an evacuated container, the final volume becomes 49.6 L. Calculate ΔH , ΔS and ΔG for the process. (9)

(b) The Maxwell distribution function for the distribution of speeds of molecules in gaseous systems is given as,

$$f(c) = 4\pi \left(\frac{m}{2\pi kT} \right)^{3/2} c^2 \exp\left(-mc^2/2kT\right)$$

Show that the most probable speed, $c_{mps} = \left(2kT/m\right)^{1/2}$ (6)

41. (a) At 600 K and 200 bar, a 1:3 (molar ratio) mixture of A_2 and B_2 react to form an equilibrium mixture containing $x_{AB_3} = 0.60$. Assuming ideal gas behaviour, calculate K_p for the reaction




- (b) A 50 mL 0.05 M solution of Fe(II) is titrated with 0.05 M solution of Ce(IV) in the presence of dilute H_2SO_4 at 25 °C. Calculate the equivalence point potential and the equilibrium constant K in terms of $\log K$.

$$[E^0(Fe^{3+}/Fe^{2+}) = +0.75 \text{ V}, E^0(Ce^{4+}/Ce^{3+}) = +1.45 \text{ V}] \quad (6)$$

42. (a) The vapour pressure of D_2O at $20\text{ }^\circ\text{C}$ is 745 mm Hg . When 15 g of a non-volatile compound is dissolved in 200 g of D_2O , the pressure changes to 730 mm Hg . Assuming the applicability of Raoult's law, calculate the molecular weight of the compound. (9)
- (b) An enzyme following Michaelis-Menten kinetics was found to have highest activity at $37\text{ }^\circ\text{C}$ and $\text{pH } 7.0$. If the maximum velocity V_{max} for this enzyme was $2.4 \times 10^{-4}\text{ mol L}^{-1}\text{s}^{-1}$ with an initial enzyme concentration $[E]_0 = 2.4\text{ nM}$, calculate the turnover frequency. (6)


43. (a) Consider the 4π electrons in cyclobutadiene to be free particles in a 2-dimensional square box of length 2 \AA . Calculate the wavelength of the electronic transition from the highest occupied molecular orbital (HOMO) to the lowest unoccupied molecular orbital (LUMO). Also write down the normalised wavefunctions for the occupied degenerate states. (9)

(b) The reaction



is first order in both directions. At 25°C , the equilibrium constant (K) of this reaction is 0.40 . If $0.115 \text{ mol.dm}^{-3}$ of *cis*-isomer is allowed to equilibrate, calculate the equilibrium concentration of each isomer. (6)

44. (a) With i, j and k as the unit vectors along X, Y and Z axes, express the vector $\overrightarrow{P_1P_2}$ in the given figure in terms of the coordinates of P_1 and P_2 . Also determine the dot products of the unit vectors i, j, k .



(9)

- (b) Deduce whether the matrices A and B commute or not.

$$A = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

(6)