(Physical and Inorgonic Chemistry) Q. 1. Select and write the most appropriate answer from the given alternatives (8) for each sub-question: (i) The enthalpies of all elements in their standard states are ..... (a) less than zero (b) zero (d) greater than unity (c) unity (ii) A conjugate acid-base pair differs by S.T.I.O.NPAPERS (a) one electron (b) one neutron (c) one proton (d) one electron pair (iii) The unit of electrochemical equivalent is ..... (d) gram coulomb -1 (b) coulomb gram-1 (c) gram amp-1 (a) gram (iv) The minimum amount of fissionable material required to continue the chain reaction is called ......

CHEMISTRY PAPER- I

Question Paper : October 2010

Max. Marks: 40

(d) atomic mass

(d) +890 kJ

(2)

(a) fission mass (b) critical mass (c) active mass (v) In a zero order reaction, the rate of reaction is independent of .......

(a) concentration of products (b) catalyst (c) concentration of reactants (d) temperature

(vi) The chemical formula of willemite is ......

(a) ZnS (b) ZnCO<sub>3</sub> (c) ZnO (d) Zn<sub>2</sub>SiO<sub>4</sub>

(vii) The volume of water to be added to 0.1dm3 of 0.5 N H2SO4 to get decinormal solution is

(a) 0.1 dm<sup>3</sup> (b) 0.4 dm<sup>3</sup> (c) 0.45 dm<sup>3</sup> (d) 0.5 dm<sup>3</sup>

(c) +445 kJ

(viii) The enthalpy change for combustion of methane is -890 kJ / mol. What is the amount of heat liberated on combustion of 8 x 10-3 kg of methane?

(a) -890 kJ (b) -445 kJ Q. 2 (A) Attempt any ONE :

Date: 04-10-2010

Time: 2 Hrs.

(i) Derive an expression for the effect of temperature on the heat of reaction at constant pressure.

WWW.UNIVERSITYQUESTIONPAPERS.COM

<ul><li>(ii) Draw a neat labelled diagram of Berkeley and Hartley method to measure</li></ul>	
- osmotic pressure.	5
	(2)
(i) Define : (a) Solubility product (b) Corrosion.	
<ul><li>(ii) Distinguish between molecularity of reaction and order of reaction.</li></ul>	1
	(4)
(i) Explain the following characteristics of transition elements.	
(a) Tendency to form complexes. (b) Catalytic property	
(ii) Define osmotic pressure. Derive an equation for determination of molecular	
weight from osmotic pressure.	
	(3)
<ul> <li>(i) Describe Landsberger and Walker method for determination of molecular weight from boiling point elevation.</li> </ul>	
(ii) Define isotopes. Explain the use of radio-isotope in carbon dating.	
	(3)
(i) Derive an integrated rate equation for the first order reaction.	ं
(ii) What is the action of following on zinc metal?	
(a) Concentrated HCl (b) Hot solution of NaOH (c) Ammonia	
	(3)
Define electrochemical series. Explain its 'one' application.	10,
	(4)
Q. 4. (A) Answer the following:	(+)
Define: (i) Enthalpy (ii) Molar heat capacity at constant volume.	
Show that $\Delta H = \Delta E + P \Delta V$ , at constant pressure.	
(B) Attempt any ONE :	(4)
(i) Define hydrolysis of a salt. Show that, for the salt of weak acid-weak base	
K	
salt, $K_h = \frac{K_w}{K_a K_h}$	
(ii) Define single electrode potential. Explain Nernst theory of electronation and	
de-electronation with suitable example.	
Q. 5. (A) Attempt any ONE :	(4)
(i) Calculate the heat of formation of sucrose from the following data -	
(1) C, H, O, H, H 12 O, S 12 CO, S 11H, O, PAH = 5834 16 kJ	
(2) $C_{(a)} + O_{2(g)} \rightarrow CO_{2(g)}$ ; $\Delta H_2 = -394.96 \text{ kJ}$	
(3) $H_{2(g)} + \frac{1}{2}O_{2(g)} \rightarrow H_2O_{(f)}$ ; $\Delta H_3 = -286.18 \text{ kJ}$	
(ii) The pH of decimolar solution of NH <sub>4</sub> Cl is 5.1276. Calculate K <sub>h</sub> , h and K <sub>b</sub> for	2.0
same solution.	1
(Given: K <sub></sub> = 1 × 10 <sup>-14</sup> )	
ID) AN THIO	(4)
(i) Calculate mass defect and binding energy per nucleon of 59 Co if the isotopic	1.,
(i) Calculate mass defect and binding energy per nucleon of Con the isotopic	
mass of cobalt is 58.997 a. m. u.	
Given: Mass of proton = 1.0078 a. m. u.	
Mass of neutron = 1.0086 a. m. u.	
(ii) Calculate the amount of electricity required to reduce all silver ions from	
1dm³ of 0.25 M silver nitrate solution.	
(At. Wt. Ag = 108, N = 14 and O = 16. 1F = 96500 C)	
(iii) $1 \times 10^{-3}$ kg of a non-volatile substance, when dissoved in $5.05 \times 10^{-2}$ kg of	
benzene, freezing point of the solvent was lowered by 0.4 K.	
If the freezing point depression constant of benzene is 5.12 K. kg. mol-1,	
calculate molecular mass of the solute.	