

## TEST CODE: MM 2007

### SYLLABUS

Convergence and divergence of sequence and series;  
Cauchy sequence and completeness;  
Bolzano-Weierstrass theorem;  
continuity, uniform continuity, differentiability,  
directional derivatives, Jacobians, Taylor Expansion;  
integral calculus of one variable – existence of Riemann integral,  
Fundamental theorem of calculus, change of variable;  
elementary topological notions for metric space – open, closed and  
compact sets, connectedness;  
elements of ordinary differential equations.

Equivalence relations and partitions;  
vector spaces, subspaces, basis, dimension, direct sum;  
matrices, systems of linear equations, determinants;  
diagonalization, triangular forms;  
linear transformations and their representation as matrices;  
groups, subgroups, quotients, homomorphisms, products,  
Lagrange's theorem, Sylow's theorems;  
rings, ideals, maximal ideals, prime ideals, quotients,  
integral domains, unique factorization domains, polynomial rings;  
fields, algebraic extensions, separable and normal extensions, finite fields.

### SAMPLE QUESTIONS

1. Let  $k[x, y]$  be the polynomial ring in two variables  $x$  and  $y$  over a field  $k$ . Prove that any ideal of the form  $(x - a, y - b)$  for  $a, b \in k$  is a maximal ideal of this ring.
2. Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a linear transformation. Show that there is a line  $L$  such that  $T(L) = L$ .
3. Let  $A \subseteq \mathbb{R}^n$  and  $f : A \rightarrow \mathbb{R}^m$  be a uniformly continuous function. If  $\{x_n\}_{n \geq 1} \subseteq A$  is a Cauchy sequence then show that  $\lim_{n \rightarrow \infty} f(x_n)$  exists.
4. Let  $N > 0$  and let  $f : [0, 1] \rightarrow [0, 1]$  be denoted by  $f(x) = 1$  if  $x = 1/i$  for some integer  $i \leq N$  and  $f(x) = 0$  for all other values of  $x$ . Show that  $f$  is Riemann integrable.

5. Let  $F : \mathbf{R}^n \rightarrow \mathbf{R}$  be defined by

$$F(x_1, x_2, \dots, x_n) = \max\{|x_1|, |x_2|, \dots, |x_n|\}.$$

Show that  $F$  is a uniformly continuous function.

6. Show that every isometry of a compact metric space into itself is onto.
7. Let  $\mathbf{T} = \{z \in \mathbf{C} : |z| = 1\}$  and  $f : [0, 1] \rightarrow \mathbf{C}$  be continuous with  $f(0) = 0, f(1) = 2$ . Show that there exists at least one  $t_0$  in  $[0, 1]$  such that  $f(t_0)$  is in  $\mathbf{T}$ .
8. Let  $f$  be a continuous function on  $[0, 1]$ . Evaluate

$$\lim_{n \rightarrow \infty} \int_0^1 x^n f(x) dx.$$

9. Find the most general curve whose normal at each point passes through  $(0, 0)$ . Find the particular curve through  $(2, 3)$ .
10. Suppose  $f$  is a continuous function on  $\mathbf{R}$  which is periodic with period 1, that is,  $f(x + 1) = f(x)$  for all  $x$ . Show that
- (i) the function  $f$  is bounded above and below,
  - (ii) it achieves both its maximum and minimum and
  - (iii) it is uniformly continuous.
11. Let  $A = (a_{ij})$  be an  $n \times n$  matrix such that  $a_{ij} = 0$  whenever  $i \geq j$ . Prove that  $A^n$  is the zero matrix.
12. Determine the integers  $n$  for which  $\mathbf{Z}_n$ , the set of integers modulo  $n$ , contains elements  $x, y$  so that  $x + y = 2, 2x - 3y = 3$ .
13. Let  $a_1, b_1$  be arbitrary positive real numbers. Define

$$a_{n+1} = \frac{a_n + b_n}{2}, b_{n+1} = \sqrt{a_n b_n}$$

for all  $n \geq 1$ . Show that  $a_n$  and  $b_n$  converge to a common limit.

14. Show that the only field automorphism of  $\mathbf{Q}$  is the identity. Using this prove that the only field automorphism of  $\mathbf{R}$  is the identity.
15. Consider a circle which is tangent to the  $y$ -axis at 0. Show that the slope at any point  $(x, y)$  satisfies  $\frac{dy}{dx} = \frac{y^2 - x^2}{2xy}$ .
16. Consider an  $n \times n$  matrix  $A = (a_{ij})$  with  $a_{12} = 1, a_{ij} = 0 \forall (i, j) \neq (1, 2)$ . Prove that there is no invertible matrix  $P$  such that  $PAP^{-1}$  is a diagonal matrix.

17. Let  $G$  be a nonabelian group of order 39. How many subgroups of order 3 does it have?
18. Let  $n \in \mathbf{N}$ , let  $p$  be a prime number and let  $\mathbf{Z}_{p^n}$  denote the ring of integers modulo  $p^n$  under addition and multiplication modulo  $p^n$ . Let  $f(x)$  and  $g(x)$  be polynomials with coefficients from the ring  $\mathbf{Z}_{p^n}$  such that  $f(x) \cdot g(x) = 0$ . Prove that  $a_i b_j = 0 \forall i, j$  where  $a_i$  and  $b_j$  are the coefficients of  $f$  and  $g$  respectively.
19. Show that the fields  $\mathbf{Q}(\sqrt{2})$  and  $\mathbf{Q}(\sqrt{3})$  are isomorphic as  $\mathbf{Q}$ -vector spaces but not as fields.
20. Suppose  $a_n \geq 0$  and  $\sum a_n$  is convergent. Show that  $\sum 1/(n^2 a_n)$  is divergent.