

This question paper contains 8 printed pages]

Your Roll No

5165

B.Sc. Prog./B.Sc. (Hons.)/I J
M.A. 107-B – MATHEMATICS
(For Life Sciences)
(NC – Admission of 2008 onwards)

Time : 3 Hours

Maximum Marks : 75

(Write your Roll No on the top immediately on receipt of this question paper)

There are three Sections in this question paper

Attempt any **two** questions from each Section.

Students are allowed to use calculators

Section – I

- 1 (a) Consider a spherical cell of volume V and surface S Express V as a function of S . Is it a linear function ? **4**
- (b) A culture of bacteria initially weighs 1 gm and is doubling in size every hour How long will it take to reach a weight of 3 gms **4**

- (c) The weight of a certain stock of fish is given by $W = nw$, where n is the size of the stock and w the average weight of each fish. If n and w change with time t according to the formulas $n = (2t^2 + 3)$ and $w = (t^2 - t + 2)$, find the rate of change of W w.r.t. time t . 4½

- 2 (a) Assume that a population of size 25000 (at time $t = 0$) grows according to the formula $N = 25000 + 45t^2$ where the time t is measured in days. Find the average growth rate in the time intervals from $t = 0$ to $t = 2$. 4½

(b) Find :

(i) $\lim_{h \rightarrow 0} \frac{4 - h}{2 + 7h}$

(ii) $\lim_{h \rightarrow 0} \frac{4 - (2 + h)^2}{1 - (1 - h)^2}$ 4

(c) Show that for Fibonacci numbers

$$a_1 + a_2 + \dots + a_n = a_{n+2} - 1$$
 4

3 (a) Integrate

(i) $\int (3x - 7)^5 dx$

$$(ii) \int \sin(5 - 3x) dx$$

$$(iii) \int \frac{\log x}{x} dx \quad 7\frac{1}{2}$$

- (b) An individual suffering from a certain disease is administered an amount x of a suitable drug. His probability of being cured is $\frac{\sqrt{x}}{3(1+x)}$.

Find the value of x that gives him the maximum probability of being cured. 5

Section - II

- 4 (a) If $A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ -1 & 1 & -1 & 1 \end{pmatrix}$ and $B^T = (a \ b \ c \ d)$ when T stands for Transpose.

Calculate

- (i) $A(B^T)^T$ (ii) $B^T A^T$, and show that

$$(AB)^T = B^T A^T \quad 4$$

- (b) A signal operated by a laboratory mouse has only two faces R = red, Y = yellow At each trial the mouse may or may not change the signal Suppose that the following transition probabilities are given

$$R \longrightarrow R \quad p_{11} = 0.8$$

$$R \longrightarrow Y \quad p_{12} = 0.2$$

$$Y \longrightarrow R \quad p_{21} = 0.6$$

$$Y \longrightarrow Y \quad p_{22} = 0.4$$

Assume further that each trial is independent of past experience. Then the outcomes of each trial form a Markov chain with two states (R and Y) Establish the transition matrix with the above probabilities Also, calculate the probabilities for two-step transitions keeping into mind the fact that under the assumption of Markov chains the multiplication rule holds 4½

(c) If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $B = \begin{pmatrix} 2 & 5 \\ 0 & 4 \end{pmatrix}$, $C = \begin{pmatrix} -1 & -2 \\ 3 & 0 \end{pmatrix}$

Find out $A(B + C)$ in two ways according to the distributive law 4

- 5 (a) If $Q = (x^2 + y^2)^{1/2}$, verify that

$$\frac{\partial^2 Q}{\partial x^2} + \frac{\partial^2 Q}{\partial y^2} = \frac{1}{Q} \quad 4$$

- (b) Some biological rhythms are described by the second order differential equation

$$\frac{d^2 x}{dt^2} + kx = 0 \quad (k > 0)$$

Show that $x = A \cos \omega t + B \sin \omega t$ is the solution of the differential equation where $\omega^2 = k$.

4½

- (c) If $z = ax^2 + 2hxy + by^2$, verify that

$$\frac{\partial^2 z}{\partial y \partial x} = \frac{\partial^2 z}{\partial x \partial y} \quad 4$$

6. (a) Show that $y = \frac{c}{x} + d$ is a solution of the

differential equation $\frac{dy}{dx} + \frac{c}{x^2} = 0$

Further, plot this solution for $c = 1, d = 0$ and $c = -1, d = 0$, take $x > 0$.

6½

- (b) Assume that a population grows in such a way that the specific growth rate $\frac{\dot{1}}{N} \frac{dN}{dt}$ remains constant. Let N_1 be the number of individuals at the time instant t_1 . Find $N = N(t)$ 6

Section – III

- 7 (a) The following are the weights (kg) of the 6 subjects in the sample studied by a scientist
83.9, 99.0, 63.8, 71.3, 65.3, 79.6
Compute the mean and standard deviation 6½
- (b) Suppose that over a period of several years the average number of deaths from a certain non-contagious disease has been 10. If the number of deaths from this disease follows the Poisson distribution, what is the probability that during the current year
- (i) exactly seven people will die from the disease
 - (ii) ten or more people will die from the disease (Given $e^{-10} = 0.000045$) 6

8. (a) The heights of a certain population of individuals are normally distributed with a mean of 70 inches and a standard deviation of 3 inches. What is the probability that a person picked at random from the group will be between 65 and 74 inches tall ?

(Area under the standard normal curve from 0 to 1.33 = 0.4082

Area under the standard normal curve from 0 to 1.67 = 0.4525)

6½

- (b) Find the equations of regression lines for the following values of x and y

x	1	2	3	4	5
y	2	5	3	8	7

Also estimate y for $x = 10$.

6

- 9 (a) In a health survey of school children, the mean haemoglobin level of 55 boys was found to be 10.2 g per 100 ml with a standard deviation 2.1 g. Can it be considered that this group of boys is identified from a population with a mean of 11.0 g / 100 ml -

6

- (b) Hearing levels in two groups of school children with normal hearing in frequency of 500 cycles per second was found as follows

	No of Children	Hearing (\bar{x}) threshold	S D (σ)
Group I	62	15.5 dB	6.5 dB
Group II	76	20 dB	7.1 dB

Test at 5% level of significance if there is any difference between hearing levels recorded in two groups. 6½
