(CHE)

CHEMICAL ENGINEERING INSTRUCTIONS TO CANDIDATES

- Candidates should write their Hall Ticket Number only in the space provided at the top left hand corner of this page, on
 the leaflet attached to this booklet and also in the space provided on the OMR Response Sheet. BESIDES WRITING,
 THE CANDIDATE SHOULD ENSURE THAT THE APPROPRIATE CIRCLES PROVIDED FOR THE
 HALL TICKET NUMBERS ARE SHADED USING H.B. PENCIL ONLY ON THE OMR RESPONSE
 SHEET. DO NOT WRITE HALL TICKET NUMBER ANY WHERE ELSE.
- Immediately on opening this Question Paper Booklet, check:
 - (a) Whether 200 multiple choice questions are printed (50 questions in Mathematics, 25 questions in Physics, 25 questions in Chemistry and 100 questions in Engineering)
 - (b) In case of any discrepancy immediately exchange the Question paper Booklet of same code by bringing the error to the notice of invigilator.
- Use of Calculators, Mathematical Tables and Log books is not permitted.
- Candidate must ensure that he/she has received the Correct Question Booklet, corresponding to his/her branch of Engineering.
- 5. Candidate should ensure that the booklet Code and the Booklet Serial Number, as it appears on this page is entered at the appropriate place on the OMR Response Sheet by shading the appropriate circles provided therein using H.B. pencil only. Candidate should note that if they fail to enter the Booklet Serial Number and the Booklet Code on the OMR Response Sheet, their Answer Sheet will not be valued. Candidate shall shade one of the circles 1, 2, 3 or 4 corresponding question on the OMR Response Sheet using H.B. Pencil only. Candidate should note that their OMR Response Sheet will be invalidated if the circles against the question are shaded using Black / Blue ink pen / Ball pen / any other pencil other than H.B. Pencil or if more than one circle is shaded against any question.
- One mark will be awarded for every correct answer. There are no negative marks.
- The OMR Response Sheet will not be valued if the candidate:
 - (a) Writes the Hall Ticket Number in any part of the OMR Response Sheet except in the space provided for the purpose.
 - (b) Writes any irrelevant matter including religious symbols, words, prayers or any communication whatsoever in any part of the OMR Response Sheet.
 - (c) Adopts any other malpractice.
- Rough work should be done only in the space provided in the Question Paper Booklet.
- 10. No loose sheets or papers will be allowed in the examination hall.
- 11. Timings of Test: 10.00 A.M. to 1.00 P.M.
- 12. Candidate should ensure that he / she enters his / her name and appends signature on the Question paper booklet, leaflet attached to this question paper booklet and also on the OMR Response Sheet in the space provided. Candidate should ensure that the invigilator puts his signature on this question paper booklet, leaflet attached to the question paper booklet and also on the OMR Response Sheet.
- 13. Before leaving the examination hall candidate should return both the OMR Response Sheet and the leaflet attached to this question paper booklet to the invigilator. Failure to return any of the above shall be construed as malpractice in the examination. Question paper booklet may be retained by the candidate.
- 14. This booklet contains a total of 32 pages including Cover page and the pages for Rough Work.

MATHEMATICS

1. If
$$A = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$
, then $A^4 = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$

- (1) 31
- (2) 91
- (3) 271
- (4) 811
- 2. If $A = \begin{bmatrix} 0 & 2 & 1 \\ -2 & 0 & -2 \\ -1 & x & 0 \end{bmatrix}$ is a skew symmetric matrix, then the value of x is
 - (1) 1
- (2) 2
- (3) 3
- (4) 4
- What is the number of all possible matrices with each entry as 0 or 1 if the order of matrices is
 - (1) 64
- (2) 268 (3) 512
- (4) 256

4. If
$$A = \begin{bmatrix} 1 & i & -i \\ i & -i & 1 \\ -i & 1 & i \end{bmatrix}$$
, then $|A| =$

- (1) 1 (2) 2 (3) 3 (4) 4

5.	The solution of a system of linear equations 2.	X.	y + 3z = 9, x + y + z =	= 6, x -	· y + z = 2 i	is
----	---	----	-------------------------	----------	---------------	----

- (1) x = -1, y = -2, z = -3
- (2) x = 3, y = 2, z = 1
- (3) x = 2, y = 1, z = 3

(4) x = 1, y = 2, z = 3

6. If
$$\frac{1}{x^2 + a^2} = \frac{A}{x + ai} + \frac{B}{x - ai}$$
 then $A =$ ______, $B =$ ______.

- (1) $\frac{1}{2ai}$, $-\frac{1}{2ai}$ (2) $-\frac{1}{2ai}$, $\frac{1}{2ai}$ (3) $\frac{1}{ai}$, $-\frac{1}{ai}$ (4) $-\frac{1}{ai}$, $\frac{1}{ai}$

7. If
$$\frac{2x+4}{(x-1)^3} = \frac{A_1}{(x-1)} + \frac{A_2}{(x-1)^2} + \frac{A_3}{(x-1)^3}$$
 then $\sum_{i=1}^3 A_i$ is equal to

- (2) 2A,
- (3) 4A,
- (4) 4A,

8. The period of the function
$$f(x) = |\sin x|$$
 is

- (1) T
- $(2) 2\pi$
- $(3) 3\pi$
- $(4) 4\pi$

- (1) 1
- (2) 0
- (3) 2 (4) -1

- (1) $\frac{\sqrt{5}+1}{4}$ (2) $\frac{\sqrt{5}+1}{2}$ (3) $\frac{\sqrt{5}-1}{2}$ (4) $\frac{\sqrt{5}-1}{4}$

11. If
$$\Lambda + B + C = \pi$$
, then $\sin 2\Lambda + \sin 2B + \sin 2C =$

(1) 4 cosA sinB cosC

(2) 4 sinA cosB sinC

(3) 4 cosA cosB cosC

(4) 4 sinA sinB sinC

12. The principal solution of Tan
$$x = 0$$
 is

(1) $x = n\pi, n \in \mathbb{Z}$

(2) x=0

- (3) $x=(2n+1) \pi' 2. n \in \mathbb{Z}$
- (4) $x = n\pi + \alpha, n \in \mathbb{Z}$

13.	The value of Tan	(2)	Tan	(3)	i
	THE VALUE OF THE	1-1		1-1	

- (2) $\frac{\pi}{2}$
- (3) $\frac{\pi}{3}$

14. If the sides of a right angle triangle are in A.P., then the ratio of its sides is

- (1) 1:2:3
- (2) 2:3:4
- (3) 3:4:5
- (4) 4:5:6

15. The value of
$$r.r_1.r_2.r_3$$
 is

- (1) Δ^2 (2) Δ^{-2}
- (3) Δ⁻³
- (4) A

16.
$$\frac{1}{r1} + \frac{1}{r2} + \frac{1}{r3} =$$

- (1) $\frac{1}{r}$ (2) $\frac{1}{2r}$
- (3) $\frac{1}{R}$
- (4) $\frac{1}{\Delta}$

17. If
$$a=6$$
, $b=5$, $c=9$, then the value of angle A is

- (1) $\cos^{-1}(2/9)$ (2) $\cos^{-1}(2/5)$ (3) $\cos^{-1}(7/9)$ (4) $\cos^{-1}(1/3)$

- (1) $\sqrt{2}e^{i\pi/4}$ (2) $\sqrt{2}e^{i\pi/4}$ (3) $\sqrt{2}e^{i\pi/4}$ (4) $\sqrt{2}e^{i\pi/2}$

if i, to, to be the cube roots of unity, then the value of 2° .2° .2° is

- (1) ()
- (2) (2)
- (3) 1
- (4) 0

... The intercept made on X-axis by the circle $x^2+y^2+2gx+2fy+c=0$ is

- (1) $\sqrt{g^2-c}$ (2) $\sqrt{f^2-c}$ (3) $2\sqrt{g^2-c}$ (4) $2\sqrt{f^2-c}$

21. If one end of the diameter of the circle $x^2+y^2-5x-8y+13=0$ is (2, 7), then the other end of the diameter is

- (1) (3.1)
- (2) (1, 3) (3) (-3, -1) (4) (-1, -3)

- 22. The radius of the circle $\sqrt{1+m^2(x^2+y^2)}-2cx-2mcy=0$ is
 - (1) 2c
- (2) 4c
- (4) c
- 23. The parametric equations of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ are
 - (1) $x = a \sec \theta, y = b \tan \theta$
- (2) $x = b \sin\theta$, $y = a \cos\theta$
- (3) $x = a \cos\theta, y = b \sin\theta$
- (4) $x = a \csc\theta$, $y = b \cot\theta$
- 24. The equation of the directrix of the parabola $2x^2 = -7y$ is
 - (1) 8y+7=0
- (2) 8y-7=0
- (3) 7y+8=0
- (4) 8x-7=0
- 25. The condition for a straight line y = mx + c to be a tangent to the hyperbola $\frac{x^2}{a^2} \frac{y^2}{b^2} = 1$ is
 - (1) c = a/m
- (2) $c^2 = a^2m^2 b^2$ (3) $c^2 = a^2m^2 + b^2$ (4) $c^2 = a/m$

- 26. $\lim_{x \to 1} \frac{\sqrt{5x-4} \sqrt{x}}{x-1}$ is
 - (1) 3
- (2) 2
- (3) 4
- (4) 1

- 27. $\log i =$
 - (1) $\pi/2$
- (2) $\pi/4$
- (3) $i\pi/2$
- (4) $i\pi/4$

- 28. $\frac{d}{dr}[\log_7 X] =$

- (1) $\frac{1}{x}$ (2) $X \log_7^e$ (3) $\frac{1}{x} \log_7^e$ (4) $\frac{1}{x} \log_7^e$
- 29. $\frac{d}{dx}[2\cosh x] =$
 - (1) $\frac{e^x + e^{-x}}{2}$ (2) $\frac{e^x e^{-x}}{2}$ (3) $e^x + e^{-x}$ (4) $e^x e^{-x}$

$$30. \quad \frac{d}{dx} \left[\cos^{-1} \left(\frac{1 - x^2}{1 + x^2} \right) \right] =$$

- (1) $\frac{1}{1+x^2}$ (2) $\frac{-1}{1+x^2}$ (3) $\frac{2}{1+x^2}$ (4) $\frac{-2}{1+x^2}$

31. If
$$x = at^2$$
, $y = 2at$, then $\frac{dy}{dx} =$

- (1) $\sqrt{\frac{y}{x}}$ (2) $\sqrt{\frac{x}{a}}$ (3) $\sqrt{\frac{a}{x}}$ (4) $\sqrt{\frac{x}{y}}$

32. The derivative of
$$e^x$$
 with respect to \sqrt{x} is

- (1) $\frac{2\sqrt{x}}{e^x}$ (2) $2\sqrt{x}e^x$ (3) $\frac{e^x}{2\sqrt{x}}$ (4) $\sqrt{x}e^x$

33. The equation of the normal to the curve
$$y = 5x^4$$
 at the point (1, 5) is

- (1) x + 20y = 99 (2) x + 20y = 101 (3) x 20y = 99 (4) x 20y = 101

34. The angle between the curves
$$y^2 = 4x$$
 and $x^2 + y^2 = 5$ is

- (1) $\frac{\pi}{4}$ (2) $\tan^{-1}(2)$ (3) $\tan^{-1}(3)$ (4) $\tan^{-1}(4)$

35. If
$$u = x^3y^3$$
 then $\frac{\partial^3 u}{\partial x^3} + \frac{\partial^3 u}{\partial y^3} =$

- (1) $6(x^3+y^3)$ (2) $6x^3y^3$
- (3) $6x^3$
- $(4) 6y^3$

36.
$$\int \csc x dx =$$

- (1) $\log(\csc x + \cot x) + C$
- (2) $\log(\cot x/2) + C$

(3) $\log (\tan x/2) + C$

(4) $-\csc x.\cot x + C$

37.
$$\int_0^{\frac{\pi}{2}} \cos^{11} x \, dx =$$

- (1) $\frac{256}{693}$ (2) $\frac{256\pi}{693}$ (3) $\frac{\pi}{4}$

38.
$$\int f'(x) [f(x)]^n dx =$$

(1)
$$\frac{[f(x)]^{n-1}}{n-1} + C$$

(2)
$$\frac{[f(x)]^{n+1}}{n+1} + C$$

$$(3) \quad n[f(x)]^{n-1} + C$$

(1)
$$\frac{[f(x)]^{n-1}}{n-1} + C$$
 (2) $\frac{[f(x)]^{n+1}}{n+1} + C$ (3) $n[f(x)]^{n-1} + C$ (4) $(n+1)[f(x)]^{n+1} + C$

$$39. \quad \int \frac{dx}{(x+7)\sqrt{x+6}} =$$

(1)
$$Tan^{-1}(\sqrt{x+6})+C$$

(2)
$$2Tan^{-1}(\sqrt{x+6})+C$$

(3)
$$Tan^{-1}(x+7)+C$$

(4)
$$2Tan^{-1}(x+7)+C$$

40.
$$\int \tan^{-1} x \, dx =$$

(1)
$$x.Tan^{-1}x + \frac{1}{2}\log(1+x^2) + C$$

(2)
$$\frac{1}{1+x^2} + C$$

(3)
$$x^2 . Tan^{-1}x + C$$

(4)
$$x.Tan^{-1}x - \log \sqrt{1+x^2} + C$$

$$41. \quad \int \frac{dx}{1+e^{-x}} =$$

(1)
$$\log (1+e^{-x}) + C$$

(2)
$$\log (1+e^x) + C$$

(3)
$$e^{-x} + C$$

42.
$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin |x| \, dx =$$

- (1) 0
- (2) 1

- 43. Area under the curve $f(x) = \sin x$ in $[0, \pi]$ is
 - (1) 4 sq. units
- (2) 2 sq. units
- (3) 6 sq. units
- (4) 8 sq. units

- 44. The order of $x^3 \frac{d^3 y}{dx^3} + 2x^2 \frac{d^2 y}{dx^2} 3y = x$ is
 - (1) 1
- (2) 4
- (3) 3
- (4) 2

- 45. The degree of $\left[\frac{d^2 y}{dx^2} + \left(\frac{dy}{dx} \right)^2 \right]^{\frac{3}{2}} = a \frac{d^2 y}{dx^2}$ is
 - (1) 4
- (2) 2
- (3) 1
- (4) 3
- 46. The family of straight lines passing through the origin is represented by the differential equation

- (1) ydx + xdy = 0 (2) xdy ydx = 0 (3) xdx + ydy = 0 (4) xdx ydy = 0
- 47. The differential equitation $\frac{dy}{dx} + \frac{ax + hy + g}{hx + by + f} = 0$ is called
 - (1) Homogeneous (2) Exact
- (3) Linear
- (4) Legender
- 48. The solution of differential equation $\frac{dy}{dx} = e^{-x^2} 2xy$ is
 - (1) $y e^{-x^2} = x + c$ (2) $y e^x = x + c$ (3) $y e^{x^2} = x + c$ (4) y = x + c

- 49. The complementary function of $(D^3+D^2+D+1)y = 10$ is
 - (1) $C_1 \cos x + C_2 \sin x + C_3 e^{-x}$ (2) $C_1 \cos x + C_2 \sin x + C_3 e^{x}$ (3) $C_1 + C_2 \cos x + C_3 \sin x$ (4) $(C_1 + C_2 x + C_3 x^2) e^{x}$

- 50. Particular Integral of $(D-1)^4y = e^x$ is
 - (1) $x^4 e^x$

- (2) $\frac{x^4}{24}e^{-x}$ (3) $\frac{x^4}{12}e^x$ (4) $\frac{x^4}{24}e^x$

Set Code :	T2
Booklet Code :	

PHYSICS

51. Two quantities A and B are related by the relation A/B = m where m is linear mass density and A is

force. The dimensions of B will be

	 same as that of latent heat 		(2)	same as that of	f press	essure			
	(3) same as that of work				(4)	same as that of momentum			
52.	The	dimensional for	mula c	of capacitance in	terms	of M, L, T and	I is		
	(1)	$[ML^2T^2P]$	(2)	$[ML^{-2}T^4I^2]$	(3)	$[M^{\cdot l}L^3T^3l]$	(4)	[M-1L-2T4I2]	
53.	If <i>I</i> ,	m and n are the	firecti	on cosines of a v	ector,	then			
	(1)	l+m+n=1	(2)	$l^2 + m^2 + n^2 = 1$	(3)	$\frac{1}{l} + \frac{1}{m} + \frac{1}{n} = 1$	(4)	lmn = 1	
54.	The	angle between i	+j and	j+k is					
	(1)	0° ·	(2)	90°	(3)	45°	(4)	60°	
55.	A pa	article is moving s ⁻¹ northwards. T	eastw he ave	vards with a veloc erage acceleration	city o	f 5 ms ⁻¹ . In 10 so his time is	econds	the velocity changes to	
	(1)	$\frac{1}{\sqrt{2}}$ ms ⁻² towar	ds nor	th-west	(2)	zero			
	(3)	$\frac{1}{2}$ ms ⁻² toward	s nortl	1	(4)	$\frac{1}{\sqrt{2}}$ ms ⁻² towa	rds no	rth-east	
56.		linear momentu	m of a	particle varies w	rith ti	$\operatorname{me} t \operatorname{as} p = a + bt$	t+ct² w	hich of the following is	
	(1)	Force varies w	ith tim	e in a quadratic n	nanne	er.			
	(2)	Force is time-o							
	(3)	The velocity o	f the p	article is proport	ional	to time.		7.	
	(4) The displacement of the particle is proportional to t.								

(3) 3v/4 (4) 4v/3

57. A shell of mass m moving with a velocity v suddenly explodes into two pieces. One part of mass

m/4 remains stationary. The velocity of the other part is

(2) 2v

(1) v

Set Code :	T2
Booklet Code :	A

58.	The	velocity of a fi	reely fal	ling body afte	r 2s is				54
	(1)	9.8 ms ⁻¹	(2)	10.2 ms ⁻¹	(3)	18.6 ms ⁻¹	(4)	19.6 ms ⁻¹	
59.	A lar	rge number of ground on which	bullets a	re fired in all d bullets will sp	lirections read is	s with the same	speed to	. The maxim	um area on
	(1)	$\frac{\pi u^2}{g^2}$	(2)	$\frac{\pi u^4}{g^2}$	(3)	$\frac{\pi u^2}{g^4}$	(4)	$\frac{\pi u}{g^4}$	
60.	The	minimum stop	ping dis	tance for a car between the ty	of mass res and t	m, moving with the road is μ, ν	th a spec	d v along a le	vel road, if
	(1)	$\frac{v^2}{2\mu g}$	(2)	$\frac{v^2}{\mu g}$	(3)	$\frac{v^2}{4\mu g}$	(4)	$\frac{v}{2\mu g}$	
61.		en a bicycle is that it acts							
	(1)			ction on the fro					
	(2)	In the forwar	d directi	on on the from	t wheel a	nd in the back	ward dir	ection on the	rear wheel
	(3)	In the backw	ard dire	ction on both t	he front	and the rear wi	heels		
	(4)	In the forwar	d direct	ion on both the	e front ar	nd the rear who	eels		
62.	In a	perfectly inela	stic col	lision, the two	bodies				
	(1)	strike and ex	plode		(2)	explode with	out strik	ing	
	(3)	implode and	explode		(4)	combine and	move to	gether	
63.		ler the action o	f a cons	tant force, a pa	article is	experiencing	a consta	nt acceleratio	on, then the
	(1)	zero				positive			
	(3)	negative			(4)	increasing ur	niformly	with time	
			34						

			Set Code : T2 Booklet Code : A
Con	sider the following two statements:		
A:	Linear momentum of a system of par	rticles	is zero.
B:	Kinetic energy of a system of particl	es is z	ero.
Ther	1		
(1)	A implies B & B implies A	(2)	A does not imply B & B does not imply A
(3)	A implies B but B does not imply A	(4)	A does not imply B but B implies A

(4) 10s

65.	An engine develops 10 kW of power. How much time will it take to lift a mass of 200 kg to a
	height of 40 m? (Given $g = 10 \text{ ms}^{-2}$)

(3) 8s

- 66. If a spring has time period T, and is cut into n equal parts, then the time period will be
 - (1) $T\sqrt{n}$ (2) $\frac{T}{\sqrt{n}}$ (3) nT (4) T
- 67. When temperature increases, the frequency of a tuning fork

(2) 5s

64. Consider the following two statements:

(1) increases

(1) 4s

- (2) decreases
- (3) remains same
- (4) increases or decreases depending on the materials
- 68. If a simple harmonic motion is represented by $\frac{d^2x}{dy^2} + \alpha x = 0$, its time period is
 - (1) $2\pi\sqrt{\alpha}$ (2) $2\pi\alpha$ (3) $\frac{2\pi}{\sqrt{\alpha}}$ (4) $\frac{2\pi}{\alpha}$
- A cinema hall has volume of 7500 m³. It is required to have reverberation time of 1.5 seconds. The total absorption in the half should be
 - (i) 850 w-m2
- (2) 82.50 w-m²
- (3) 8.250 w-m²
- (4) 0.825 w-m²

70.	To a	bsorb the so	ound in a ha	ll which o	ing are used				
	(1)	Glasses, s	tores		(2)	Carpets, ci	urtains		
	(3)	Polished s	urfaces		(4)	Platforms		+11	
71.	IfN	represents a	wagadro's	number, th	en the numb	er of molec	ules in 6 gr	n of hydroge	n at NTP is
		2N		3N	(3)			N/6	
72.	The	mean transl	lational kin	etic energ	y of a perfec	t gas molec	ule at the te	emperature T	Kis
	(1)	$\frac{1}{2}kT$	(2)	kT	(3)	$\frac{3}{2}kT$	(4)	2kT	
			E				1 100		
73.	The	amount of h	neat given t	o a body w	hich raises				
	(1) water equivalent				(2)	thermal heat capacity			
	(3)	specific he	eat		(4)	temperatu	re gradient		
74.		ing an adiah				s is found to	o be propoi	tional to the	cube of its
	(1)	$\frac{3}{2}$	(2)	$\frac{4}{3}$	(3)	2	(4)	$\frac{5}{3}$	
75.	(1) (2)	to protect	the fiber fi the fiber f	rom mecha rom corros	nical stress				

(4) to protect the fiber from electromagnetic guidance

Set Code :	T2
Booklet Code :	A

CHEMISTRY

76.	The	valency electro	nic co	nfiguration of	Phospho	prous atom (At.)	No. 15) is	
	(1)	$3s^2 3p^3$	(2)	3s1 3p3 3d1	(3)	$3s^23p^23d^1$	(4)	$3s^1 3p^2 3d^2$	
77.	An	element 'A' of A	t.No.12	2 combines wi	th an ele	ment 'B' of At.N	o.17. 7	The compound for	ned is
	(1)	covalent AB	(2)	ionic AB ₂	(3)	covalent AB ₂	(4).	ionic AB	
78.	The	number of neut	rons p	resent in the a	tom of	Ba ¹³⁷ is			
	(1)	56	(2)	137	(3)	193	(4)	81	
79.	Hyd	rogen bonding	in wate	r molecule is	responsi	ble for			
	(1)	decrease in its	freezi	ng point	(2)	increase in its	degree	of ionization	
	(3)	increase in its	boiling	g point	(4)	decrease in its	boilin	g point	
80.	In th	e HCl molecule	, the b	onding between	en hydro	gen and chlorine	is		
	(1)	purely covalen	it (2)	purely ionic	(3)	polar covalent	(4)	complex coordin	nate
81.	Pota	ssium metal and	d potas	sium ions				*	
	(1)	both react with	water	-	(2)	have the same	numbe	er of protons	
	(3)	both react with	chlori	ine gas	(4).	have the same	electro	onic configuration	
82.	stane	dard flask. 10 ml er into 100 ml of	ofthis	solution were	pipetted ntration o	out into another f of the sodium chl	lask ar	made upto 100 m nd made up with dis solution now is	
	(1)	0.1 M	(2)	1.0 M	(3)	0.5 M	(4)	0.25 M	
83.	Con	centration of a 1	1.0 M s	olution of pho	osphoric	acid in water is			
	(1)	0.33 N	(2)	1.0 N	(3)	2.0 N	(4)	3.0 N	
84.	Whi	ch of the follow	ing is a	Lewis acid?					
	(1)	Ammonia			(2)	Berylium chlor	ide		
	(3)	Boron trifluori	ide		(4)	Magnesium oxi	ide		
					14-A				

Set Code :	T2
Booklet Code :	A

85.	Which of the following constitutes the components of a buffer solution?										
	(1)			nd potassium hyd	lroxid	le					
	(2)	(2) Sodium acetate and acetic acid									
	(3)	•	-	and sulphuric aci	d						
	(4)	Calcium chlor	ide and	l calcium acetate							
86.	Whi	ch of the follow	ving is	an electrolyte?							
	(1)	Acetic acid	(2)	Glucose	(3)	Urea	(4)	Pyridine			
87.	Calculate the Standard emf of the cell, $Cd/Cd^{+2}//Cu^{+2}/Cu$ given that E^0 $Cd/Cd^{+2} = 0.44V$ and E^0 $Cu/Cu^{+2} = (-) 0.34$ V.										
	(1)	(-) 1.0 V	(2)	1.0 V	(3)	(-) 0.78 V	(4)	0.78 V			
88.	Asc	lution of nicke	l ċhlori	de was electrolys	sed us	sing Platinum e	lectrod	es. After electrolysis,			
	(1)							ed at the cathode			
	(3)	H ₂ gas will be	liberat	ed at the anode	(4)	nickel will be	deposi	ted on the cathode			
89.	Which of the following metals will undergo oxidation fastest?										
	(1)	Cu	(2)	Li	(3)	Zinc	(4)	Iron			
90.	Which of the following cannot be used for the sterilization of drinking water?										
	(1)	Ozone			(2)						
	(3)	Potassium Ch	loride		(4)	Chlorine water					
91.		ater sample sho			g/litro	e of magnesiun	n sulpha	te. Then, its hardness in			
		1.0 ppm		1.20 ppm	(3)	0.60 ppm	(4)	2.40 ppm			
92.	Sod	a used in the L-	S proce	ess for softening	of wa	ter is, Chemica	illy.				
	(1)	sodium bicarb	onate		(2)	sodium carbo	nate dec	cahydrate			
	(3)	sodium carbo	nate		(4)	sodium hydro	oxide (4	0%)			
93.	The	process of cem	entatio	n with zinc powd	er is l	known as					
		sherardizing		zincing			g (4)	electroplating			

94.	Carr	osion of a metal is fast	est in		
	(1)	rain-water (2)	acidulated water	(3)	distilled water (4) de-ionised water
95.	Whi	ch of the following is a	thermoset polyr	ner?	*
	(1)	Polystyrene		(2)	PVC
	(3)	Polythene		(4)	Urea-formaldehyde resin
96.	Che	mically, neoprene is			
	(1)	polyvinyl benzene		(2)	polyacetylene
	(3)	polychloroprene		(4)	poly-1,3-butadiene
97.	Vuk	anization involves hear	ting of raw rubber	with	e e
	(1)	selenium element		(2)	elemental sulphur
	(3)	a mixture of Se and el	emental sulphur	(4)	a mixture of selenium and sulphur dioxide
					·°
98.		ol largely contains			
	(1)	a mixture of unsatura			C ₈
	(2)	a mixture of benzene,			n .
	(3)	a mixture of saturated			
	(4)	a mixture of saturated	l hydrocarbons C	- C ₈	
99.	Whi	ch of the following gas	ses is largely resp	onsil	ble for acid-rain?
	(1)	SO ₂ & NO ₂		(2)	CO ₂ & water vapour
	(3)	CO ₂ & N ₂		(4)	N ₂ & CO ₂
100.	BOI	O stands for			
	(1)	Biogenetic Oxygen D	emand	(2)	Biometric Oxygen Demand
	(2)	Piological Ovygen D		(4)	Biospecific Oxygen Demand

Set Code :	T2
Booklet Code :	Α

CHEMICAL ENGINEERING

101. T	The	property of met	als or	alloys which	describe	s its ability to be	e draw	n into wires is	known as
(1)	tenacity			(2)				
(:	3)	porosity			(4)	malleability			
102. S	tair	nless steel, in ad	dition	to iron and c	arbon co	ntains			
		aluminium		zinc		lead	(4)	chromium	
103. W	Vhi	ch of the followi	ng ha	s maximum n	nelting p	oint?		2.	e:
		wrought iron			(2)		n		
(3	3)	steel .		*	(4)				
104. TI	he j	process of prote	ction	of iron by coa	ating with	h zinc is called			
(1		tempering			. (2)	galvanizing			
(3	3)	nitriding			(4)	smelting			
105. Tł	he r	nost commonly	used	resin for mak	ing reinf	orced plastic is			
(1		unsaturated poly			(2)	polypropylene			
(3)	polyurethane			(4)	nylon-6			
06. Pr	ese	nce of mangane	se in	allov steel im	proves it	is.			
		corrosion resist			(2)	cutting ability			
(3)		abrasion resistar		7	(4)		reep r	resistance	
07. Eq	jual essi	weights of meth	ane a	nd oxygen are gen is	mixed i	n a empty reactor	r at 25	°C. The fraction	on of total
(1))	1/3	(2)	1/2	(3)	3/2	(4)	$\frac{1}{2} \times \frac{273}{298}$	
					17-À			-	(CHE)

123	. The	major constituent in black liquor is		
	(1)	sodium carbonate	(2)	sodium sulphate
	(3)	silica	(4)	iron oxide
124	. Cor	ntact process for the manufacture of s	ulphur	ic acid yields
	(1)	80% H ₂ SO ₄ only	(2)	98% H ₂ SO ₄ and higher
	(3)	95% H ₂ SO ₄ only	(4)	90% H ₂ SO ₄ only
125	. The	constituents of water gas are		
	(1)	CO and H ₂ O	(2)	CO ₂ and N ₂
	(3)	CO and H ₂	(4)	CO and N ₂
126.	The	principal raw materials for the manuf	acture	of soda ash by Solvay process are
	(1)	limestone and potassium chloride	(2)	dolomite and sodium hydroxide
	(3)	limestone, brine and coal	(4)	coal and caustic soda •
127.	Clin	ker is the mass obtained by heating		
	(1)	powdered limestone and clay	(2)	gypsum
	(3)	dolomite	(4)	sand, limestone and washing soda
128.	Who	en temporary hard water is boiled, one	of the	substances formed is
	(1)	calcium bicarbonate	(2)	calcium sulphate
	(3)	hydrogen chloride	(4)	carbon dioxide
129.	Nitri	ic acid is manufactured by catalytic or	cidation	of ammonia. This process is called
	(1)	Solvay process	(2)	Haber's process
	(3)	Ostwald's process	(4)	Bosch process

130.	Var	nish does not co	ntain								
	(1)	pigment				(2)	thinner				
	(3)	dryer				(4)	anti skinning	agent			
131.	Cata	alyst used in oxi	dation	of amn	nonia is						
	(1)	Platinum-Bery	llium			(2)	Platinum-Rho	dium			
	(3)	Cobalt-Molyb	denum			(4)	Platinum-Mo	lybdeni	ım		
132.	Flui	ds which show	an appa	arent in	crease ir	viscos	ity with time ar	e calle	d		
	(1)	rheopectic	(2)	thixot	ropic	(3)	ideal fluids	(4)	newtonian fluids		
133.	Ben	noulli's theorem	deals	with the	e law of	conser	vation of				
	(1)	energy				(2)	mass				
	(3)	momentum				(4)	gravity				
134.	Pito	t tube is used to	meası	ıre							
	(1)	local velocity	at a po	int		(2)	volumetric flow rate				
	(3)	average veloci	ty			(4)	pressure at a point				
135.	Stok	e (St) is the uni	t of kir	nematic	viscosit	y and o	one stoke is equ	al to			
	(1)	$1 m^2/s$	(2)	1ft ² /s		(3)	1 cm ² /s	(4)	1 mm ² /s		
136.	For	laminar flow in	a pipe,	the val	ue of ki	netic er	nergy correction	n factor	r(α) is		
	(1)	1	(2)	1.01		(3)	1.33	(4)	2		
137.	Whi	ch of the follow	ing equ	ations	is applic	able fo	r the flow of flu	uid thro	ugh packed bed?		
		Hagen-Poiseui				(2)					
		Nikuradse equa				(4)	Ergun equation		14		

Set Code :	T2
Booklet Code :	A

138.		ch of the following pumps is preferre		
	(1)	Gear pump	(2)	Lobe pump
	(3)	Screw pump	(4)	Centrifugal pump
139.	Min	imum porosity for fluidization is		
	(1)	that corresponding to static bed		
	(2)	that corresponding to completely fl	uidized	bed
	(3)	the porosity of the bed when true fle		
	(4)	less than that of the static bed		
140	For	turbulent flow in smooth pipe of diar	neter D.	the transition length is taken as
140.		0.05 D (2) 50 D		150 D (4) 0.5 D
141.		e of that factor were increased? temperature difference	(2)	transfer rate by conduction to decrease, if the thermal conductivity
	(3)	area	(4)	thickness
142.	The	rmal conductivity is minimum for		
	(1)	asphalt	(2)	water
	(3)	petroleum coke	(4)	air
143.	. In fo	orced convection, fluid moves under	the influ	nence of
	(1)	changes in fluid pressure produced	by exter	nal work
	(2)	buoyant forces arising from change	es in den	sity
	(3)	elastic forces		*1
	(4)	surface tension forces		
144	. The	Graetz number is associated with		
	(1)	heat transfer by radiation	(2)	heat transfer in laminar flow
	(3)	heat transfer in turbulent flow	(4)	mass transfer operations
			22-A	(CHE)

145	5. Dr	op-wise condens	sation	usually occurs	on				
	(1)	smooth surface	ce		(2)	oily surface	:		
	(3)	coated surfac	e		(4)	glazed surfa	ice		٠
146	5. The	e presence of sm	all am	ounts of non-c	ondensir	ng gas in a con	densing	vapor	
	(1)			rate of conde					
	(2)			e rate of conde					
2.5	(3)	does not affect the rate of condensation							
	(4)	increases the	conde	nsing film coe	fficient				
147	. The	heat flux in the	free c	onvection regi	me of no	ool boiling var	ies as th	ie.	
		ΔT^3		ΔT ^{5/4}		ΔT ²		ΔT1/4	
148	, In a	single effect ev	aporat	ion, to evapora	ate 1 lb o	of water from a	a solutio	on calls for	
		1 to 1.3lb of st				1.5 to 2 lb st			
	(3)	2 to 2.5 lb of s		(4)	0.5 to 0.8 lb				
149.	. The	total emissivity	ofare	eal surface is					
	(1)	less than zero			(2)	greater than	one		
	(3)	equal to one			(4)	greater than			
150.	The	units of fouling	factor	are					
	(1)	m ² .K/W	(2)	W/(m ² .K)	(3)	m.K/W	(4)	$m^2 K^4/W$	
151.	Crus	shing efficiency	of a si	ze reduction ec	quipmen	t ranges betwe	en		
	(1)	0.1 to 2%	(2)	10 to 20%	(3)	40 to 50%	(4)	70 to 80%	٠.
152.	Whi	ch of the followi	ng scr	eens has the m	aximum	capacity?			
	(1)	Grizzlies	-			Trommels			
	(3)	Vibrating screen	n		(4)	Stationery sc	reen		
					23-A				(CHE)

Set Code :	T2
Booklet Code :	A

153.	In a	ball mill most of	of the re	ductin is do	one by			1		
	(1)	slow compres	sion		(2)	cutting				
	(3)	attrition			(4)	impact				
154.	In a	rotary-drum fil	ter, the	fractional s	ubmergeno	e of the drum	in the sl	urry is abou	ıt	
	(1)	0.03	(2)	0.30	(3)	0.90	(4)	0.15		
155.	Indu	strially, the pro	cess of	sedimentat	ion is cond	lucted on a lar	ge scale	in equipme	nt called	
	(1)	sorting classi	fiers		(2)	cyclones				
	(3)	thickeners			(4)	filters				
156.	The	speed, in rpm,	of a con	tinuous rot	ary vacuum	n filter may be				
	(1)	1	(2)	100	(3)	1000	(4)	10000	50	
157.	Frot	th flotation is m	nost suit	able for tre	ating					
	(1)	iron ores			(2)	sulphide ore	s			
	(3)	quartzite			. (4)	nitride ores			ia .	
158.	The	most efficient	equipm	ent for remo	oval of sub-	micron dust p	articles	form blast f	urnace gas is	
	(1)	venturi scrubl	ber		(2)	gravity settling chamber				
	(3)	electrostatic	precipit	ator	(4)	cyclone sepa	arator		¥8	
159.	Cha	inge of state, e.g	g. freezi	ng, melting	, evaporati	on and conden	sation, i	s an		
	(1)	adiabatic pro	cess		(2)	isobaric pro	cess			
	(3)	isothermal pr	rocess		(4)	isochoric pr	rocess			
160.	Mat	hematical state	ment o	second lav	v of thermo	odynamics is				
	(1)	$\Delta S = 0$	(2)	$\Delta S > 0$	(3)	$\Delta S \le 0$	(4)	$\Delta S \ge 0$		
					24-A				(CHE)	

								5	Set Code : T2		
								Book	et Code : A		
161.	-	variation of heat	of re	action with temp	eratur	e at constant pro	essure	or at co	onstant volume is		
	(1)	Kirchoff's law	(2)	Fourier's law	(3)	Laplace law	(4)	Hess	's law		
162.	The	principal of refri	gerat	ion is based on							
	(1)	Zeroth law of th	ermo	dynamics	(2)	first law of the	rmody	namic	s		
	(3)	second law of the	nermo	odynamics	(4)	thrid law of the	ermody	namio	rs ·		
163.	_	as is termed an id w deviation from			gas co	uation PV = RT	. When	do yo	ou expect a gas to		
	(1)	At high pressure	es and	low temperatur	es						
	(2)	At low pressure	s and	low temperature	es						
	(3)	At high pressures and high temperatures									
	(4)	At low pressure	s and	high temperatur	es						
164.	From		plot	ln k versus 1/T g	ives a:	straight line with	a slop	e of (-	E/R). The units of		
	(1)	K	(2)	cal	(3)	cal/K		(4)	K/cal		
165.	The	rate constant of a	ny re	action depends of	on						
	(1)	the temperature	of th	e system	(2)	the time of rea	ction				
	(3)	the extent of re-	action	1	(4)	the initial cond	centrat	ion of	the reactants		
166.	A ca	talyst is a substar	ice w	hich							
	(1)	increases the eq	uilib	rium concentrati	on of	the product					
	(2)	changes the equ	ilibri	um constant of t	he rea	ction					
	(3)										
	(4)	supplies energy									

25-A (CHE)

Set Code :	T2
Booklet Code:	A

	of th	ne initial concen	tration	, the ord	er of the							
	(1)	zero	(2)	one		(3)	two		(4)	three		
168.	The	irreversible rea	ction is	simply	the spec	ial ca	se of the re	eversible	e rea	action if		
	(1)	the concentrat	ion of	the react	ant at eq	uilibr	ium condi	tions is	zero) ii		
	(2)	the fractional	conver	sion of th	he reacta	int at	equilibriur	n condit	tions	s is zero		
	(3)	the equilibrium	n const	tant is ze	ro							
	(4)	the equilibrium	n const	tant is on	ie							
169.		steady state tem				nall ar	nount of lie	quid eva	pora	ating into a	large amou	ınt
	(1)	dry-bulb tempe	erature									
	(2)	dew point										
	(3)	wet-bulb temp	erature	:						•		
	(4)	bubble point										
170.	Rela	ative volatility, o	ι, for a	binary sy	ystem							
	(1)	decreases with	increa	ase in pre	essure	97						
	(2)	increases with	increa	se in pre	ssure							
	(3)	increases with	increa	se in tem	peratur	e at co	onstant pre	ssure				
	(4)	has no signific	ance ii	n distillat	tion ope	ration						
171.	At n	ninium reflux ra	tio the	operatin	ig cost o	f a dis	stillation c	olumn is	S	¥1		
	(1)	maximum	(2)	optimu	m	(3)	minimum	n	(4)	infinite		
172.	For	all useful liquid	-liquid	extraction	on opera	itions	the selecti	ivity of s	olve	ent must be	e	
	(1)	more than zero				(2)	more tha	n one				
	(3)	less than one				(4)	less than	or equa	l to	one	+	
											"	HE)

Set Code :	T2
Booklet Code :	A

173.	At f	ixed temperature, the solubility of	gases in so	olvent
	(1)	remains constant with change in	pressure	
	(2)	decreases with increase in pressu	ıre	
	(3)	increases with increase in pressu	ire	
	(4)	decreases exponentially with inc	rease in pr	essure
174.	Veg	etable oils are recovered from oil s	seeds by le	aching with
	(1)	hot sulphuric acid	(2)	cold water
	(3)	nitric acid	(4)	hexane
175.	The	ratio of momentum diffusivity to r	nass diffus	sivity is known as
	(1).	Schmidt number	(2)	Sherwood number
	(3)	Lewis number	(4)	Stanton number
176.	Whi	ch one of the cooling tower is mo	st efficient	?
	(1)	Chimney type natural draft coolin	ng tower	
	(2)	Atmospheric circulation type coo	oling tower	r
	(3)	Induced draft cooling tower.		
	(4)	Forced draft cooling tower		
177.	Gran	nular or crystalline material can be	dried in	
	(1)	37.00 to 10.00 to 10	(2)	rotary dryer
	(3)		(4)	screw-conveyor dryer
178.	Swe	nson-Walker crystallizer is a		
	(1)	continuous unit	(2)	batch unit
	(3)	semi-batch unit	(4)	cooling (adiabatic)-cum-evaporation device
179.	Whi	ch one of the following is a static	characteris	stic of instruments?
	(1)	Fidelity (2) Time lag	(3)	Dynamic error (4) Reproducibility

Set Code :	T2
Booklet Code :	A

180.	Wh	ich of the following is most suitable	to meas	ure a temperature of 2000°C?
	(1)	Ordinary mercury-in-glass thermo	meter	
	(2)	Platinum resistance thermometer		
	(3)	Radiation pyrometer		
	(4)	Constant-volume hydrogen thermo	ometer	
181.	Offs	set is zero for		
	(1)	P-controller only	(2)	P-D controller only
	(3)	P- and P-D controllers	(4)	P-I and P-I-D controllers only
182.	On-	off control is a special case of		
	(1)	proportional control	(2)	proportional-integral control
	(3)	proportional-derivative control	(4)	proporțional-integral-derivative control
183.	Abs	olute pressure is measured by		
	(1)	a bourdon gauge	(2)	an ancroid barometer
	(3)	a differential manometer	. (4)	a vacuum gauge
184.	Resp	ponse of a linear control system for	a change	in set point is called
	(1)	frequency response	(2)	transient response
	(3)	servo problem	(4)	regulator problem
185.	Deg is ca		the chan	ges in measured variable withou dynamic error
	(1)	speed of response	(2)	reproducibility of instrument
	(3)	fidelity	(4)	its static characteristics
186.	Step	response of a first-order system is		
	(1)	under damped	(2)	critically damped
	(3)	over damped	(4)	undamped
			28-A	(CHE)

Set Code :	T2
Booklet Code :	A

						7974				
187.		ar radiation flux							ial to	
	(1)	1 cal/cm ²	(2)	1 Btu/ft ²	(3)	1 j/m ²	(4)	1 Btu/in ²		
188.	Cor	version efficie	ncies fo	or silicon cel	ls (i.e., so	olar cells) rang	ge betwe	en		
	(1)	10 and 15%			(2)	30 and 35%	,			
	(3)	90 and 95%			(4)	95% and 99	%			
189.	The	function of a w	vindmil	l is to extract	energy fr	om the wind a	and to pro	oduce		
	(1)	mechanical er			(2)					
	(3)	electrical ene	rgy		(4)	chemical en	377			
190.		maximum pov		lable in the	wind is di	rectly proport	tional to	the velocity o	f the wind	
	(1)	1	(2)	2	(3)	3	(4)	4		
191.	Liqu	uefied petroleur	m gas (l	LPG) is a						
	(1)	primary liquio	fuel		(2)	primary gase	eous fuel			
	(3)	secondary gas	seous fu	iel	(4)	secondary li	quid fuel			
192.	Whi	ch of the follow	ving iso	tope is a raw	material	for the produc	ction of F	u ²³⁹ nuclear	fuel?	
		U^{238}	(2)	U^{235}	(3)	000	(4)			
193.	The	quality of a goo	od fuel i	is						
	(1)	high calorific			(2)	low cost				
	(3)	easily availabl	e		(4)	no ash		9		
194.	Whi	ch one of the fo	llowin	g is the most	severe ai	r pollutant?				
		SO ₂		NO _X		СО	(4)	CH_4		
	. ,	-2	(=)	X	(-)	363/070	,			
					29-A			5	(CHE)	

Set Code : T2

Booklet Code : A

195.	The	rmal power plants are the major so	urce of	
	(1)	SO ₂ pollutants	(2)	ammonia pollutants
	(3)	NO _X pollutants	(4)	phosgene pollutants
196.	The	fire of electrical equipments can b	e extinguis	shed with the use of
	(1)	soda-acid extinguisher	(2)	carbon dioxide extinguisher
	(3)	foam extinguisher	(4)	antifreeze extinguisher
197.	Fire	is a proper combination of		
	(1)		(2)	fuel and oxygen
	(3)		(4)	oxidizing material and air
		Landles days a don shad oon bo ye	ed to store	
198.		I ventilated wooden shed can be us		
	(1)	oxidizing materials	(2)	flammable liquids
	(3)	acids	(4)	compressed gases
199.	The	biochemical treatment of sewage	effluents i	s essentially a process of
	(1)		(2)	
	(3)	dehydration	(4)	alkalinization
200.		ich one of the following chemica ution?		t in the form of inorganic impurity in water
	(1)	Proteins (2) Fats	(3)	Salts of metals (4) Carbohydrates