

CBSE Question Paper – 2010 MATHEMATICS (Theory) Class – XII

Time allowed: 3 hours

cbse

iess

Maximum Marks: 100

Instructions: Attempt all questions. All questions carry equal marks.

- 1. If f: $R \rightarrow R$ defined by $f(x) = (3 x^3)^{1/3}$, then find fof(x).
- 2. Write the principal value of sec⁻¹(-2).
- 3. What positive value of X makes the following pair of determinants equal?

 $\begin{vmatrix} 2x & 3 \\ 5 & x \end{vmatrix}, \begin{vmatrix} 16 & 3 \\ 5 & 2 \end{vmatrix}$

4. Evaluate :

$$\int \sec^2 (7 - 4x) dx$$

Other Educational Websites:

ICSEGuess.com | NIOSGuess.com | IGNOUGuess.com | IITGuess.com | MagicSense.com | AIPMTGuess.com | AIEEEGuess.com | IndiaGuess.com






5. Write the adjoin of the following matrix :

$$\begin{pmatrix} 2 & -1 \\ 4 & 3 \end{pmatrix}$$

6. write the value of the following integral



- 7. A is a square matrix of order 3 and |A| = 7. Write the value of $|adj \cdot A|$.
- 8. Write the distance of the following plane from the origin :

2x - y + 2z+1=0


9. Write a vector of magnitude 9 unites in the direction of vector

$$-2\hat{i} + \hat{j} + 2\hat{k}.$$
10. Find \times if $(2\hat{i} + 6\hat{j} + 14\hat{k}) \times (\hat{i} - \lambda\hat{j} + 7\hat{k}) = \vec{0}$
SECTION B

- 11. A family has 2 children. Find the probability that both are boys, if it is known that
 - (i) At least one of the children is a boy,

Other Educational Websites:

ICSEGuess.com | NIOSGuess.com | IGNOUGuess.com | IITGuess.com | MagicSense.com | AIPMTGuess.com | AIEEEGuess.com | IndiaGuess.com



iess

- 12. Show that the relation S in the set A = { $xEZ : 0 \le x \le 12$ } given by S = {(a, b): a, b E z. |a b| is divisible by 4} is an equivalence relation. Find the set of all elements related to 1.
- 13. Prove the following :

cbse

$$\tan^{-1} x + \tan^{-1} \left(\frac{2x}{1 - x^2} \right) = \tan^{-1} \left(\frac{3x - x^3}{1 - 3x^2} \right)$$

Prove the following:

Cos [tan⁻¹{sin(cot⁻¹ x)] =
$$\sqrt{\frac{1}{2}}$$

14. Express the following matrix as the sum of a symmetric and skew symmetric matrix, and verify your result :

OR

$$\begin{pmatrix} 3 & -2 & -4 \\ 3 & -2 & -5 \\ -1 & 1 & 2 \end{pmatrix}$$

Other Educational Websites:

ICSEGuess.com | NIOSGuess.com | IGNOUGuess.com | IITGuess.com | MagicSense.com | AIPMTGuess.com | AIEEEGuess.com | IndiaGuess.com

15.

cbse

If $\vec{a} = i + j + k$, b = 4i - 2j + 3k and c = i - 2j + k, find a vector of magnitude 6 units which is parallel to the vector $2\vec{a} - \vec{b} + 3\vec{c}$.

- OR
- Let $\overrightarrow{a} = \hat{i} + 4\hat{j} + 2\hat{k}$, $\overrightarrow{b} = 3\hat{i} 2\hat{j} + 7\hat{k}$ and $\overrightarrow{c} = 2\hat{i} - \hat{j} + 4\hat{k}$. Find a vector \overrightarrow{d} which is perpendicular to both \overrightarrow{a} and \overrightarrow{b} and $\overrightarrow{c} \cdot \overrightarrow{d} = 18$.
- 16. Find the points on the line $\frac{x+2}{3} = \frac{y+1}{2} = \frac{z-3}{2}$ at a distance of 5 units from the point P (1, 3, 3).

Find the distance of the point p(6,5,9) From the plane determined by the points A(3,-1,2), B(5,2,4) and C (-1,-1,6).

17. Solve the following differential equation :

$$(x^{2}-1)\frac{dy}{dx} + 2xy = \frac{1}{x^{2}-1}; |x| \neq 1$$

$$\sqrt{1 + x^2 + y^2 + x^2y^2} + xy\frac{dy}{dx} = 0$$

OR

Other Educational Websites:

ICSEGuess.com | NIOSGuess.com | IGNOUGuess.com | IITGuess.com | MagicSense.com | AIPMTGuess.com | AIEEEGuess.com | IndiaGuess.com



- 18. Show that the differential equation (x-y) $\frac{dy}{dx} = X + 2y$, is homogeneous and solve it.
- 19. Evaluate the following ;

$$\int \frac{x+2}{\sqrt{(x-2)(x-3)}} dx$$

20. Evaluate the following :

$$\int_{1}^{2} \frac{5x^2}{x^2 + 4x + 3} \, dx$$

21. If $y = e^{a \sin -1x}$, $-1 \le x \le 1$, then show that

(1-x²)
$$\frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2y = 0.$$

22. If
$$y = \cos^{-1}\left(\frac{3x+4\sqrt{1-x^2}}{5}\right)$$
, find $\frac{dy}{dx}$

SECTION C

23. Using properties of determinants, prove the following :

$$\begin{vmatrix} x & x^2 & 1 + px^3 \\ y & y^2 & 1 + py^3 \\ z & z^2 & 1 + pz^3 \end{vmatrix} = (1 + pxyz) (x-y) (y-z) (z-x)$$

Other Educational Websites:

ICSEGuess.com | NIOSGuess.com | IGNOUGuess.com | IITGuess.com | MagicSense.com | AIPMTGuess.com | AIEEEGuess.com | IndiaGuess.com



OR

Find the inverse of the following matrix using elementary operations:

$$\mathsf{A} = \begin{bmatrix} 1 & 2 & -2 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$$

- 24. A bag contains 4 balls two balls are drawn at random, and are found to be white. What is the probability that all balls are white?
- 25. One kind of cake requires 300 g of flour and 15 g of fat, another kind of cake requires 150 g of flour and 30 g of fat. Find the maximum number of cakes which can be made from 7.5 kg of flour and 600g of fat, assuming that there is no shortage of the other ingredients' used in making the cakes. Make it as an L.P.P and solve it graphically.
- 26. Find the coordinates of the foot of the perpendicular and the perpendicular distance of the point p (3, 2, 1) from the plane 2x-y+z+1=0. Find also, the image of the point in the plane.

27. Find the area of the circle $4x^2 + 4y^2 = 9$ which is interior to the parabola $x^2 = 4y$.

OR

Using integration, find the area of the triangle ABC, coordinates of whose vertices are A (4, 1), B (6, 6) and C (8, 4).

Other Educational Websites:

ICSEGuess.com | NIOSGuess.com | IGNOUGuess.com | IITGuess.com | MagicSense.com | AIPMTGuess.com | AIEEEGuess.com | IndiaGuess.com



28. If the length of three sides of a trapezium other than the base is 10 cm each , find the area of the trapezium, when it is maximum.

R

- 29. Find the intervals in which the following function is
 - (a) Strictly increasing,
 - (b) Strictly decreasing.

Other Educational Websites: