



7. If  $z = \frac{1-i\sqrt{3}}{1+i\sqrt{3}}$ , then  $\arg(z)$  is  
 (a)  $60^\circ$       (b)  $120^\circ$   
 (c)  $240^\circ$       (d)  $300^\circ$
8. If  $f(x) = \sqrt{\log_{10} x^2}$ . The set of all values of  $x$  for which  $f(x)$  is real, is  
 (a)  $[-1, 1]$   
 (b)  $[1, \infty)$   
 (c)  $(-\infty, -1]$   
 (d)  $(-\infty, -1] \cup [1, \infty)$
9. For what values of  $m$  can the expression  

$$2x^2 + mxy + 3y^2 - 5y - 2$$
 be expressed as the product of two linear factors?  
 (a) 0      (b)  $\pm 1$   
 (c)  $\pm 7$       (d) 49
10. If  $B$  is a non-singular matrix and  $A$  is a square matrix, then  $\det(B^{-1}AB)$  is equal to  
 (a)  $\det(A^{-1})$       (b)  $\det(B^{-1})$   
 (c)  $\det(A)$       (d)  $\det(B)$
11. If  $f(x)$ ,  $g(x)$  and  $h(x)$  are three polynomials of degree 2 and  

$$\Delta(x) = \begin{vmatrix} f(x) & g(x) & h(x) \\ f'(x) & g'(x) & h'(x) \\ f''(x) & g''(x) & h''(x) \end{vmatrix},$$
 then  $\Delta(x)$  is a polynomial of degree  
 (a) 2      (b) 3  
 (c) 0      (d) atmost 3
12. The chances of defective screws in three boxes  $A$ ,  $B$ ,  $C$  are  $\frac{1}{5}, \frac{1}{6}, \frac{1}{7}$  respectively. A box is selected at random and a screw drawn from it at random is found to be defective. Then, the probability that it came from box  $A$ , is  
 (a)  $\frac{16}{29}$       (b)  $\frac{1}{15}$   
 (c)  $\frac{27}{59}$       (d)  $\frac{42}{107}$
13. The value of  $\frac{\cos \theta}{1 + \sin \theta}$  is equal to  
 (a)  $\tan\left(\frac{\theta}{2} - \frac{\pi}{4}\right)$       (b)  $\tan\left(-\frac{\pi}{4} - \frac{\theta}{2}\right)$   
 (c)  $\tan\left(\frac{\pi}{4} - \frac{\theta}{2}\right)$       (d)  $\tan\left(\frac{\pi}{4} + \frac{\theta}{2}\right)$
14. If  $3 \sin \theta + 5 \cos \theta = 5$ , then the value of  $5 \sin \theta - 3 \cos \theta$  is equal to  
 (a) 5      (b) 3  
 (c) 4      (d) None of these
15. The principal value of  $\sin^{-1} \left\{ \sin \frac{5\pi}{6} \right\}$  is  
 (a)  $\frac{\pi}{6}$       (b)  $\frac{5\pi}{6}$   
 (c)  $\frac{7\pi}{6}$       (d) None of these
16. A rod of length  $l$  slides with its ends on two perpendicular lines. Then, the locus of its mid point is  
 (a)  $x^2 + y^2 = \frac{l^2}{4}$       (b)  $x^2 + y^2 = \frac{l^2}{2}$   
 (c)  $x^2 - y^2 = \frac{l^2}{4}$       (d) None of these
17. The equation of straight line through the intersection of line  $2x + y = 1$  and  $3x + 2y = 5$  and passing through the origin is  
 (a)  $7x + 3y = 0$       (b)  $7x - y = 0$   
 (c)  $3x + 2y = 0$       (d)  $x + y = 0$
18. The line joining  $(5, 0)$  to  $(10 \cos \theta, 10 \sin \theta)$  is divided internally in the ratio  $2 : 3$  at  $P$ . If  $\theta$  varies, then the locus of  $P$  is  
 (a) a straight line  
 (b) a pair of straight lines  
 (c) a circle  
 (d) None of the above
19. If  $2x + y + k = 0$  is a normal to the parabola  $y^2 = -8x$ , then the value of  $k$ , is  
 (a) 8      (b) 16  
 (c) 24      (d) 32
20.  $\lim_{n \rightarrow \infty} \left[ \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \dots + \frac{1}{n(n+1)} \right]$  is equal to  
 (a) 1      (b) -1  
 (c) 0      (d) None of these
21. The condition that the line  $lx + my = 1$  may be normal to the curve  $y^2 = 4ax$ , is  
 (a)  $al^3 - 2alm^2 = m^2$   
 (b)  $al^2 + 2alm^3 = m^2$   
 (c)  $al^3 + 2alm^2 = m^3$   
 (d)  $al^3 + 2alm^2 = m^2$

22. If  $\int f(x) dx = f(x)$ , then  $\int \{f(x)\}^2 dx$  is

- equal to  
 (a)  $\frac{1}{2} \{f(x)\}^2$       (b)  $\{f(x)\}^3$   
 (c)  $\frac{\{f(x)\}^3}{3}$       (d)  $\{f(x)\}^2$

23.  $\int \sin^{-1} \left\{ \frac{(2x+2)}{\sqrt{4x^2 + 8x + 13}} \right\} dx$  is equal to

- (a)  $(x+1) \tan^{-1} \left( \frac{2x+2}{3} \right)$   
 $- \frac{3}{4} \log \left( \frac{4x^2 + 8x + 13}{9} \right) + c$   
 (b)  $\frac{3}{2} \tan^{-1} \left( \frac{2x+2}{3} \right)$   
 $- \frac{3}{4} \log \left( \frac{4x^2 + 8x + 13}{9} \right) + c$   
 (c)  $(x+1) \tan^{-1} \left( \frac{2x+2}{3} \right)$   
 $- \frac{3}{2} \log (4x^2 + 8x + 13) + c$   
 (d)  $\frac{3}{2} (x+1) \tan^{-1} \left( \frac{2x+2}{3} \right)$   
 $- \frac{3}{4} \log (4x^2 + 8x + 13) + c$

24. If the equation of an ellipse is  $3x^2 + 2y^2 + 6x - 8y + 5 = 0$ , then which of the following are true?

- (a)  $e = \frac{1}{\sqrt{3}}$   
 (b) centre is  $(-1, 2)$   
 (c) foci are  $(-1, 1)$  and  $(-1, 3)$   
 (d) All of the above

25. The equation of the common tangents to the two hyperbolas  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and  $\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$ , are

- (a)  $y = \pm x \pm \sqrt{b^2 - a^2}$   
 (b)  $y = \pm x \pm \sqrt{a^2 - b^2}$   
 (c)  $y = \pm x \pm \sqrt{a^2 + b^2}$   
 (d)  $y = \pm x \pm (a^2 - b^2)$

26. Domain of the function  $f(x) = \log_x \cos x$ , is

- (a)  $\left( -\frac{\pi}{2}, \frac{\pi}{2} \right) - \{1\}$       (b)  $\left[ -\frac{\pi}{2}, \frac{\pi}{2} \right] - \{1\}$   
 (c)  $\left( -\frac{\pi}{2}, \frac{\pi}{2} \right)$       (d) None of these

27. Range of the function  $y = \sin^{-1} \left( \frac{x^2}{1+x^2} \right)$ , is

- (a)  $\left( 0, \frac{\pi}{2} \right)$       (b)  $\left[ 0, \frac{\pi}{2} \right)$   
 (c)  $\left[ 0, \frac{\pi}{2} \right]$       (d)  $\left[ 0, \frac{\pi}{2} \right]$

28. If  $x = \sec \theta - \cos \theta$ ,  $y = \sec^n \theta - \cos^n \theta$ , then

- $(x^2 + 4) \left( \frac{dy}{dx} \right)^2$  is equal to  
 (a)  $n^2 (y^2 - 4)$       (b)  $n^2 (4 - y^2)$   
 (c)  $n^2 (y^2 + 4)$       (d) None of these

29. If  $y = \sqrt{x + \sqrt{y + \sqrt{x + \sqrt{y + \dots \infty}}}}$ , then  $\frac{dy}{dx}$  is equal to

- (a)  $\frac{y+x}{y^2 - 2x}$       (b)  $\frac{y^3 - x}{2y^2 - 2xy - 1}$   
 (c)  $\frac{y^3 + x}{2y^2 - x}$       (d) None of these

30. If  $\int_1^x \frac{dt}{|t| \sqrt{t^2 - 1}} = \frac{\pi}{6}$ , then  $x$  can be equal to

- (a)  $\frac{2}{\sqrt{3}}$       (b)  $\sqrt{3}$   
 (c) 2      (d) None of these

31. The area bounded by the curve  $y = |\sin x|$ ,  $x$ -axis and the lines  $|x| = \pi$ , is

- (a) 2 sq unit      (b) 1 sq unit  
 (c) 4 sq unit      (d) None of these

32. The degree of the differential equation of all curves having normal of constant length  $c$  is

- (a) 1      (b) 3  
 (c) 4      (d) None of these

33. If  $\vec{a} = 2\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $\vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$  and  $\vec{c} = 3\hat{i} + \hat{j}$ , then  $\vec{a} + t\vec{b}$  is perpendicular to  $\vec{c}$ , if  $t$  is equal to

- (a) 2      (b) 4  
 (c) 6      (d) 8

34. The distance between the line  $\vec{r} = 2\hat{\mathbf{i}} - 2\hat{\mathbf{j}} + 3\hat{\mathbf{k}} + \lambda(\hat{\mathbf{i}} - \hat{\mathbf{j}} + 4\hat{\mathbf{k}})$  and the plane  $\vec{r} \cdot (\hat{\mathbf{i}} + 5\hat{\mathbf{j}} + \hat{\mathbf{k}}) = 5$ , is
- (a)  $\frac{10}{3}$       (b)  $\frac{10}{\sqrt{3}}$   
 (c)  $\frac{10}{3\sqrt{3}}$       (d)  $\frac{10}{9}$
35. The equation of sphere concentric with the sphere  $x^2 + y^2 + z^2 - 4x - 6y - 8z - 5 = 0$  and which passes through the origin, is
- (a)  $x^2 + y^2 + z^2 - 4x - 6y - 8z = 0$   
 (b)  $x^2 + y^2 + z^2 - 6y - 8z = 0$   
 (c)  $x^2 + y^2 + z^2 = 0$   
 (d)  $x^2 + y^2 + z^2 - 4x - 6y - 8z - 6 = 0$
36. If the lines  $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z-1}{4}$  and  $\frac{x-3}{1} = \frac{y-k}{2} = \frac{z}{1}$  intersect, then the value of  $k$ , is
- (a)  $\frac{3}{2}$       (b)  $\frac{9}{2}$   
 (c)  $-\frac{2}{9}$       (d)  $-\frac{3}{2}$
37. The two curves  $y = 3^x$  and  $y = 5^x$  intersect at an angle
- (a)  $\tan^{-1}\left(\frac{\log 3 - \log 5}{1 + \log 3 \log 5}\right)$   
 (b)  $\tan^{-1}\left(\frac{\log 3 + \log 5}{1 - \log 3 \log 5}\right)$   
 (c)  $\tan^{-1}\left(\frac{\log 3 + \log 5}{1 + \log 3 \log 5}\right)$   
 (d)  $\tan^{-1}\left(\frac{\log 3 - \log 5}{1 - \log 3 \log 5}\right)$
38. The equation  $\lambda x^2 + 4xy + y^2 + \lambda x + 3y + 2 = 0$  represents a parabola, if  $\lambda$  is
- (a) 0      (b) 1  
 (c) 2      (d) 4
39. If two circles  $2x^2 + 2y^2 - 3x + 6y + k = 0$  and  $x^2 + y^2 - 4x + 10y + 16 = 0$  cut orthogonally, then the value of  $k$  is
- (a) 41      (b) 14  
 (c) 4      (d) 1
40. If  $A(-2, 1)$ ,  $B(2, 3)$  and  $C(-2, -4)$  are three points. Then, the angle between  $BA$  and  $BC$  is
- (a)  $\tan^{-1}\left(\frac{2}{3}\right)$       (b)  $\tan^{-1}\left(\frac{3}{2}\right)$   
 (c)  $\tan^{-1}\left(\frac{1}{3}\right)$       (d)  $\tan^{-1}\left(\frac{1}{2}\right)$

## Answer Key

1. d	2. c	3. a	4. a	5. c	6. b	7. c	8. d	9. c	10. c
11. c	12. d	13. c	14. b	15. a	16. a	17. a	18. c	19. c	20. a
21. d	22. a	23. a	24. d	25. b	26. d	27. b	28. c	29. d	30. a
31. c	32. d	33. d	34. c	35. a	36. b	37. a	38. d	39. c	40. a