egnado to insburPOST-GRADUATE COURSEd world (d)

Term End Examination — December, 2007

frequency distributimon.M

BASIC STATISTICAL CONCEPT & TOOLS

PAPER VII

41 ovoda 41 - 21 21 - 01 Time — 2 hours

W-75

Full marks—50

P.T.O.

(Weightage of marks—80%)

Special credit will be given for accuracy and relevance in the answer. Marks will be deducted for incorrect spelling, untidy work and illegible handwriting. The weightage for each question has been indicated in the margin.

Rank by Judge-II aluboM

Answer any two questions.

1.(a) Lives of two models of refrigerators in a recent survey

Life	No. of refrigerators				
(No. of years)	Mient, between	Model A Model B			
0 - 2		d deviction	(c) $\hat{\mathbf{g}}$ tandar		
2 -4 = 1		V 16 d s do a	(d) Nost p		
4 - 6		13	12		
formul8 - Salcula					
8 - 10		nedw v to	ealue o		
10 - 12 040		04 2200	1		

What is the average life of each model of these refrigerators? Which model has greater uniformity?

 $N = 1000, T_A = 470, T_B = 020, T_{AB} = 320.$

PG CO-VII (2)

(b) Show that standard deviation is independent of change of origin but depends on scale. 8½+4

2.(a) Compute suitable averages from the following frequency distribution:

Monthly income : JADITETTATE DIZAB

(1000 Rs.)	be	low 2	2 - 4	4 - 6	
No. of famil	ies	5	12	30	
6 - 8	8 - 10	10 - 12	12 - 14	above	14
42	48	31	18	10	

8

(b) Compute Sperman's rank correlation from the following data and comment:

Candidate: A B C D E F G H

Rank by Judge-I: 5 2 8 1 4 6 3 7

Rank by Judge-II: 4 5 7 3 2 8 1 6

41/2

- 3. You are given that the variance of x is 9. The regression equations are 8x 10y + 66 = 0 and 40x 18y = 214. Find
 - (a) Average values of x and y.
 - (b) Correlation coefficient between two variables.
 - (c) Standard deviation of y.
 - (d) Most probable value of y when x = 25.

 $12\frac{1}{2}$

4.(a) Using suitable interpolation formula, calculate the value of y when x = 25.

x: 7 11 15 19 23 27 $y: 20256 \ 20625 \ 21296 \ 22407 \ 24098 \ 26511$

8

(b) Find whether attributes A and B are independent, positively or negatively associated in the cases N = 1000, $f_A = 470$, $f_B = 620$, $f_{AB} = 320$. $4\frac{1}{2}$

8. Write short notes all sluboM om the following

Answer any two questions. (6)

5.(a) What is chain base method of construction of index numbers and how does it differ from fixed base method.

(b) Calculate Laspeyres' and Paasche's price lindex numbers from the following data:

Commodity	Base Year		Current Year	
	Price	quantity	Price	quantity
A	50	200	65	180
В	70	300	82	275
C	60	220	78	250
D	20	160	27	170
			•	, 8

6. Compute seasonal indices by suitable method from the following data:

Year / Quarter	I	II	III	IV
2004	135	185	160	229
2005	143	205	180	259
2006	152	231	195	279

121/2

7. The values of sample mean (\bar{x}) and the ranges (R) for 10 samples of size 5 each are given below. Draw appropriate mean chart and comment on the state of control of the process.

Sample No.: 1 2 3 4 5 6 7 8 9 10 \overline{x} : 25 29 23 26 28 31 25 22 29 24 R: 6 5 4 6 5 4 7 8 4 6

121/2

P.T.O.

PG CO-VII (4)

8. Write short notes on any 3 from the following:

(a) Index number formulae.

(b) Multiple and partial correlation.

(c) Scatter diagram.

(d) Total quality management.

(e) Moving average method.

(f) Relative dispersion.

 $3\times4+\frac{1}{2}$