

1. If  $f : [2, 3] \rightarrow R$  is defined by  $x^3 + 3x - 2$ , then the range  $f(x)$  is

contained in the interval

- (a)  $[1, 12]$       (b)  $[12, 34]$   
(c)  $[35, 50]$       (d)  $[-12, 12]$

2. The number of subsets of  $\{1, 2, 3, \dots, 9\}$  containing at least one odd number is

- (a) 324      (b) 396  
(c) 496      (d) 512

3. A binary sequence is an array of 0's and 1's. The number of  $n$ -digit binary sequences which contain even number of 0's is

- (a)  $2^{n-1}$       (b)  $2^n - 1$   
(c)  $2^{n-1} - 1$       (d)  $2^n$

4. If  $x$  is numerically so small so that  $x^2$  and higher powers of  $x$  can be neglected, then

$$\left(1 + \frac{2x}{3}\right)^{3/2} \cdot (32 + 5x)^{-1/5}$$

is approximately equal to

(a)  $\frac{32 + 31x}{64}$

(b)  $\frac{31 + 32x}{64}$

(c)  $\frac{31 - 32x}{64}$

(d)  $\frac{1 - 2x}{64}$

5. The roots of

$$(x - a)(x - a - 1) + (x - a - 1)(x - a - 2) + (x - a)(x - a - 2) = 0$$

$a \in R$  are always

- (a) equal      (b) imaginary  
(c) real and distinct      (d) rational and equal

6. Let  $f(x) = x^2 + ax + b$ , where  $a, b \in R$ . If  $f(x) = 0$  has all its roots imaginary, then the roots of  $f(x) + f'(x) + f''(x) = 0$  are

- (a) real and distinct      (b) imaginary  
(c) equal      (d) rational and equal

7. If  $f(x) = 2x^4 - 13x^2 + ax + b$  is divisible by  $x^2 - 3x + 2$ , then  $(a, b)$  is equal to

- (a)  $(-9, -2)$       (b)  $(6, 4)$   
(c)  $(9, 2)$       (d)  $(2, 9)$

8. If  $x, y, z$  are all positive and are the  $p$ th,  $q$ th and  $r$ th terms of a geometric progression respectively, then the value of the determinant  
[www.jobssalart.in](http://www.jobssalart.in)

$$\begin{vmatrix} \log x & p & 1 \\ \log y & q & 1 \\ \log z & r & 1 \end{vmatrix}$$

equals

- (a)  $\log xyz$       (b)  $(p-1)(q-1)(r-1)$   
 (c)  $pqr$       (d) 0

9. The locus of  $z$  satisfying the inequality  

$$\left| \frac{z+2i}{2z+i} \right| < 1$$
, where  $z = x + iy$ , is

- (a)  $x^2 + y^2 < 1$       (b)  $x^2 - y^2 < 1$   
 (c)  $x^2 + y^2 > 1$       (d)  $2x^2 + 3y^2 < 1$

10. If  $n$  is an integer which leaves remainder one when divided by three, then  $(1 + \sqrt{3}i)^n + (1 - \sqrt{3}i)^n$  equals

- (a)  $-2^{n+1}$       (b)  $2^{n+1}$   
 (c)  $-(-2)^n$       (d)  $-2^n$

11. The period of  $\sin^4 x + \cos^4 x$  is

- (a)  $\frac{\pi^4}{2}$       (b)  $\frac{\pi^2}{2}$   
 (c)  $\frac{\pi}{4}$       (d)  $\frac{\pi}{2}$

12. If  $3 \cos x \neq 2 \sin x$ , then the general solution of  $\sin^2 x - \cos 2x = 2 - \sin 2x$  is

- (a)  $n\pi + (-1)^n \frac{\pi}{2}, n \in \mathbb{Z}$   
 (b)  $\frac{n\pi}{2}, n \in \mathbb{Z}$   
 (c)  $(4n \pm 1) \frac{\pi}{2}, n \in \mathbb{Z}$   
 (d)  $(2n - 1)\pi, n \in \mathbb{Z}$

13.  $\cos^{-1}\left(\frac{-1}{2}\right) - 2\sin^{-1}\left(\frac{1}{2}\right) + 3\cos^{-1}\left(\frac{-1}{\sqrt{2}}\right) - 4\tan^{-1}(-1)$  equals

- (a)  $\frac{19\pi}{12}$       (b)  $\frac{35\pi}{12}$   
 (c)  $\frac{47\pi}{12}$       (d)  $\frac{43\pi}{12}$

14. In a  $\Delta ABC$

$$\frac{(a+b+c)(b+c-a)(c+a-b)(a+b-c)}{4b^2c^2}$$

equals

- (a)  $\cos^2 A$       (b)  $\cos^2 B$   
 (c)  $\sin^2 A$       (d)  $\sin^2 B$

15. The angle between the lines whose direction cosines satisfy the equations  $l+m+n=0$ ,  $l^2+m^2-n^2=0$  is

- (a)  $\frac{\pi}{6}$       (b)  $\frac{\pi}{4}$   
 (c)  $\frac{\pi}{3}$       (d)  $\frac{\pi}{2}$

16. If  $m_1, m_2, m_3$  and  $m_4$  are respectively the magnitudes of the vectors

$$\vec{a}_1 = 2\hat{i} - \hat{j} + \hat{k}, \quad \vec{a}_2 = 3\hat{i} - 4\hat{j} - 4\hat{k},$$

$$\vec{a}_3 = \hat{i} + \hat{j} - \hat{k} \text{ and } \vec{a}_4 = -\hat{i} + 3\hat{j} + \hat{k},$$

then the correct order of  $m_1, m_2, m_3$  and  $m_4$  is

- (a)  $m_3 < m_1 < m_4 < m_2$   
 (b)  $m_3 < m_1 < m_2 < m_4$   
 (c)  $m_3 < m_4 < m_1 < m_2$   
 (d)  $m_3 < m_4 < m_2 < m_1$

17. If  $X$  is a binomial variate with the range  $\{0, 1, 2, 3, 4, 5, 6\}$  and  $P(X=2)=4P(X=4)$ , then the parameter  $p$  of  $X$  is

- (a)  $\frac{1}{3}$       (b)  $\frac{1}{2}$   
 (c)  $\frac{2}{3}$       (d)  $\frac{3}{4}$

18. The area (in square unit) of the circle which touches the lines  $4x + 3y = 15$  and  $4x + 3y = 5$  is

- (a)  $4\pi$       (b)  $3\pi$   
 (c)  $2\pi$       (d)  $\pi$

19. The area (in square unit) of the triangle formed by  $x + y + 1 = 0$  and the pair of straight lines  $x^2 - 3xy + 2y^2 = 0$  is

- (a)  $\frac{7}{12}$       (b)  $\frac{5}{12}$   
 (c)  $\frac{1}{12}$       (d)  $\frac{1}{6}$

20. The pairs of straight lines  $x^2 - 3xy + 2y^2 = 0$  and  $x^2 - 3xy + 2y^2 + x - 2 = 0$  form a

- (a) square but not rhombus  
 (b) rhombus  
 (c) parallelogram  
 (d) rectangle but not a square

21. The equations of the circle which pass through the origin and makes intercepts of lengths 4 and 8 on the  $x$  and  $y$ -axes respectively are

- (a)  $x^2 + y^2 \pm 4x \pm 8y = 0$   
 (b)  $x^2 + y^2 \pm 2x \pm 4y = 0$   
 (c)  $x^2 + y^2 \pm 8x \pm 16y = 0$   
 (d)  $x^2 + y^2 \pm x \pm y = 0$

22. The point  $(3, -4)$  lies on both the circles

$$\text{www.jobalert.in } x^2 + y^2 - 2x + 8y + 13 = 0$$

$$\text{and } x^2 + y^2 - 4x + 6y + 11 = 0$$

Then, the angle between the circles is

$$(a) 60^\circ \quad (b) \tan^{-1} \left( \frac{1}{2} \right)$$

$$(c) \tan^{-1} \left( \frac{3}{5} \right) \quad (d) 135^\circ$$

23. The equation of the circle which passes through the origin and cuts orthogonally each of the circles  $x^2 + y^2 - 6x + 8 = 0$  and  $x^2 + y^2 - 2x - 2y = 7$  is

$$(a) 3x^2 + 3y^2 - 8x - 13y = 0 \\ (b) 3x^2 + 3y^2 - 8x + 29y = 0 \\ (c) 3x^2 + 3y^2 + 8x + 29y = 0 \\ (d) 3x^2 + 3y^2 - 8x - 29y = 0$$

24. The number of normals drawn to the parabola  $y^2 = 4x$  from the point  $(1, 0)$  is

$$(a) 0 \quad (b) 1 \\ (c) 2 \quad (d) 3$$

25. If the circle  $x^2 + y^2 = a^2$  intersects the hyperbola  $xy = c^2$  in four points  $(x_i, y_i)$ , for  $i = 1, 2, 3$  and  $4$ , then  $y_1 + y_2 + y_3 + y_4$  equals

$$(a) 0 \quad (b) c \\ (c) a \quad (d) c^4$$

26. The mid point of the chord  $4x - 3y = 5$  of the hyperbola  $2x^2 - 3y^2 = 12$  is

$$(a) \left( 0, -\frac{5}{3} \right) \quad (b) (2, 1) \\ (c) \left( \frac{5}{4}, 0 \right) \quad (d) \left( \frac{11}{4}, 2 \right)$$

27. The perimeter of the triangle with vertices at  $(1, 0, 0)$ ,  $(0, 1, 0)$  and  $(0, 0, 1)$  is

$$(a) 3 \quad (b) 2 \\ (c) 2\sqrt{2} \quad (d) 3\sqrt{2}$$

28. If a line in the space makes angle  $\alpha, \beta$  and  $\gamma$  with the coordinate axes, then

$$\cos 2\alpha + \cos 2\beta + \cos 2\gamma + \sin^2 \alpha + \sin^2 \beta \\ + \sin^2 \gamma \text{ equals}$$

$$(a) -1 \quad (b) 0 \\ (c) 1 \quad (d) 2$$

29. The radius of the sphere

$$x^2 + y^2 + z^2 = 12x + 4y + 3z \text{ is}$$

$$(a) 13/2 \quad (b) 13 \\ (c) 26 \quad (d) 52$$

30.  $\lim_{x \rightarrow \infty} \left( \frac{x+5}{x+2} \right)^{x+3}$  equals

$$(a) e \quad (b) e^2 \\ (c) e^3 \quad (d) e^5$$

31. If  $f : R \rightarrow R$  is defined by

$$f(x) = \begin{cases} \frac{2 \sin x - \sin 2x}{2x \cos x}, & \text{if } x \neq 0 \\ a, & \text{if } x = 0 \end{cases}$$

then the value of  $a$  so that  $f$  is continuous at  $0$  is

$$(a) 2 \quad (b) 1 \\ (c) -1 \quad (d) 0$$

32.  $x = \cos^{-1} \left( \frac{1}{\sqrt{1+t^2}} \right)$ ,  $y = \sin^{-1} \left( \frac{t}{\sqrt{1+t^2}} \right) \Rightarrow \frac{dy}{dx}$

is equal to

$$(a) 0 \quad (b) \tan t \\ (c) 1 \quad (d) \sin t \cos t$$

33.  $\frac{d}{dx} \left[ a \tan^{-1} x + b \log \left( \frac{x-1}{x+1} \right) \right] = \frac{1}{x^4 - 1}$

$\Rightarrow a - 2b$  is equal to

$$(a) 1 \quad (b) -1 \\ (c) 0 \quad (d) 2$$

34.  $y = e^{a \sin^{-1} x} \Rightarrow (1-x^2)y_{n+2} - (2n+1)xy_{n+1}$  is equal to

$$(a) -(n^2 + a^2)y_n \quad (b) (n^2 - a^2)y_n \\ (c) (n^2 + a^2)y_n \quad (d) -(n^2 - a^2)y_n$$

35. The function  $f(x) = x^3 + ax^2 + bx + c$ ,  $a^2 \leq 3b$  has

$$(a) \text{one maximum value} \\ (b) \text{one minimum value} \\ (c) \text{no extreme value} \\ (d) \text{one maximum and one minimum value}$$

36.  $\int \left( \frac{2 - \sin 2x}{1 - \cos 2x} \right) e^x dx$  is equal to

$$(a) -e^x \cot x + c \quad (b) e^x \cot x + c \\ (c) 2e^x \cot x + c \quad (d) -2e^x \cot x + c$$

37. If  $I_n = \int \sin^n x dx$ , then  $nI_n - (n-1)I_{n-2}$  equals

$$(a) \sin^{n-1} x \cos x \\ (b) \cos^{n-1} x \sin x \\ (c) -\sin^{n-1} x \cos x \\ (d) -\cos^{n-1} x \sin x$$

38. The line  $x = \frac{\pi}{4}$  divides the area of the region

bounded by  $y = \sin x$ ,  $y = \cos x$  and  $x$ -axis  
 $\left(0 \leq x \leq \frac{\pi}{2}\right)$  into two regions of areas  $A_1$  and  $A_2$ .

Then  $A_1 : A_2$  equals

- |           |           |
|-----------|-----------|
| (a) 4 : 1 | (b) 3 : 1 |
| (c) 2 : 1 | (d) 1 : 1 |

39. The solution of the differential equation

$$\frac{dy}{dx} = \sin(x+y) \tan(x+y) - 1$$

- (a)  $\text{cosec}(x+y) + \tan(x+y) = x + c$
- (b)  $x + \text{cosec}(x+y) = c$
- (c)  $x + \tan(x+y) = c$
- (d)  $x + \sec(x+y) = c$

40. If  $p \Rightarrow (\sim p \vee q)$  is false, the truth value of  $p$  and  $q$  are respectively

- (a) F, T
- (b) F, F
- (c) T, F
- (d) T, T

## Answer Key

1. b	2. c	3. a	4. a	5. c	6. b	7. c	8. d	9. c	10. c
11. d	12. c	13. d	14. c	15. c	16. a	17. a	18. d	19. c	20. c
21. a	22. d	23. b	24. b	25. a	26. b	27. d	28. c	29. a	30. c
31. d	32. c	33. b	34. c	35. c	36. a	37. c	38. d	39. b	40. c