

PART A

1. A physiological disorder X always leads to the disorder Y. However, disorder Y may occur by itself. A population shows 4% incidence of disorder Y. Which of the following inferences is valid?

1. 4% of the population suffers from both X & Y
2. Less than 4% of the population suffers from X
3. At least 4% of the population suffers from X
4. There is no incidence of X in the given population

2. Exposing an organism to a certain chemical can change nucleotide bases in a gene, causing mutation. In one such mutated organism if a protein had only 70% of the primary amino acid sequence, which of the following is likely?






1. Mutation broke the protein
2. The organism could not make amino acids
3. Mutation created a terminator codon
4. The gene was not transcribed

3. The speed of a car increases every minute as shown in the following Table. The speed at the end of the 19th minute would be .


Time (minutes)	Speed (m/sec)
1	1.5
2	3.0
3	4.5
.	.
.	.
24	36.0
25	37.5

1. 26.5
2. 28.0
3. 27.0
4. 28.5

4. If V_{input} is applied to the circuit shown, the output would be



5. Water is dripping out of a tiny hole at the bottom of three flasks whose base diameter is the same, and are initially filled to the same height, as shown




Which is the correct comparison of the rate of fall of the volume of water in the three flasks?

1. A fastest, B slowest
2. B fastest, A slowest
3. B fastest, C slowest
4. C fastest, B slowest

6. A reference material is required to be prepared with 4 ppm calcium. The amount of $CaCO_3$ (molecular weight = 100) required to prepare 1000 g of such a reference material is.

1. 10 μ g
2. 4 μ g
3. 4 mg
4. 10 mg

7.



The normal boiling point of a solvent (whose vapour pressure curve is shown in the figure) on a planet whose normal atmospheric pressure is 3 bar, is about

1. 100 K
2. 273 K
3. 400 K
4. 500 K

8. How many σ bonds are present in the following molecule?



1. 4
2. 6
3. 10
4. 13


9. The reason for the hardness of diamond is

1. extended covalent bonding
2. layered structure
3. formation of cage structures
4. formation of tubular structures


10. The acidity of normal rain water is due to

1. SO_2
2. CO_2
3. NO_2
4. NO

11. A ball is dropped from a height h above the surface of the earth. Ignoring air drag, the curve that best represents its variation of acceleration is



12.



The cumulative profits of a company since its inception are shown in the diagram. If the net worth of the company at the end of 4th year is 99 crores, the principal it had started with was

1. 9.9 crores
2. 91 crores
3. 90 crores
4. 9.0 crores

13. Diabetic patients are advised a low glycaemic index diet. The reason for this is

1. They require less carbohydrate than healthy individuals
2. They cannot assimilate ordinary carbohydrates
3. They need to have slow, but sustained release of glucose in their blood stream
4. They can tolerate lower, but not higher than normal blood sugar levels

14. Standing on a polished stone floor one feels colder than on a rough floor of the same stone. This is because

1. Thermal conductivity of the stone depends on the surface smoothness
2. Specific heat of the stone changes by polishing it
3. The temperature of the polished floor is lower than that of the rough floor
4. There is greater heat loss from the soles of the feet when in contact with the polished floor than with the rough floor


15. Popular use of which of the following fertilizers increases the acidity of soil?

1. Potassium Nitrate
2. Urea
3. Ammonium sulphate
4. Superphosphate of lime

16. If the atmospheric concentration of carbon dioxide is doubled and there are favourable conditions of water, nutrients, light and temperature, what would happen to water requirement of plants?

1. It decreases initially for a short time and then returns to the original value
2. It increases
3. It decreases
4. It increases initially for a short time and then returns to the original value

17.




The graph represents the depth profile of temperature in the open ocean; in which region this is likely to be prevalent?

1. Tropical region
2. Equatorial region
3. Polar region
4. Sub-tropical region

18. Glucose molecules diffuse across a cell of diameter d in time τ . If the cell diameter is tripled, the diffusion time would

1. increase to 9τ
2. decrease to $\tau/3$
3. increase to 3τ
4. decrease to $\tau/9$

19. Identify the figure which depicts a first order reaction.



20. Which of the following particles has the largest range in a given medium if their initial energies are the same?

1. alpha
2. electron
3. positron
4. gamma

PART B

21. According to crystal field theory, Ni^{2+} can have two unpaired electrons in
1. octahedral geometry only
 2. square-planar geometry only
 3. tetrahedral geometry only
 4. both octahedral and tetrahedral geometry
22. $[\text{Ni}(\text{CN})_4]^{2-}$ and $[\text{NiCl}_4]^{2-}$ complex ions are
1. both diamagnetic
 2. both paramagnetic
 3. diamagnetic and paramagnetic respectively
 4. antiferromagnetic and diamagnetic respectively
23. Which of the following spectroscopic techniques will be useful to distinguish between M-SCN and M-NCS binding modes?
1. NMR
 2. IR
 3. EPR
 4. Mass
24. Which of the following compounds show a charge-transfer band?
1. lanthanum nitrate
 2. ceric ammonium nitrate
 3. manganese(II) acetate
 4. copper(II) sulphate pentahydrate
25. Among SF_4 , BF_4^- , XeF_4 , and ICl_4^- the number of species having two lone pair of electrons on the central atom according to VSEPR theory is:
1. 2
 2. 3
 3. 4
 4. 0
26. The FALSE statement for a polarographic measurement procedure is
1. O_2 is removed
 2. dropping mercury electrode is working electrode
 3. I_d is proportional to concentration of electroactive species
 4. Residual current is made zero by adding supporting electrolyte
27. The ligand system present in vitamin B_{12} is
1. porphyrin
 2. corrin
 3. phthalocyanine
 4. crown ether
28. Which one of the following exhibits rotational spectra?
1. H_2
 2. N_2
 3. CO
 4. CO_2
29. In Ziegler-Natta catalysis the commonly used catalyst system is:
1. TiCl_4 , $\text{Al}(\text{C}_2\text{H}_5)_3$
 2. $(\eta^5\text{-Cp})_2\text{TiCl}_2$, $\text{Al}(\text{OEt})_3$
 3. $\text{VO}(\text{acac})_2$, $\text{Al}_2(\text{CH}_3)_6$
 4. TiCl_4 , BF_3
30. Oxidation occurs very easily in case of
1. $(\eta^5\text{-C}_5\text{H}_5)_2\text{Fe}$
 2. $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}$
 3. $(\eta^5\text{-C}_5\text{H}_5)_2\text{Ru}$
 4. $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}^+$
31. Complex in which organic ligand is having only σ -bond with metal is:
1. $\text{W}(\text{CH}_3)_6$
 2. $(\eta^5\text{-C}_5\text{H}_5)_2\text{Fe}$
 3. $\text{K}[\text{PtCl}_3(\text{C}_2\text{H}_4)]$
 4. $(\eta^6\text{-C}_6\text{H}_6)_2\text{Ru}$

32. In molecules H_2O , NH_3 and CH_4

1. the bond angles are same
2. the bond distances are same
3. the hybridizations are same
4. the shapes are same

33. The correct order of stability of difluorides is

1. $\text{GeF}_2 > \text{SiF}_2 > \text{CF}_2$
2. $\text{CF}_2 > \text{SiF}_2 > \text{GeF}_2$
3. $\text{SiF}_2 > \text{GeF}_2 > \text{CF}_2$
4. $\text{CF}_2 > \text{GeF}_2 > \text{SiF}_2$

34. The number of possible isomers for $[\text{Ru}(\text{bpy})_2\text{Cl}_2]$ is (bpy = 2,2'-bipyridine)

1. 2
2. 3
3. 4
4. 5

35. The species ^{19}Ne and ^{14}C emit a positron and β -particle respectively. The resulting species formed are respectively

1. ^{19}Na and ^{14}B
2. ^{19}F and ^{14}N
3. ^{19}Na and ^{14}N
4. ^{19}F and ^{14}B

36. *Cis* and *trans* complexes of the type $[\text{PtA}_2\text{X}_2]$ are distinguished by

1. Chromyl chloride test
2. Carbylamine test
3. Kurnakov test
4. Ring test

37. The term symbol of a molecule with electronic configuration $(1\sigma_g)^2(1\sigma_u)^2(2\sigma_g)^2(2\sigma_u)^2(1\pi_u)^1(1\pi_u)^1$ is

1. $^1\Sigma_g^+$
2. $^3\Sigma_g^-$
3. $^1\Sigma_g^-$
4. $^3\Sigma_g^+$

38. A process is carried out at constant volume and at constant entropy. It will be spontaneous if:

1. $\Delta G < 0$
2. $\Delta H < 0$
3. $\Delta U < 0$
4. $\Delta A < 0$

39. The half life of a zero order reaction ($A \rightarrow P$) is given by (k = rate constant):

1. $t_{1/2} = \frac{[A]_0}{2k}$
2. $t_{1/2} = \frac{2.303}{k}$
3. $t_{1/2} = \frac{[A]_0}{k}$
4. $t_{1/2} = \frac{1}{k[A]_0}$

40. For an aqueous solution at 25°C , the Debye-Hückel limiting law is given by

1. $\log \gamma_{\pm} = 0.509 |Z_+ Z_-| \sqrt{\mu}$
2. $\log \gamma_{\pm} = 0.509 |Z_+ Z_-| \mu$
3. $\log \gamma_{\pm} = -0.509 |Z_+ Z_-| \sqrt{\mu}$
4. $\log \gamma_{\pm} = -0.509 |Z_+ Z_-| \mu^2$

41. The microwave spectrum of a molecule yields three rotational constants. The molecule is

1. prolate symmetric top
2. spherical top
3. asymmetric top
4. oblate symmetric top


42. The Q band in the vibrational spectrum of acetylene is observed in the

1. C - C stretching mode
2. C - H symmetric stretching mode
3. bending mode
4. C - H antisymmetric stretching mode

43. The Stark splitting for a given field is larger for a molecule AX as compared to BX. Which one of the following is true? (μ is the dipole moment)


1. $\mu_{AX} = \mu_{BX}$
2. $\mu_{AX} > \mu_{BX}$
3. $\mu_{AX} < \mu_{BX}$
4. $\mu_{BX} = 2\mu_{AX}$

44. The adsorption of a gas on a solid surface exhibits the following isotherm. Which one of the following statements is true?



1. Heat of adsorption is independent of coverage
2. Adsorption is multilayer
3. Heat of adsorption varies monotonically with coverage
4. Heat of adsorption varies exponentially with coverage

45. In a chemical reaction



the total pressure at equilibrium is 6 atm. The value of equilibrium constant is

1. 1/2
2. 9
3. 1
4. 36

46. A molecule, AX, has a vibrational energy of 1000 cm^{-1} and rotational energy of 10 cm^{-1} . Another molecule, BX, has a vibrational energy of 400 cm^{-1} and rotational energy of 40 cm^{-1} . Which one of the following statements about the coupling of

vibrational and rotational motion is true?

1. The coupling is stronger in BX.
2. The coupling is stronger in AX.
3. Magnitude of coupling is same in both AX and BX.
4. There is no coupling in both AX and BX.

47. At room temperature, which molecule has the maximum rotational entropy?

1. H_2
2. O_2
3. D_2
4. N_2

48. The normalized hydrogen atom 1s wavefunction is given by

$$\psi_{1s} = \frac{1}{\sqrt{\pi}} \left(\frac{\zeta}{a_0} \right)^{3/2} e^{-\zeta r/a_0} \text{ where } \zeta = 1 \text{ and}$$

energy is -0.5 au . If we use a normalized wavefunction of the above form with $\zeta \neq 1$, the average value of energy of the ground state of hydrogen atom is

1. greater than -0.5 au
2. equal to -0.5 au
3. less than -0.5 au
4. equal to ζ times -0.5 au

49. A constant of motion is defined by the equation:

1. $[H, A] = 0$
2. $\langle [H, A] \rangle = 0$
3. $A = f(H)$
4. $A^\dagger = A$

50. The hermitian conjugate of operator d/dx , called $(d/dx)^\dagger$, is actually equal to

1. $-d/dx$
2. d/dx
3. $i(d/dx)$
4. $-i(d/dx)$


51. An ideal gas expands by following an equation $PV^a = \text{constant}$. In which case does one expect heating?

1. $3 > a > 2$
2. $2 > a > 1$
3. $0 < a < 1$
4. $-1 < a < 0$

52. If $y^2 = 4x$ and 0.1% error is incurred for x , the percentage error involved in y will be


1. 0.4
2. 0.025
3. 0.1
4. 0.05

53. The configurations at the two stereocentres in the compound given below are




1. 1R, 4R
2. 1R, 4S
3. 1S, 4R
4. 1S, 4S

54. The two compounds given below are



1. enantiomers ✓
2. identical ✓
3. diastereomers
4. regioisomers

55. A suitable catalyst for bringing out the transformation given below is




1. $\text{BF}_3 \cdot \text{Et}_2\text{O}$
2. NaOEt ✓
3. tungsten lamp
4. dibenzoyl peroxide

56. Thermolysis of allyl phenyl ether generate


1. *o*-allylphenol only
2. *o*- and *p*-allylphenols
3. *o*-, *m*- and *p*-allylphenols
4. *m*-allylphenol only

57. The major product formed in the reaction given below is




- 1.
- 2.
- 3.
- 4.

58. The most suitable reagent for the following transformation is



1. LiAlH_4
2. $\text{NH}_2\text{NH}_2 / \text{KOH}$
3. $\text{NaBH}_4 / \text{CeCl}_3$
4. Li / liq. NH_3 ✓

59. The intermediate involved in the reaction given below is




1. free radical
2. carbocation
3. carbanion
4. carbene ✓

60. In the most stable conformation of *trans*-1-*t*-butyl-3-methylcyclohexane, the substituents at C-1 and C-3, respectively, are

1. axial and equatorial
2. equatorial and equatorial
3. equatorial and axial
4. axial and axial

61. Among the carbocations given below




1. A is homoaromatic, B is antiaromatic and C is aromatic.
2. A is aromatic, B is antiaromatic and C is homoaromatic.
3. A is antiaromatic, B is aromatic and C is homoaromatic.
4. A is homoaromatic, B is aromatic and C is antiaromatic.

62. The order of carbonyl stretching frequency in the IR spectra of ketone, amide and anhydride is


1. anhydride > amide > ketone
2. ketone > amide > anhydride
3. amide > anhydride > ketone
4. anhydride > ketone > amide

63. In the mass spectrum of the compound given below, during the α -cleavage, the order of preferential loss of groups is



1. Me > C₃H₇ > Et
2. C₃H₇ > Et > Me
3. Et > Me > C₃H₇
4. Et > C₃H₇ > Me

64. The reaction given below is an example of



1. 1,3-sigmatropic hydrogen shift
2. 1,3-sigmatropic methyl shift
3. 1,5-sigmatropic hydrogen shift
4. 1,5-sigmatropic methyl shift

65. The concerted photochemical reaction between two olefins leading to a cyclobutane ring is

1. $\pi 2_s + \pi 2_a$ cycloaddition
2. $\pi 2_s + \pi 2_s$ cycloaddition
3. $\sigma 2_s + \sigma 2_s$ cycloaddition
4. $\pi 2_s + \sigma 2_a$ cycloaddition


66. Addition of BH₃ to a carbon-carbon double bond is

1. *anti*-Markovnikov *syn* addition
2. *anti*-Markovnikov *anti* addition
3. Markovnikov *syn* addition
4. Markovnikov *anti* addition

67. The absorption at λ_{\max} 279 nm ($\epsilon = 15$) in the UV spectrum of acetone is due to


1. $\pi-\pi^*$ transition
2. $n-\pi^*$ transition
3. $\sigma-\sigma^*$ transition
4. $\pi-\sigma^*$ transition

68. The reaction given below is an example of




1. E₂-elimination
2. E₁-elimination
3. *syn*-elimination
4. E_{1cb}-elimination

69. The suitable reagent for the following conversion is



1. *m*-CPBA
2. $\text{H}_2\text{O}_2 / \text{AcOH}$
3. $\text{tBuOOH} / \text{HCl}$
4. $\text{H}_2\text{O}_2 / \text{NaOH}$

70. The relative rates of solvolysis of iodides A-C are



1. $\text{C} > \text{A} > \text{B}$
2. $\text{C} > \text{B} > \text{A}$
3. $\text{B} > \text{C} > \text{A}$
4. $\text{B} > \text{A} > \text{C}$

PART C

71. Alkali metal superoxides are obtained by the reaction of

1. oxygen with alkali metals in liquid ammonia
2. water with alkali metals in liquid ammonia
3. H_2O_2 with alkali metals
4. H_2O_2 with alkali metals in liquid ammonia

72. H_2O_2 reduces

- (A) $[\text{Fe}(\text{CN})_6]^{3-}$ (B) KIO_4
(C) $\text{Ce}(\text{SO}_4)_2$ (D) SO_3^{2-}

1. A and B only
2. B and C only
3. C and D only
4. B and D only

73. Match list I (compounds) with list II (application) and select the correct answer using the codes given below the lists

	List I		List II
A.	Trisodium phosphate	(i)	Plasticizer
B.	Triarylphosphates	(ii)	Water softener
C.	Triethylphosphate	(iii)	Toothpaste
D.	Calcium hydrogen phosphate	(iv)	Insecticides

1. (A-ii) (B-i) (C-iv) (D-iii)
2. (A-i) (B-ii) (C-iv) (D-iii)
3. (A-ii) (B-iii) (C-iv) (D-i)
4. (A-iii) (B-i) (C-ii) (D-iv)

74. Among the following statements, identify the correct ones for complexes of lanthanide(III) ion.

- (A) Metal-ligand bond is significantly ionic.
(B) Complexes rarely show isomerism.
(C) The coordination number is not more than 8.
(D) The magnetic moments are not accounted even approximately by spin only value for majority of lanthanides

1. (A), (B) and (D) only.
2. (A), (B) and (C) only.
3. (B) and (C) only.
4. (A) and (D) only.

75. According to VSEPR theory, the molecule/ion having ideal tetrahedral shape is:

1. SF_4
2. SO_4^{2-}
3. S_2Cl_2
4. SO_2Cl_2

76. The complex $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$ has very light pink colour. The best reason for it is

1. the complex does not have a charge-transfer transition
2. d-d transitions here are orbital-forbidden but spin-allowed
3. d-d transitions here are orbital-allowed but spin-forbidden
4. d-d transitions here are both orbital-forbidden and spin-forbidden

77. The highest occupied MO in N_2 and O_2^+ respectively are (take x-axis as internuclear axis)

1. $\sigma 2p_x, \pi^* 2p_y$
2. $\pi 2p_y, \pi 2p_z$
3. $\sigma^* 2p_x, \sigma 2p_x$
4. $\pi^* 2p_y, \pi^* 2p_z$

78. The correct order of LMCT energies is:

1. $MnO_4^- < CrO_4^{2-} < VO_4^{3-}$
2. $MnO_4^- > CrO_4^{2-} > VO_4^{3-}$
3. $MnO_4^- > CrO_4^{2-} < VO_4^{3-}$
4. $MnO_4^- < CrO_4^{2-} > VO_4^{3-}$

79. Carboxypeptidase contains:

1. Zn(II) and hydrolyses CO_2 .
2. Zn(II) and hydrolyses peptide bonds.
3. Mg(II) and hydrolyses CO_2 .
4. Mg(II) and hydrolyses peptide bonds.

80. In the EPR spectrum of tetragonal Cu(II) complex, when $g_{\parallel} > g_{\perp} > g_e$ the unpaired electron resides in the orbital:

1. d_{xy}
2. $d_{x^2-y^2}$
3. d_z
4. d_{xz}

81. The oxidative addition and reductive elimination steps are favoured by:

1. electron rich metal centers
2. electron deficient metal centers
3. electron deficient and electron rich metal centers respectively
4. electron rich and electron deficient metal centers respectively

82. Identify the order according to increasing stability of the following organometallic compounds, $TiMe_4$, $Ti(CH_2Ph)_4$, $Ti(i-Pr)_4$ and $TiEt_4$.

(Me = methyl, Ph = phenyl, *i*-Pr = isopropyl, Et = ethyl)

1. $Ti(CH_2Ph)_4 < Ti(i-Pr)_4 < TiEt_4 < TiMe_4$
2. $TiEt_4 < TiMe_4 < Ti(i-Pr)_4 < Ti(CH_2Ph)_4$
3. $Ti(i-Pr)_4 < TiEt_4 < TiMe_4 < Ti(CH_2Ph)_4$
4. $TiMe_4 < TiEt_4 < Ti(i-Pr)_4 < Ti(CH_2Ph)_4$


83. Among the metals, Mn, Fe, Co and Ni, the ones those would react in its native form directly with CO giving metal carbonyl compounds are:

1. Co and Mn
2. Mn and Fe
3. Fe and Ni
4. Ni and Co

84. The molecule with highest number of lone-pairs and has a linear shape based on VSEPR theory is

1. CO_2
2. I_3^-
3. NO_2^-
4. NO_2^+

85. Given



A 100 ml solution is 1080 mg with respect to Ag^+ and 635 mg with respect to Cu^{2+} . If 0.1 mg Ag^+ left in the solution is considered to be the complete deposition of Ag^+ , the cathode potential, so that no copper is deposited during the process, is:

1. 0.16 V
2. 0.84 V
3. 0.31 V
4. -0.16 V

86. In the $H_2Ru_6(CO)_{18}$ cluster, containing 8-coordinated Ru centers, the hydrogen atoms are:

1. both terminal
2. one terminal and the other bridging
3. both bridging between two Ru centers
4. both bridging between three Ru centers

87. In the hydroformylation reaction, the intermediate $CH_3CH_2CH_2Co(CO)_4$:

1. forms an acyl intermediate $CH_3CH_2CH_2COCO(CO)_3$
2. forms an adduct with an olefin reactant
3. reacts with H_2
4. eliminates propane

88. **Statement I:** The sizes of Zr and Hf are similar
Statement II: Size of Hf is affected by lanthanide contraction
1. Statements I and II are correct and II is correct explanation of I
 2. Statement I and II are correct but II is not a correct explanation of I
 3. Statement I is correct and II is incorrect.
 4. Statements I and II both are incorrect.
89. Consider the compounds, (A) SnF₄, (B) SnCl₄ and (C) R₃SnCl
 The nuclear quadrupole splitting are observed for
1. (A), (B) and (C)
 2. (A) and (B) only
 3. (B) and (C) only
 4. (A) and (C) only
90. Consider two redox pairs (a) Cr(II)/Ru(III) and (b) Cr(II)/Co(III). The rate of acceleration in going from a outer-sphere to a inner-sphere mechanism is lower for (a) relative to (b). Its correct explanation is :
1. HOMO/LUMO are σ^* and σ^* respectively
 2. HOMO/LUMO are σ^* and π^* respectively
 3. HOMO/LUMO are π^* and σ^* respectively
 4. HOMO/LUMO are π^* and π^* respectively
91. The correct value of isomer shift (in Mössbauer spectra) and its explanation for Fe(II)-TPP and Fe(III)-TPP respectively from the following are:
 (TPP = tetraphenylporphyrinate)
- (A) 0.52 mms⁻¹
 - (B) 0.45 mms⁻¹,
 - (C) increase in s electron density
 - (D) decrease in s electron density
1. (A) and (D) ; (B) and (C)
 2. (A) and (C) ; (B) and (C)
 3. (B) and (D) ; (A) and (D)
 4. (B) and (D) ; (A) and (C)
92. In IR spectrum of [Co(CN)₅H]³⁻ the Co-H stretch is observed at 1840 cm⁻¹. The Co-D stretch in [Co(CN)₅D]³⁻ will appear at nearly
1. 1300 cm⁻¹
 2. 1400 cm⁻¹
 3. 1500 cm⁻¹
 4. 1600 cm⁻¹
93. For the complexes (A) [Ni(H₂O)₆]²⁺, (B) [Mn(H₂O)₆]²⁺, (C) [Cr(H₂O)₆]³⁺ and (D) [Ti(H₂O)₆]³⁺, the ideal octahedral geometry will **not** be observed in
1. (A) and (D)
 2. (C) and (D)
 3. (B) only
 4. (D) only
94. Among the following, the number of anhydrides of acids are:
 CO, NO, N₂O, B₂O₃, N₂O₅, SO₃ and P₄O₁₀.
1. 3
 2. 4
 3. 5
 4. 6
95. For a given nuclear fission reaction of ²³⁵U
- $${}_{92}^{235}\text{U} + {}_0^1\text{n} \rightarrow {}_{56}^{142}\text{Ba} + {}_{36}^{91}\text{Kr} + 3{}_0^1\text{n}$$
- the amount of energy (in kJ/mol) released during this process is (given ²³⁵U = 235.0439 amu, ¹⁴²Ba = 141.9164 amu, ⁹¹Kr = 90.9234 amu, neutron = 1.00866 amu)
1. 3.12 × 10¹²
 2. 2.8 × 10¹¹
 3. 1.0 × 10⁹
 4. 1.68 × 10¹⁰
96. The decomposition of gaseous acetaldehyde at T(K) follows second order kinetics. The half-life of this reaction is 400 s when the initial pressure is 250 Torr. What will be the rate constant (in Torr⁻¹ s⁻¹) and half-life (in s) respectively, if the initial pressure of the acetaldehyde is 200 Torr at the same temperature?
1. 10⁵ and 500
 2. 10⁻⁵ and 400
 3. 10⁻⁴ and 400
 4. 10⁻⁵ and 500

97) For an enzyme catalyzed reaction, a Lineweaver-Burk plot gave the following data:

$$\text{slope} = 40 \text{ s}$$

$$\text{intercept} = 4 \text{ (mmol dm}^{-3} \text{ s}^{-1})^{-1}$$

If the initial concentration of enzyme is $2.5 \times 10^{-9} \text{ mol dm}^{-3}$, what is the catalytic efficiency (in $\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$) of the reaction?

1. 10^5
2. 10^6
3. 10^7
4. 10^4

98. A hydrogenic orbital with radial function of the form $r^\alpha \exp[-\beta r]$ and ϕ -part as $\exp[-3i\phi]$ corresponds to

1. $n > 4, l = 3, m = 3$
2. $n = 4, l = 3, m = -3$
3. $n = 4, l = 3, m = 3$
4. $n > 4, l = 3, m = -3$

99. For an assembly of molecules (molar mass = M) at temperature T, the standard deviation of Maxwell's speed is approximately

1. $0.7 \sqrt{\frac{RT}{M}}$
2. $1.4 \sqrt{\frac{RT}{M}}$
3. $0.7 \sqrt{\frac{M}{RT}}$
4. $1.4 \sqrt{\frac{M}{RT}}$

100. The unperturbed energy levels of a system are $\epsilon_0 = 0, \epsilon_1 = 2$ and $\epsilon_2 = 4$. The second-order correction to energy for the ground state in presence of the perturbation V for which $V_{10} = 2, V_{20} = 4$ and $V_{12} = 6$, has been found to be

1. -6
2. 0
3. +6
4. -8

101) Given the character table of the point group C_{3v} .

	E	$2C_3$	$3\sigma_v$	
A_1	1	1	1	z
A_2	1	1	-1	
E	2	-1	0	(x,y)

Consider the reducible representation, Γ

	E	$2C_3$	$3\sigma_v$
Γ	6	3	0

Its irreducible components are

1. $E + 2A_1 + 2A_2$
2. $2E + A_1 + A_2$
3. $3A_1 + 3A_2$
4. $E^2 + 2A_1$

102. Refer to the character table of the point group C_{3v} given above. Find which of the following transition is forbidden

1. $a_1 \leftrightarrow a_1$
2. $a_1 \leftrightarrow e$
3. $a_2 \leftrightarrow e$
4. $a_1 \leftrightarrow a_2$

103. The electronic configuration for gadolinium (Gd) is $[\text{Xe}]4f^7 5d^1 6s^2$, where as that of Gd^{2+} is:

1. $[\text{Xe}]4f^6 5d 6s^2$
2. $[\text{Xe}]4f^6 6s^2$
3. $[\text{Xe}]4f^6 5d^1 6s^1$
4. $[\text{Xe}]4f^7 5d^1$

104) The possible J values for 3D term symbol are

1. 2
2. 3
3. 4
4. 5

105) The energy levels for cyclobutadiene are $\alpha + 2\beta, \alpha, \alpha$ and $\alpha - 2\beta$. The delocalization energy in this molecule is

1. 0
2. -4β
3. -8β
4. 4α

106. The variation of equilibrium constant (K) of a certain reaction with temperature (T)

$$\text{is } \ln k = 3.0 + \frac{2.0 \times 10^4}{T}$$

given $R = 8.3 \text{ J k}^{-1} \text{ mol}^{-1}$, the values of ΔH° and ΔS° are:

1. 166 kJ mol^{-1} and $24.9 \text{ Jk}^{-1} \text{ mol}^{-1}$
2. 166 kJ mol^{-1} and $-24.9 \text{ Jk}^{-1} \text{ mol}^{-1}$
3. -166 kJ mol^{-1} and $24.9 \text{ Jk}^{-1} \text{ mol}^{-1}$
4. -166 kJ mol^{-1} and $-24.9 \text{ Jk}^{-1} \text{ mol}^{-1}$

107. The chemical potential of component 1 in a solution of binary mixture is $\mu_1 = \mu_1^0 + RT \ln p_1$, when p_1 is the partial pressure of component 1 vapour phase. The standard state μ_1^0 is

1. Independent of temperature and pressure
2. depends on temperature and pressure both
3. depends on temperature only
4. depends on pressure only

108. Debye - Hückel screening length (κ^{-1}) is a measure of size of diffuse ion cloud around an ion, provided

$$\sqrt{\frac{2e^2 N_A}{\epsilon_0 k_B T}} \approx 30 \left(nm \sqrt{\text{mol K}^{-1}} \right)^{-1} \text{ at}$$

298K, which of the following values of κ^{-1} is true for a 0.03 molal solution for Na_2SO_4 in water ($\epsilon_r \approx 100$)?

1. $\frac{10}{9} nm$
2. $\frac{9}{10} nm$
3. $\frac{10\sqrt{2}}{9} nm$
4. $\frac{9}{10\sqrt{2}} nm$

109. If the ratio of composition of oxidized and reduced species in electrochemical cell,

$$\text{is given as } \frac{[O]}{[R]} = e^2, \text{ the correct}$$

potential difference will be



$$1. E - E^{0'} = + \frac{2RT}{nF}$$

$$2. E - E^{0'} = - \frac{2RT}{nF}$$


$$3. E - E^{0'} = \frac{RT}{nF}$$

$$4. E - E^{0'} = - \frac{RT}{nF}$$

110. If the equilibrium constants for the reactions 1 and 2



are K_1 and K_2 , the equilibrium constant for the reaction



1. $K_1 + K_2$
2. $K_1 - K_2$
3. $K_1 K_2$
4. K_1 / K_2

111. The virial expansion for a real gas can be written in either of the following forms:

$$\frac{P\bar{V}}{RT} = 1 + B_p P + C_p P^2 + \dots$$

$$= 1 + B_v V + C_v V^2 + \dots$$

If $B_v = \alpha B_p$, the value of α would be

1. PV/RT
2. RT/PV
3. PV
4. RT

112. A certain system of noninteracting particles has the single-particle partition function

$$f = A \frac{T^m}{V} \text{ where } A \text{ is some constant. The}$$

average energy per particle will be

1. $m\kappa T$
2. $A\kappa T$
3. $\kappa T/m$
4. $\kappa T/A$

113. Observe the following aqueous solutions of same compound. All the measurements are made at same wavelength and same temperature.

Solution A: The transmittance of 0.1 mol dm^{-3} using 1 cm cell is 0.5

Solution B: The optical density of 0.5 mol dm^{-3} is measured using 1 mm cell.

Solution C: The transmittance of this solution is 0.1 .

The optical density of these solutions follow the order:

($\log 20 = 1.3010$; $\log 30 = 1.4771$;
 $\log 50 = 1.6990$)

1. $A > B > C$
2. $B > C > A$
3. $B > A > C$
4. $C > A > B$

114. The rotational constant of $^{14}\text{N}_2$ is 2 cm^{-1} . The wave number of incident radiation in a Raman spectrometer is 20487 cm^{-1} . What is the wave number of first scattered Stokes line (in cm^{-1}) of $^{14}\text{N}_2$?

1. 20479
2. 20475
3. 20499
4. 20495

115. For a certain particle encountering a barrier, the tunneling probability is approximately e^{-10} . If the mass is halved and width of the barrier (rectangular) doubled, approximate value of the tunneling probability will be

1. $e^{-10/\sqrt{2}}$
2. $e^{-10\sqrt{2}}$
3. $e^{-20\sqrt{2}}$
4. e^{-10}

116. An operator A is defined as $A = -\frac{d}{dx} + x$.

Which one of the following statements is true?

1. A is a Hermitian operator
2. A^\dagger is an antihermitian operator

3. Both $A A^\dagger$ and $A^\dagger A$ are Hermitian operators

4. $A A^\dagger$ is Hermitian, but $A^\dagger A$ is antihermitian

117. Isotherm which has fractional coverage, linearly dependent on pressure at low pressures but almost independent at high pressure is called

1. BET isotherm.
2. Langmuir isotherm.
3. Freundlich isotherm.
4. Temkin isotherm.

118. A one-dimensional crystal of lattice dimension 'a' is metallic. If the structure is distorted in such a way that the lattice dimension is enhanced to '2a',

1. the electronic structure remains unchanged.
2. the width of conduction band decreases and a band gap is generated.
3. the width of conduction band increases.
4. the width of the conduction band remains unchanged.

119. For a H_2 molecule, the ground state wavefunction is $\psi(1,2) = \phi(1,2)\sigma(1,2)$ where ϕ refers to the space part and σ to the spin part. Given that $\phi(1,2) = \phi(2,1)$, the form of $\sigma(1,2)$ would be

1. $\alpha(1)\beta(2)$
2. $\alpha(2)\beta(1)$
3. $\alpha(1)\beta(2) - \alpha(2)\beta(1)$
4. $\alpha(1)\beta(2) + \alpha(2)\beta(1)$

120. There are several types of mean molar masses for polymer and they are dependent on experimental methods like: (a) osmometry, (b) light scattering and (c) sedimentation. Correct relation between mean molar masses and experimental method is

1. $\bar{M}_n \Leftrightarrow (c), \bar{M}_w \Leftrightarrow (b), \bar{M}_z \Leftrightarrow (a)$
2. $\bar{M}_n \Leftrightarrow (b), \bar{M}_w \Leftrightarrow (c), \bar{M}_z \Leftrightarrow (a)$
3. $\bar{M}_n \Leftrightarrow (a), \bar{M}_w \Leftrightarrow (b), \bar{M}_z \Leftrightarrow (c)$
4. $\bar{M}_n \Leftrightarrow (a), \bar{M}_w \Leftrightarrow (c), \bar{M}_z \Leftrightarrow (b)$


121. An organic compound ($C_7H_{12}O_2$) exhibited the following data in the 1H NMR spectrum.

δ 7.10 (1 H, dt, $J = 16$ and 7.2 Hz), 5.90 (1 H, dt, $J = 16$ and 2 Hz), 4.1 (2H, q, $J = 7.2$ Hz), 2.10 (2 H, m), 1.25 (3 H, t, $J = 7.2$ Hz), 0.90 (3 H, t, $J = 7.2$ Hz) ppm.

The compound, among the choices given below, is

- 1.
- 2.
- 3.
- 4.

122. In the broad band decoupled ^{13}C NMR spectrum, the number of signals appearing for the bicyclooctane A-C, respectively, are




1. five, four and eight
2. three, two and five
3. five, four and five
4. three, two and eight

123. In the mass spectrum of dichlorobenzene the ratio of the peaks at m/z 146, 148 and 150, is


1. 1 : 1 : 1
2. 3 : 3 : 1
3. 1 : 2 : 1
4. 9 : 6 : 1

124. The major compound X formed in the following reaction exhibited a strong absorption at ν_{max} 1765 cm^{-1} in the IR spectrum. The structure of X is.




- 1.
- 2.
- 3.
- 4.

125. The correct order of acidity of the following compounds A-C, is




1. $B > C > A$
2. $C > B > A$
3. $A > C > B$
4. $A > B > C$

126. The major product formed in the reaction sequence is




-
-
-
-

127: For the following allylic oxidation reaction, the appropriate statement, among the choices given below, is




- suitable reagent is KMnO_4 and the major product is **A**.
- suitable reagent is KMnO_4 and the major product is **B**.
- suitable reagent is SeO_2 and the major product is **A**.
- suitable reagent is SeO_2 and the major product is **B**.

128. The intermediate **A** and the major product **B** in the following conversion are




- A** is a carbocation and **B** is
- A** is a carbanion and **B** is
- A** is a free radical and **B** is
- A** is a benzyne and **B** is

129. The major product formed in the following reaction is



-
-
-
-

130. The major product formed in the following reaction is




- 1.
- 2.
- 3.
- 4.

131. The major product formed in the reaction of glucose with benzaldehyde and p-TSA is

- 1.
- 2.
- 3.
- 4.

132. Papaverine on oxidation with potassium permanganate gives a ketone, which on fusion with potassium hydroxide gives








- 1.
- 2.
- 3.
- 4.






133. The major product formed on nitration ($\text{HNO}_3/\text{H}_2\text{SO}_4$) of uridine followed by reduction with tin and HCl is

- 1.
- 2.
- 3.
- 4.

134. In the following reaction sequence, the correct structures for the major products X and Y are




135. The major product formed in the following reaction is



136. Match the following:

Column I


Column II



(i) chiral centre



(ii) chiral axis







(iii) chiral plane


(iv) helical chirality

- | | | | |
|----|-----------|-----------|-----------|
| 1. | (a)-(iii) | (b)-(ii) | (c)-(iv) |
| 2. | (a)-(iv) | (b)-(iii) | (c)-(ii) |
| 3. | (a)-(ii) | (b)-(iv) | (c)-(iii) |
| 4. | (a)-(ii) | (b)-(iii) | (c)-(iv) |



137. The gauche interaction values for Me/Me, Me/Br and Br/Br are 3.3, 0.8 and 3.0 kJ/mol, respectively. Among the following, the most stable conformation of 2,3-dibromobutane is





138. The major product formed in the reaction of (S)-1,2,4-butanetriol with 3-pentanone in the presence of a catalytic amount of p-TSA is





139. The major product formed in the following reaction is




140. The major product formed in the following transformation is



141. The major product formed in the following transformation is



142. The structures of the major products X and Y in the following transformation are




1. X = Y =
2. X = Y =
3. X = Y =
4. X = Y =

143. Match the following:

Column I	Column II
a. pyrrole	i. Pictet-Spengler
b. 1,4-dihydropyridine	ii. Chichibabin
c. isoquinoline	iii. Paal-Knorr
	iv. Hantzsch

1. (a)-(i) (b)-(ii) (c)-(iii)
 2. (a)-(ii) (b)-(iii) (c)-(iv)
 3. (a)-(iv) (b)-(i) (c)-(ii)
 4. (a)-(iii) (b)-(iv) (c)-(i)

144. Consider the following reaction:



In an experiment, 1.99 g of bromide A on reaction with ethanolic potassium hydroxide gave 1.062 g of a mixture of the olefins B and C. If the ratio of olefins B:C formed is 2:1, the yields for their formation, respectively, are

1. 60 and 30%
 2. 50 and 25%
 3. 66 and 33%
 4. 54 and 27%

145. An organic compound A ($C_8H_{16}O_2$) on treatment with an excess of methylmagnesium chloride generated two alcohols B and C, whereas reaction of A with lithium aluminum hydride generated only a single alcohol C. Compound B on treatment with an acid yielded an olefin (C_6H_{12}), which exhibited only a singlet at δ 1.6 ppm in the 1H NMR spectrum. The compound A is

- 1.
- 2.
- 3.
- 4.