






PAPER II (2007)
PART I
SECTION –I (Physics)
Straight Objective Type


This section contains 9 multiple choice questions numbered 1 to 9. Each question has 4 choices (A), (B), (C) and (D), out of which ONLY ONE is correct

Q.1 A magnetic field $\vec{B} = B_0 \hat{j}$ exists in the region $a < x < 2a$ and where $\vec{B} = -B_0 \hat{j}$ in the region $2a < x < 3a$, B_0 is a positive constant. A positive point charge moving with a velocity $\vec{v} = v_0 \hat{i}$, where v_0 is a positive constant, enters the magnetic field at $x = a$. The trajectory of the charge in this region can be like.



Ans. (A)

Q.2 A spherical portion has been removed from a solid sphere having a charge distributed uniformly in its volume as shown in the figure. The electric field inside the emptied space is



- (A) zero everywhere
 (C) non uniform


- (B) non-zero and uniform
 (D) zero only at its center

Ans. (B)


Q.3 Electrons with de-Broglie wavelength λ fall on the target in an X-ray tube. The cut-off wavelength of the emitted X-rays is

- (A) $\lambda_0 = \frac{2mc\lambda^2}{h}$ (B) $\lambda_0 = \frac{2h}{mc}$ (C) $\lambda_0 = \frac{2m^2c^2\lambda^3}{h^2}$ (D) $\lambda_0 = \lambda$ **Ans. (A)**

- Q.4** A small object of uniform density rolls up a curved surface with an initial velocity v . It reaches up to a maximum height of $\frac{3v^2}{4g}$ with respect to the initial position. The object is



- (A) ring (B) solid sphere (C) hollow sphere (D) disc **Ans. (D)**
- Q.5** Positive and negative point charges of equal magnitude are kept at $\left(0, 0, \frac{a}{2}\right)$ and $\left(0, 0, \frac{-a}{2}\right)$, respectively. The work done by the electric field when another positive point charge is moved from $(-a, 0, 0)$ to $(0, a, 0)$ is
 (A) positive
 (B) negative
 (C) zero
 (D) depends on the path connecting the initial and final positions **Ans. (C)**
- Q.6** A student performs an experiment to determine the Young's modulus of a wire, exactly 2 m long, by Searle's method. In a particular reading, the student measures the extension in the length of the wire to be 0.8 mm with an uncertainty of ± 0.05 mm at a load of exactly 1.0 kg. The student also measures the diameter of the wire to be 0.4 mm with an uncertainty of ± 0.01 mm. Take $g = 9.8 \text{ m/s}^2$ (exact). The Young's modulus obtained from the reading is
 (A) $(2.0 \pm 0.3) \times 10^{11} \text{ N/m}^2$ (B) $(2.0 \pm 0.2) \times 10^{11} \text{ N/m}^2$
 (C) $(2.0 \pm 0.1) \times 10^{11} \text{ N/m}^2$ (D) $(2.0 \pm 0.05) \times 10^{11} \text{ N/m}^2$
Ans. (B)
- Q.7** Water is filled up to a height h in a beaker of radius R as shown in the figure. The density of water is ρ , the surface tension of water is T and the atmospheric pressure is P_0 . Consider a vertical section ABCD of the water column through a diameter of the beaker. The force on water on one side of this section by water on the other side of this section has magnitude



- (A) $|2P_0Rh + \pi R^2 \rho gh - 2RT|$ (B) $|2P_0Rh + R^2 \rho gh - 2RT|$
 (C) $|P_0\pi R^2 + R\rho gh^2 - 2RT|$ (D) $|P_0\pi R^2 + R\rho gh^2 + 2RT|$
- Q.8** In the experiment to determine the speed of sound using a resonance column,
 (A) prongs of the tuning fork are kept in a vertical plane
 (B) prongs of the tuning fork are kept in a horizontal plane
 (C) in one of the two resonances observed, the length of the resonating air column is close to the wavelength of sound in air
 (D) in one of the two resonances observed, the length of the resonating air column is close to half of the wavelength of sound in air
Ans. (B)


- Q.9** A particle moves in the X –Y plane under the influence of a force such that its linear momentum is $\vec{p}(t) = A [\hat{i} \cos(kt) - \hat{j} \sin(kt)]$, where A and k are constants. The angle between the force and the momentum is
- (A) 0° (B) 30° (C) 45° (D) 90°

Ans. (D)

SECTION – II (Physics) Assertion – Reason Type

This section contains 4 questions numbered 10 to 13. Each question contains STATEMENT–1 (Assertion) and STATEMENT–2 (Reason). Each question has 4 choices (A), (B), (C) and (D) out of which ONLY ONE is correct.

- 10. STATEMENT–1**
If there is no external torque on a body about its centre of mass, then the velocity of the center of mass remains constant.
because
STATEMENT –2
The linear momentum of an isolated system remains constant.
- (A) Statement–1 is True, Statement–2 is True; Statement–2 is a correct explanation for Statement–1
(B) Statement–1 is True, Statement–2 is True; Statement–2 is **NOT** a correct explanation for Statement–1
(C) Statement–1 is True, Statement–2 is False
(D) Statement–1 is False, Statement–2 is True.
- 11. STATEMENT–1**
A vertical iron rod has a coil of wire wound over it at the bottom end. An alternating current flows in the coil. The rod goes through a conducting ring as shown in the figure. The ring can float at a certain height above the coil.



because

STATEMENT –2

In the above situation, a current is induced in the ring which interacts with the horizontal component of the magnetic field to produce an average force in the upward direction

- (A) Statement–1 is True, Statement–2 is True; Statement–2 is a correct explanation for Statement–1
(B) Statement–1 is True, Statement–2 is True; Statement–2 is **NOT** a correct explanation for Statement–1
(C) Statement–1 is True, Statement–2 is False
(D) Statement–1 is False, Statement–2 is True.

Ans. (A)

12 STATEMENT –1

A cloth covers a table. Some dishes are kept on it. The cloth can be pulled out without dislodging the dishes from the table.

because

STATEMENT –2

For every action there is an equal and opposite reaction.

- (A) Statement–1 is True, Statement–2 is True; Statement–2 is a correct explanation for Statement–1
(B) Statement–1 is True, Statement–2 is True; Statement–2 is NOT a correct explanation for Statement–1
(C) Statement–1 is True, Statement–2 is False
(D) Statement–1 is False, Statement–2 is True.

Ans. (B)

13 STATEMENT – 1

The total translational kinetic energy of all the molecules of a given mass of an ideal gas is 1.5 times the product of its pressure and its volume.

because

STATEMENT – 2

The molecules of a gas collide with each other and the velocities of them molecules change due to the collision.

- (A) Statement–1 is True, Statement–2 is True; Statement–2 is a correct explanation for Statement–1
(B) Statement–1 is True, Statement–2 is True; Statement–2 is NOT a correct explanation for Statement–1
(C) Statement–1 is True, Statement–2 is False
(D) Statement–1 is False, Statement–2 is True.


Ans. (B)

SECTION – III (Physics) Linked Comprehension Type

This section contains 2 paragraphs P14–16 and P 17–19. Based upon each paragraph, 3 multiple choice questions have to be answered. Each question has 4 choices (A), (B), (C) and (D), out of which ONLY ONE is correct.

P 17–19 : Paragraph for Question Nos. 14 to 16

The figure shows surface XY separating two transparent media, medium –1 and medium – 2. The lines ab and cd represent wavefronts of a light wave traveling in medium –1 and incident on XY. The lines ef and gh represent wavefronts of the light wave in medium –2 after refraction.



- 14 Light travels as a
(A) parallel beam in each medium
(B) convergent beam in each medium
(C) divergent beam in each medium
(D) divergent beam in one medium and convergent beam in the other medium

Ans. (A)

15. The phases of the light wave at c, d, e and f are ϕ_c , ϕ_d , ϕ_e and ϕ_f respectively. It is given that $\phi_c \neq \phi_f$:
- (A) ϕ_c cannot be equal to ϕ_d (B) ϕ_d can be equal to ϕ_e
 (C) $(\phi_d - \phi_f)$ is equal to $(\phi_c - \phi_e)$ (D) $(\phi_d - \phi_c)$ is not equal to $(\phi_f - \phi_e)$

Ans. (C)


16. Speed of light is
- (A) the same in medium -1 and medium -2 (B) larger in medium -1 than in medium -2
 (C) larger in medium -2 than in medium -1 (D) different at b and d

Ans. (B)

P 17-19 : Paragraph for Question Nos. 17 to 19

Two trains A and B are moving with speeds 20 m/s and 30 m/s respectively in the same direction on the same straight track, with B ahead of A. The engines are at the front ends. The engines of train A blows a long whistle.


Assume that the sound of the whistle is composed of components varying in frequency from $f_1 = 800$ Hz to $f_2 = 1120$ Hz, as shown in the figure. The spread in the frequency (highest frequency - lowest frequency) is thus 320 Hz. The speed of sound in still air is 340 m/s.



17. The speed of sound of the whistle is
- (A) 340 m/s for passengers in A and 310 m/s for passengers in B
 (B) 360 m/s for passengers in A and 310 m/s for passengers in B
 (C) 310 m/s for passengers in A and 360 m/s for passengers in B
 (D) 340 m/s for passengers in both the trains

Ans. (B)

18. The distribution of the sound intensity of the whistle as observed by the passengers in train A is best represented by



Ans. (A)

19. The spread of frequency as observed by the passengers in train B is
- (A) 310 Hz (B) 330 Hz (C) 350 Hz (D) 290 Hz

Ans. (A)





SECTION – IV (Physics)

Matrix – Match Type

This section contains 3 questions. Each question contains statements given in two columns which have to be matched. Statements (A ,B ,C ,D) in **Column I** have to be matched with statements (p,q,r,s) in **Column II**. The answers to these questions have to be appropriately bubbled as illustrated in the following example. If the correct matches are A –p , A –s , B –q , B –r , C –p , C –q and D –s , then the correctly bubbled 4 x 4 matrix should be as follows :

	p	q	r	s
A	<input checked="" type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input type="radio"/> s
B	<input type="radio"/> p	<input checked="" type="radio"/> q	<input checked="" type="radio"/> r	<input type="radio"/> s
C	<input type="radio"/> p	<input checked="" type="radio"/> q	<input type="radio"/> r	<input type="radio"/> s
D	<input checked="" type="radio"/> p	<input checked="" type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s

20. Two wires each carrying a steady current I are shown in four configurations in **Column I**. Some of the resulting effects are described in **Column II**. Match the statements in **Column I** with the statements in **Column II** and indicate your answer by darkening appropriate bubbles in the 4×4 matrix given in the ORS.

Column – I	Column – II
<p>(A) Point P is situated midway between the wires.</p> 	<p>(p) The magnetic fields (B) at P due to the currents in the wires are in the same direction.</p>
<p>(B) Point P is situated at the mid-point of the line joining the centres of the circular wires, which have same radii.</p> 	<p>(q) The magnetic field (B) at P due to the currents in the wires are in opposite directions</p>
<p>(C) Point P is situated at the mid-point of the line joining the centers of the circular wires, which have same radii.</p> 	<p>(r) There is no magnetic field at P.</p>
<p>(D) Point P is situated at the common center of the wires.</p> 	<p>(s) The wires repel each other.</p>

Ans. A (q,r); B – p; C – (q, r); D (q)

21. **Column I** gives some devices and **Column II** gives some processes on which the functioning of these devices depend. Match the devices in **Column I** with the processes in **Column II** and indicate your answer by darkening appropriate bubbles in the 4×4 matrix given in the O R S.

Column – I	Column – II
(A) Bimetallic strip (B) Steam engine (C) Incandescent lamp (D) Electric fuse	(p) Radiation from a hot body (q) Energy conversion (r) Melting (s) Thermal expansion of solids

Ans. A–s; B – q; C – p; D – r

22. **Column I** describes some situations in which a small object moves. **Column II** describes some characteristics of these motions. Match the situations in **Column I** with the characteristics in **Column II** and indicate your answer by darkening appropriate bubbles in the 4×4 matrix in the ORS.

Column – I	Column – II
(A) The object moves on the x–axis under a conservative force in such way that its “speed” and “position satisfy $v = c_1\sqrt{c_2 - x^2}$, where c_1 and c_2 are positive constants.	(p) The object executes a simple harmonic motion.
(B) The object moves on the x–axis in such way that its velocity and its displacement from the origin satisfy $v = -kx$, where k is a positive constant.	(q) The object does not change its direction.
(C) The object is attached to one end of a mass–less spring of a given spring constant. The other end of the spring is attached to the ceiling of an elevator. Initially everything is at rest. The elevator starts going upward with a constant acceleration a . The motion of the object is observed from the elevator during the period it maintains this acceleration.	(r) The kinetic energy of the object keeps on decreasing.
(D) The object is projected from the earth’s surface vertically upwards with a speed $2\sqrt{GM_e/R_e}$, where M_e is the mass of the earth and R_e is the radius of the earth. Neglect forces from objects other than the earth.	(s) The object can change its direction only once.

Ans. A–p ; B – q,r; C – p; D – r,q

PAPER (II)-2007

PART-II

Useful data :

Gas Constant $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ $F = 96500 \text{ C}$

Atomic Numbers : H = 1, Li = 3, B = 5, C = 6, N = 7, O = 8, F = 9, Na = 11, P = 15, S = 16, Cl = 17, Ar = 18, K = 19, V = 23, Cr = 24, Mn = 25, Fe = 26, Co = 27, Ni = 28, Cu = 29, Zn = 30, Ge = 32, Br = 35, Ag = 47, I = 53, Xe = 54, Pt = 78, Hg = 80, Pb = 82.

SECTION - I (Chemistry)

Straight Objective Type

This section contains 9 multiple choice questions numbered 23 to 31. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct.

23. For the process $\text{H}_2\text{O} (l) (1 \text{ bar}, 373 \text{ K}) \rightarrow \text{H}_2\text{O} (g) (1 \text{ bar}, 373 \text{ K})$, the correct set of thermodynamic parameters is:

(A) $\Delta G = 0, \Delta S = +ve$





(B) $\Delta G = 0, \Delta S = -ve$

(D) $\Delta G = +ve, \Delta S = 0$

(D) $\Delta G = -ve, \Delta S = +ve$

Ans. (A)

24. Among the following, the least stable resonance structure is :



Ans. (A)

25. Consider a reaction $aG + bH \rightarrow \text{Products}$. When concentration of both the reactants G and H is doubled, the rate increases by eight times. However, when concentration of G is doubled keeping the concentration of H fixed, the rate is doubled. The overall order of the reaction is :

(A) 0





(B) 1

(C) 2

(D) 3

Ans. (D)

26. Cyclohexene on ozonolysis followed by reaction with zinc dust and water gives compound E. Compound E on further treatment with aqueous KOH yields compound F. Compound F is :



Ans. (A)

27. A positron is emitted from ${}^{23}_{11}\text{Na}$. The ratio of the atomic mass and atomic number of the resulting nuclide is :

(A) 22/10




(B) 22/11

(C) 23/10

(D) 23/12

Ans. (C)

28. Among the following metal carbonyls, the C–O bond order is lowest in :



Ans. (B)

29. The number of stereoisomers obtained by bromination of trans-2-butene is :
 (A) 1 (B) 2 (C) 3 (D) 4 **Ans. (A)**
30. Consider a titration of potassium dichromate solution with acidified Mohr's salt solution using diphenylamine as indicator. The number of moles of Mohr's salt required per mole of dichromate is :
 (A) 3 (B) 4 (C) 5 (D) 6 **Ans. (D)**
31. A solution of a metal ion when treated with KI gives a red precipitate which dissolves in excess KI to give a colourless solution. Moreover, the solution of metal ion on treatment with a solution of cobalt(II) thiocyanate gives rise to a deep blue crystalline precipitate. The metal ion is :
 (A) Pb^{2+} (B) Hg^{2+} (C) Cu^{2+} (D) Co^{2+} **Ans. (B)**

SECTION - II (Chemistry)

Assertion - Reason Type

This section contains 4 questions numbered 32 to 35. Each question contains **STATEMENT -1** (Assertion) and **STATEMENT-2** (Reason). Each question has 4 choices (A), (B), (C) and (D) out of which **ONLY ONE** is correct.


32. **STATEMENT -1** : Alkali metals dissolve in liquid ammonia to give blue solutions.
because
STATEMENT -2 : Alkali metals in liquid ammonia give solvated species of the type $[M(NH_3)_n]^+$ (M = alkali metals).
 (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 (C) Statement-1 is True, Statement-2 is False
 (D) Statement-1 is False, Statement-2 is True **Ans. (B)**
33. **STATEMENT-1** : Molecules that are not superimposable on their mirror images are chiral.
because
STATEMENT -2 : All chiral molecules have chiral centres.
 (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 (C) Statement-1 is True, Statement-2 is False
 (D) Statement-1 is False, Statement-2 is True **Ans. (C)**
34. **STATEMENT-1** : Glucose gives a reddish-brown precipitate with Fehling's solution.
because
STATEMENT -2 : Reaction of glucose with Fehling's solution gives CuO and gluconic acid.
 (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 (C) Statement-1 is True, Statement-2 is False
 (D) Statement-1 is False, Statement-2 is True **Ans. (C)**
35. **STATEMENT -1** : Band gap in germanium is small.
because
STATEMENT-2 : The energy spread of each germanium atomic energy level is infinitesimally small.
 (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 (C) Statement-1 is True, Statement-2 is False
 (D) Statement-1 is False, Statement-2 is True **Ans. (C)**

SECTION - III (Chemistry)


Linked Comprehension Type

This section contains 2 paragraphs C 36-38 and C 39-41. Based upon each paragraph, 3 multiple choice questions have to be answered. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct. **C 36-38 : Paragraph for Question Nos. 36 to 38**

Riemer-Tiemann reaction introduces an aldehyde group, on to the aromatic ring of phenol, ortho to the hydroxyl group. This reaction involves electrophilic aromatic substitution. This is a general method for the synthesis of substituted salicylaldehydes as depicted below .




36. Which one of the following reagents is used in the above reaction ?
 (A) aq. NaOH + CH₃Cl (B) aq. NaOH + CH₂Cl₂
 (C) aq. NaOH + CHCl₃ (D) aq. NaOH + CCl₄ Ans. (C)
37. The electrophile in this reaction is :
 (A) : CHCl (B) ⁺CHCl₂ (C) : CCl₂ (D) · CCl Ans. (C)
38. The structure of the intermediate I is :



Ans. (B)

C 39-41 : Paragraph for Question Nos. 39 to 41

Redox reactions play a pivotal role in chemistry and biology. The values of standard redox potential (E°) of two half-cell reactions decide which way the reaction is expected to proceed. A simple example is a Daniel cell in which zinc goes into solution and copper gets deposited. Given below are a set of half-cell reactions (acidic medium) along with their E° (V with respect to normal hydrogen electrode) values. Using this data obtain the correct explanations to Questions 36-38.



39. Among the following, identify the correct statement :
 (A) Chloride ion is oxidised by O₂ (B) Fe²⁺ is oxidised by iodine
 (C) Iodide ion is oxidised by chlorine (D) Mn²⁺ is oxidised by chlorine Ans. (C)
40. While Fe³⁺ is stable, Mn³⁺ is not stable in acid solution because :
 (A) O₂ oxidises Mn²⁺ to Mn³⁺ (B) O₂ oxidises both Mn²⁺ to Mn³⁺ and Fe²⁺ to Fe³⁺
 (C) Fe³⁺ oxidises H₂O to O₂ (D) Mn³⁺ oxidises H₂O to O₂ Ans. (D)
41. Sodium fusion extract, obtained from aniline, on treatment with iron (II) sulphate and H₂SO₄ in presence of air gives a Prussian blue precipitate. The blue colour is due to the formation of :
 (A) Fe₄[Fe(CN)₆]₃ (B) Fe₃[Fe(CN)₆]₂ (C) Fe₄[Fe(CN)₆]₂ (D) Fe₃[Fe(CN)₆]₃ Ans. (A)

SECTION - IV (Chemistry)

Matrix - Match Type

This section contains 3 questions. Each question contains statements given in two columns which have to be matched. Statements (A, B, C, D) in **Column-I** have to be matched with statements (p, q, r, s) in **Column-II**. The answers to these questions have to be appropriately bubbled as illustrated in the following example. If the correct matches are A -p, A -s, B -q, B -r, C -p, C -q and D -s, then the correctly bubbled 4 x 4 matrix should be as follows:

	p	q	r	s
A	<input checked="" type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s
B	<input type="radio"/> p	<input checked="" type="radio"/> q	<input checked="" type="radio"/> r	<input type="radio"/> s
C	<input checked="" type="radio"/> p	<input checked="" type="radio"/> q	<input type="radio"/> r	<input type="radio"/> s
D	<input type="radio"/> p	<input type="radio"/> q	<input type="radio"/> r	<input checked="" type="radio"/> s

42. Match the compounds/ions in **Column I** with their properties/reactions in **Column II**. Indicate your answer by darkening the appropriate bubbles of the 4 × 4 matrix given in the O R S

Column I

- (A) C₆H₅CHO
 (B) CH₃C ≡ CH
 (C) CN⁻
 (D) I⁻

Column II

- (p) gives precipitate with 2,4 dinitrophenylhydrazine
 (q) gives precipitate with AgNO₃
 (r) is anucleophile
 (s) is involved in cyanohydrin formation

Ans. A (p,q,s); B (q); C (q,r,s), D (q,r)

43. Match the crystal system/unit cells mentioned in Column I with their characteristic feature mentioned in Column II. Indicate your answer by darkening the appropriate bubbles of the 4×4 matrix given in the ORS.

Column I

- (A) simple cubic and face centred cubic
 (B) cubic and rhombohedral
 (C) cubic and tetragonal
 (D) hexagonal and monoclinic

Column II

- (p) have these cell parameters
 $a = b = c$ and $\alpha = \beta = \gamma$
 (q) are two crystal systems
 (r) have only two crystallographic angles of 90°
 (s) belong to same crystal system

Ans. A (p, s) ; B (p,q); C (q), D (q,r)

44. Match the reactions in column I with nature of the reactions/type of the products in Column II. Indicate your answer by darkening the appropriate bubbles of the 4×4 matrix given in the ORS.

Column I

- (A) $O_2^- \rightarrow O_2 + O_2^{2-}$
 (B) $CrO_4^{2-} + H^+ \rightarrow$
 (C) $MnO_4^- + NO_2^- + H^+ \rightarrow$
 (D) $NO_3^- + H_2SO_4 + Fe^{2+} \rightarrow$

Column II

- (p) redox reaction
 (q) one of the products has Trigonal planar structure
 (r) dimeric bridged tetrahedral metal ion
 (s) disproportionation

Ans. A (p, s); B (r); C (p,q), D (p)

PAPER (II)-2007
PART III
SECTION – I (Mathematics)
Straight Objective Type

This section contains 9 multiple choice questions numbered 45 to 53. Each question has 4 choices (A), (B), (C) and (D), out of which ONLY ONE is correct.

45. Let $f(x) = \frac{x}{(1+x^n)^{1/n}}$ for $n \geq 2$ and $g(x) = \underbrace{(\text{fofo} \dots \text{of})}_{\text{f occurs n times}}(x)$. Then $\int x^{n-2}g(x)dx$ equals-

- (A) $\frac{1}{n(n-1)}(1+nx^n)^{1-\frac{1}{n}} + K$ (B) $\frac{1}{n-1}(1+nx^n)^{1-\frac{1}{n}} + K$
 (C) $\frac{1}{n(n+1)}(1+nx^n)^{1+\frac{1}{n}} + K$ (D) $\frac{1}{n+1}(1+nx^n)^{1+\frac{1}{n}} + K$

Ans. (A)

46. Let \vec{a} , \vec{b} , \vec{c} be unit vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$. Which one of the following is correct ?

- (A) $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} = \vec{0}$
 (B) $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} \neq \vec{0}$
 (C) $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{a} \times \vec{c} \neq \vec{0}$
 (D) $\vec{a} \times \vec{b}, \vec{b} \times \vec{c}, \vec{c} \times \vec{a}$

are mutually perpendicular

Ans. (B)

47. The letters of the word **COCHIN** are permuted and all the permutations are arranged in an alphabetical order as in an English dictionary. The number of words that appear before the word **COCHIN** is

- (A) 360 (B) 192 (C) 96 (D) 48 **Ans. (C)**

48. $\frac{d^2x}{dy^2}$ equals

- (A) $\left(\frac{d^2y}{dx^2}\right)^{-1}$ (B) $-\left(\frac{d^2y}{dx^2}\right)^{-1}\left(\frac{dy}{dx}\right)^{-3}$
 (C) $\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-2}$ (D) $-\left(\frac{d^2y}{dx^2}\right)\left(\frac{dy}{dx}\right)^{-3}$ **Ans. (D)**

49. Let ABCD be a quadrilateral with area 18, with side AB parallel to the side CD and $AB = 2 CD$. Let AD be perpendicular to AB and CD. If a circle is drawn inside the quadrilateral ABCD touching all the sides, then its radius is

- (A) 3 (B) 2 (C) $\frac{3}{2}$ (D) 1 **Ans. (B)**

50. If $|z| = 1$ and $z \neq \pm 1$, then all the values of $\frac{z}{1-z^2}$ lie on
 (A) a line not passing through the origin
 (B) $|z| = \sqrt{2}$
 (C) the x-axis
 (D) the y-axis Ans. (D)

51. The differential equation $\frac{dy}{dx} = \frac{\sqrt{1-y^2}}{y}$ determines a family of circles with
 (A) variable radii and a fixed centre at (0, 1)
 (B) variable radii and a fixed centre at (0, -1)
 (C) fixed radius 1 and variable centres along the x-axis
 (D) fixed radius 1 and variable centres along the y-axis Ans. (C)

52. Let O (0, 0), P(3, 4), Q (6, 0) be the vertices of the triangle OPQ. The point R inside the triangle OPQ is such that the triangles OPR, PQR, OQR are of equal area. The coordinates of R are
 (A) $\left(\frac{4}{3}, 3\right)$ (B) $\left(3, \frac{2}{3}\right)$ (C) $\left(3, \frac{4}{3}\right)$ (D) $\left(\frac{4}{3}, \frac{2}{3}\right)$
Ans. (C)

53. Let E^c denote the complement of an event E. Let E, F, G be pairwise independent events with $P(G) > 0$ and $P(E \cap F \cap G) = 0$. Then $P(E^c \cap F^c | G)$ equals
 (A) $P(E^c) + P(F^c)$ (B) $P(E^c) - P(F^c)$ (C) $P(E^c) - P(F)$ (D) $P(E) - P(F^c)$
Ans. (C)

SECTION – II (Mathematics) Assertion–Reason Type

This section contains 4 questions numbered 54 to 57. Each question contains STATEMENT–1 (Assertion) and STATEMENT–2 (Reason). Each question has 4 choices (A), (B), (C) and (D) out of which ONLY ONE is correct.

54. Consider the planes $3x - 6y - 2z = 15$ and $2x + y - 2z = 5$
STATEMENT–1 : The parametric equations of the line of intersection of the given planes are $x = 3 + 14t$, $y = 1 + 2t$, $z = 15t$.
because
STATEMENT–2 : The vector $14\hat{i} + 2\hat{j} + 15\hat{k}$ is parallel to the line of intersection of given planes.
 (A) Statement–1 is True, Statement–2 is True; Statement–2 is a correct explanation for Statement–1.
 (B) Statement–1 is True, Statement–2 is True; Statement–2 is NOT a correct explanation for Statement–1
 (C) Statement–1 is True, Statement–2 is False
 (D) Statement–1 is False, Statement–2 is True

Ans. (D)

55. Let $f(x) = 2 + \cos x$ for all real x
STATEMENT-1 : For each real t , there exists a point c in $[t, t + \pi]$ such that $f'(c) = 0$.
because
STATEMENT-2 : $f(t) = f(t + 2\pi)$ for each real t .
 (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 (C) Statement-1 is True, Statement-2 is False
 (D) Statement-1 is False, Statement-2 is True

Ans. (B)

56. **STATEMENT-1** : The curve $y = \frac{-x^2}{2} + x + 1$ is symmetric with respect to the line $x = 1$
because
STATEMENT-2 : A parabola is symmetric about its axis.
 (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 (C) Statement-1 is True, Statement-2 is False
 (D) Statement-1 is False, Statement-2 is True

Ans. (A)

57. Lines $L_1 : y - x = 0$ and $L_2 : 2x + y = 0$ intersect the line $L_3 : y + 2 = 0$ at P and Q, respectively. The bisector of the acute angle between L_1 and L_2 intersects L_3 at R.

STATEMENT-1 : The ratio PR : RQ equals $2\sqrt{2} : \sqrt{5}$

because

STATEMENT-2 : In any triangle, bisector of an angle divides the triangle into two similar triangles.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 (C) Statement-1 is True, Statement-2 is False
 (D) Statement-1 is False, Statement-2 is True

Ans. (C)

SECTION – III

Linked Comprehension Type

This section contains 2 paragraphs M_{58-60} and M_{61-63} . Based upon each paragraph, 3 multiple choice questions have to be answered. Each question has 4 choices (A), (B), (C) and (D) out of which **ONLY ONE** is correct.

M_{58-60} : Paragraph for Question Nos. 58 to 60

If a continuous function f defined on the real line R , assumes positive and negative values in R then the equation $f(x) = 0$ has a root in R . For example, if it is known that a continuous function f on R is positive at some point and its minimum value is negative then the equation $f(x) = 0$ has a root in R .

Consider $f(x) = ke^x - x$ for all real x where k is a real constant.

58. The line $y = x$ meets $y = ke^x$ for $k \leq 0$ at
 (A) no point (B) one point
 (C) two points (D) more than two points

Ans. (B)

59. The positive value of k which $ke^x - x = 0$ has only one root is
 (A) $\frac{1}{e}$ (B) 1 (C) e (D) $\log_e 2$ **Ans. (A)**
60. For $k > 0$, the set of all values of k for which $ke^x - x = 0$ has two distinct roots is
 (A) $\left(0, \frac{1}{e}\right)$ (B) $\left(\frac{1}{e}, 1\right)$ (C) $\left(\frac{1}{e}, \infty\right)$ (D) $(0, 1)$ **Ans. (A)**

M 61–63 : Paragraph for Question Nos. 61 to 63

Let A_1, G_1, H_1 denote the arithmetic, geometric and harmonic means, respectively, of two distinct positive numbers. For $n \geq 2$, let A_{n-1} and H_{n-1} have arithmetic, geometric and harmonic means as A_n, G_n, H_n respectively.

61. Which one of the following statement is correct
 (A) $G_1 > G_2 > G_3 > \dots$ (B) $G_1 < G_2 < G_3 < \dots$
 (C) $G_1 = G_2 = G_3 = \dots$ (D) $G_1 < G_3 < G_5 < \dots$ and $G_2 > G_4 > G_6 > \dots$ **Ans. (C)**
62. Which one of the following statements is correct ?
 (A) $A_1 > A_2 > A_3 > \dots$
 (B) $A_1 < A_2 < A_3 < \dots$
 (C) $A_1 > A_3 > A_5 > \dots$ and $A_2 < A_4 < A_6 < \dots$
 (D) $A_1 < A_3 < A_5 < \dots$ and $A_2 > A_4 > A_6 > \dots$ **Ans. (A)**
63. Which one of the following statement is correct ?
 (A) $H_1 > H_2 > H_3 > \dots$
 (B) $H_1 < H_2 < H_3 < \dots$
 (C) $H_1 > H_3 > H_5 > \dots$ and $H_2 < H_4 < H_6 < \dots$
 (D) $H_1 < H_3 < H_5 < \dots$ and $H_2 > H_4 > H_6 > \dots$ **Ans. (B)**

SECTION – IV (Mathematics)
Matrix–Match Type

This section contains 3 questions. Each question contains statements given in two columns which have to be matched. Statements (A, B, C, D) in Column I have to be matched with statements (P, Q, R, S) in Column II. The answers to these questions have to be appropriately bubbled as illustrated in the following example.

If the correct matches are A–P, A–S, B–Q, B–R, C–P, C–Q and D–S, then the correctly bubbled 4×4 matrix should be as follows :

	P	Q	R	S
A	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>
B	<input type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>
C	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>
D	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>

64. Let (x, y) be such that $\sin^{-1}(ax) + \cos^{-1}(y) + \cos^{-1}(bxy) = \frac{\pi}{2}$
 Match the statement in Column I with statements in Column II and indicate your answer by darkening the appropriate bubbles in the 4 × 4 matrix given in the ORS.

Column I

- (A) If $a = 1$ and $b = 0$, then (x, y)
 (B) If $a = 1$ and $b = 1$, then (x, y)
 (C) If $a = 1$ and $b = 2$, then (x, y)
 (D) If $a = 2$ and $b = 2$, then (x, y)

Column II

- (P) lies on the circle $x^2 + y^2 = 1$
 (Q) lies on $(x^2 - 1)(y^2 - 1) = 0$
 (R) lies on $y = x$
 (S) lies on $(4x^2 - 1)(y^2 - 1) = 0$

Ans. A- P; B - Q, C- Q, D- S

65. Match the statements in Column I with the properties in Column II and indicate your answer by darkening the appropriate bubbles in the 4x4 matrix given the ORS.

Column I

- (A) Two intersecting circles
 (B) Two mutually external circles
 (C) Two circles, one strictly inside the other
 (D) Two branches of a hyperbola

Column II

- (P) have a common tangent
 (Q) have a common normal
 (R) do not have a common tangent
 (S) do not have a common Normal

Ans. A- P; B - P; C- Q,R D; Q, R

66. Let $f(x) = \frac{x^2 - 6x + 5}{x^2 - 5x + 6}$

- Match the expressions/statements in Column I with expression/statements in Column II and indicate your answer by darkening the appropriate bubbles in the 4x4 matrix given in the ORS.

Column I

- (A) If $-1 < x < 1$, then $f(x)$ satisfies
 (B) If $1 < x < 2$, then $f(x)$ satisfies
 (C) If $3 < x < 5$, then $f(x)$ satisfies
 (D) If $x > 5$, then $f(x)$ satisfies

Column II

- (P) $0 < f(x) < 1$
 (Q) $f(x) < 0$
 (R) $f(x) > 0$
 (S) $f(x) < 1$

Ans. A- P, R, S; B - Q, S; C- Q, S; D- P, R, S