## LIFE SCIENCES PAPER I(PART 'B')

- 41. During protein synthesis, L-amino acid binds to t-RNA through
  - α-amino group.
  - 2. hydrophobic side chain.
  - α-carboxyl group.
  - carboxyl group of the side chain.
- 42. The peptide bond is planar
  - due to restriction caused by rotation around c<sup>α</sup>-N bond
  - due to restriction around c<sup>α</sup>-c' bond
  - due to delocalization of the lone pair of electrons of the privogen onto carbonyl oxygen
  - 4. because amide protons and carbonyl oxygen are wow wed in hydrogen bonding.
- 43. Hydrogen bond length will NOT be
  - independent of the nature of don and acceptor atoms.
  - dependent on donor and acceptor a soms.
  - dependent on the solvent is which the molecule is dissolved.
  - dependent on the other an ambonded with the donor and acceptor atom.
- 44. Why a DNA duplex melts at propecific temperature (Tm) on heating?
  - Loss of base stacking energy
  - The double figlix is intrinsically unstable
  - The single helix is more stable as compared to the double helix
  - 4. The Road double helix is a co-operative structure stabilized by hydrogen bonds and base pairing
- 45. Lipid bil ayers can be formed by phospholipids which have variable head groups and fatty acyl chains. The fluidity of the membrane will depend on
  - only the nature of head groups.
  - only the length of the fatty acid chains irrespective of the extent of unsaturation.
  - only unsaturation irrespective of the length of the fatty acid chains.
  - length and degree of unsaturation of fatty acid chains.

46.	Which express	one of the following RNA molecules is involved in regulation of gene sion?
	1. 2. 3. 4.	miRNA rRNA 5S RNA tRNA
47.	In which	ch organelle is NADP <sup>+</sup> the final electron acceptor?
	1. 2. 3. 4.	Only chloroplast Only mitochondrion Both chloroplast and mitochondrion Lysosome
48.	When	the K'q = 1, ΔG° is equal to
	1. 2. 3. 4.	-1. 0. +1. 10.
49.	Which  1. 2. 3. 4.	one of the following human pathogen is a flagellated protozoan?  Trypanosoma Plasmodium Paramecium Entamenha
50.	Which  1. 2. 3. 4.	of the following mammalian cells usually does NOT divide in adult life?  Noithelial cells in lung  Nerve cells in brain  Liver cells  Osteoblast cells
51.	What h	appens to the Cdk-cyclin A complex at metaphase?
	1. 2. 3. 4.	Both cyclin A and Cdk remain undegraded Only Cdk is degraded Only cyclin A is degraded Both cyclin A and Cdk are degraded

52.	vari ou:	rerage human genome has approximately 3×10 <sup>9</sup> base pairs coding for sproteins. If an "average" protein contains 400 amino acids, what is the number of proteins that can be encoded by the human genome?
	1. 2. 3. 4.	$2.5 \times 10^{6}$ . $2.5 \times 10^{7}$ . $3.0 \times 10^{6}$ . $3.5 \times 10^{7}$ .
53.	In eukaryotes, the interaction of enhancer and promoter elements is broug closer by	
	1. 2. 3. 4.	zinc finger. DNA looping. helix turn helix. palindrome.
54.	contair	of the following cytoskeleton elements guides the movement of vesicles ning cell wall precursors from their site of familiation in Golgi to the site of all formation in a growing pollen tube?
	1. 2. 3. 4.	Myosin Actin Kinesin Dynein
55.	bilayer	ores are small hydrophobic molecules that can partition into the lipid and increase their permeability to specific inorganic ions. Which of the ing is a channel forwing ionophore?
	1. 2. 3. 4.	Valinomych. Actimoraych. Gramboldin A. Toin.
56.		GTPases regulate intracellular transport in mammalian cells through fusion?
	1. 2. 3. 4.	Rab Ran Ras Rho

57.	The region where RNA polymerase binds to promoter in prokaryotes is called
	<ol> <li>Pribnow box.</li> <li>Hogness box.</li> <li>Homeo box.</li> <li>Shine-Dalgarno box.</li> </ol>
58.	Which of the following features highlights the difference between Z-DNA and BDNA?
	<ol> <li>Double helical nature</li> <li>Orientation of phosphate backbone</li> <li>Pairing of G - C</li> <li>Antiparallel nature of two polynucleotide strands of double helix</li> </ol>
59.	Ribosomal subunits are assembled in
	1. cytoplasm. 2. nucleolus. 3. nucleus. 4. endoplasmic reticulum.
60.	The absence of sigma factor in RNA polymerase
	1. affects elongation only. 2. blocks initiation only. 3. affects both initiation and elongation. 4. does not affect transcription.
61.	Which of the following features is not required in the initiation step of protein synthesis?  1. Unino acid activation 2. Unding of mRNA to the ribosomes 3. Transfer of activated amino acid to tRNA 4. Joining together of two amino acids by peptide bond formation
62.	By which of the following mechanisms does cycloheximide inhibit protein synthesis?
	<ol> <li>Blocking the peptidyl transferase of 80S euokaryotic ribosomes</li> <li>Blocking the peptidyl transferase of 70S prokaryotic ribosomes</li> <li>By binding to DNA dependent RNA polymerase</li> </ol>

By binding to sigma factor


63.	Which of the following is a natural inducer of the lac operon in E. coli?
	<ol> <li>Lactose</li> <li>Galactose</li> <li>Allolactose</li> <li>IPTG</li> </ol>
64.	Antitermination of RNA synthesis is a major mechanism of regulation in
	<ol> <li>lytic phase of λ phage.</li> <li>lysogenic phase of λ phage.</li> <li>lac operon.</li> <li>trp operon.</li> </ol>
65.	Influenza virus binds to its host cells through which one of the two wing carbohydrate moiety.
	<ol> <li>N-acetyl glucosamine</li> <li>N-acetyl neuraminic acid</li> <li>Fucose</li> <li>Mellibiose</li> </ol>
66.	In mammals, G protein coupled receiptors (GPCR) play a major role in mediating effects of various hormones NOT through
	<ol> <li>activation of protein linase A.</li> <li>activation of aden wats cyclase.</li> <li>inactivation of alenylate cyclase.</li> <li>activation of tyrosine kinase activity.</li> </ol>
67.	Receptors for neurotransmitters are located on the
	<ol> <li>Surface.</li> <li>Surface.</li> <li>Surface.</li> <li>endosome.</li> <li>Golgi apparatus.</li> </ol>
68.	One of the major transmembrane proteins in a "tight junction" is
	<ol> <li>lectin.</li> <li>claudin.</li> <li>adherin.</li> <li>integrin.</li> </ol>

- 69. Retroviruses are well known as cancer causing agents because it
  - causes mutations in host genes involved in growth.
  - integrates their provinal DNA next to protooncogenes.
  - generates defective viruses lacking some of the viral genes.
  - integrates their T antigens into the host genome.
- 70. Select the correct statement:
  - In vertebrate development, immune and nervous systems are the prevalent sites for apoptosis.
  - Apoptosis is triggered by growth stimulus.
  - The tum or suppressor factor p53 inhibits apoptosis.
  - The soluble form of tumor necrosis factor cannot induce appropriosis.
- 71. ELISA assay
  - 1. uses complement mediated cell lysis.
  - 2. uses a radiolabeled second antibody.
  - 3. involves addition of substrate which is converted to coloured end product.
  - requires specialized red blood cells,
- 72. In mosaic development, the prospective patency of cells
  - 1. equals its prospective fata
  - is greater than prospective hate
  - is less than prospective fate
  - 4. and fate are unrallate,d
- 73. Experiments with sea unchin demonstrated species specific sperm-egg recognition through the protein
  - bidelil
  - Willin
  - 3. Stivin
  - hyalin.
- 74. Exposing a regenerating limb to which of the following chemicals results in the blastem a proximalization?
  - Ascorbic acid
  - Thyroxine
  - Retinoic acid
  - Glutamic acid

- 75. Temperature-dependent sex determination is observed in
  - Drosophila.
  - amphibians.
  - reptiles.
  - sea urchins.
- 76. Some plants require vernalization (prolonged cold treatment) for transition to flowering. For floral induction, vernalization signal is perceived primarily by
  - young leaves subtending the apical meristem.
  - 2. mature leaves near the root-shoot junction.
  - all vegetative parts.
  - shoot apical meristem.
- 77. Which phase of embryogenesis in plants is characterized by the initiation of deposition of storage reserves?
  - Globular stage
  - Heart stage
  - Torpedo stage
  - Cell enlargement stage
- 78. Which of the following statements is NOT true in relation to growth of animals?
  - 1. When all body parts grow the same rate, it is called isometric growth.
  - 2. When different body parts grow at different rates, it is called allometric growth.
  - Two-fold change in weight will cause a 1.26-fold expansion in length if growth is allemetric.
  - 4. Isometric & with cannot create dramatic changes in the structure of organisms.
- 79. Abnormative, during development caused by exogenous agents are called disruptive, and the agents are specifically called
  - morphogen.
  - teratogen.
  - allergen.
  - mutagen.

- 80. At which of the following steps does application of dichlorophenyl-dimethyl urea (DCMU) inhibit electron flow during photosynthesis?
  - P680\* → Pheophytin
  - 2.  $Q_A \rightarrow Q_B$
  - 3.  $Q_B \rightarrow Cytb_{e}f complex$
  - Cytbef complex → plastocyanin
- 81. A major functional difference between the succinyl CoA-synthetase of plant and animal cell mitochondria is that it
  - does not produce ATP in plant cell.
  - 2. does not produce GTP in plant cell.
  - produces ATP in plants and GTP in animals.
  - produces GTP in plants and ATP in animals.
- 82. A common symptom of molybdenum deficiency in plants is the accumulation of nitrate in the cytosol, which results from
  - reduced nitrite reductase activity.
  - reduced nitrate reductase activity.
  - reduced transport of nitrate into the vacuoles
  - reduced transport of nitrate into the chloroplasts.
- 83. The Rht mutations in wheat that we've pivotal for 'Green Revolution' cause reduction in plant height due to impairment in
  - gibberellic acid bij synthesis pathway.
  - gibberellic acid Magnaling pathway.
  - auxin biosymthetic pathway.
  - auxin response pathway.
- 84. Which of the full owing phytochrome controlled responses displays red/far-red reversibility.
  - 1. Pery low-fluence responses
  - 2. Low-fluence responses
  - High-irradiance responses
  - Very high-irradiance responses
- 85. At permanent wilting point, plants cannot regain turgor pressure even if transpiration stops because
  - water potential of soil (ψ<sub>w</sub>) is less than or equal to osmotic potential (ψ<sub>S</sub>)
    of the plant.
  - ψ<sub>w</sub> is higher than ψ<sub>S</sub>.
  - Ψ<sub>w</sub> and Ψ<sub>S</sub> are unaltered.
  - Ψs remains unaltered.

- 86. Alkaloid production in plants is regulated by a change in the endogenous pool of
  - gibberellins.
  - jasmonates.
  - brassinosteroi ds.
  - abscisic acid.
- 87. Flooding and submergence of plants leads to anoxic conditions. Usually, flood-tolerant species alter their developmental programme, resulting in
  - increased activity at the apical meristem.
  - growth retardation of foliage.
  - elongation growth of their stems.
  - increased proliferation of root system.
- 88. An organism with the genotype AABbCcDD is selfed. The four genes are independently assorting. What proportion of the progent will show the genotype AAbbccDD?
  - 1. 1/4
  - 2. 1/16
  - 3. 1/64
  - 4. 1/256
- 89. Two fruitflies with mutant eye color were crossed. All progenies obtained from this cross had wild type eye color with as
  - the mutations are Melac.
  - one mutation is Commant over the other.
  - the mutations are co-dominant.
  - the mutations are in two different genes.
- 90. The following pedigree represents the inheritance of a rare disorder.



Based on the above pedigree, what is the most likely mode of inheritance?

- Autosomal dominant
- X-linked recessive
- X-linked dominant
- Y-linked dominant

- 91. Which of the following pair of relatives will have the highest genetic correlation?
  - First double cousins
  - 2. Half siblings
  - Brothers
  - Brother-sister
- 92. a<sub>1</sub>, a<sub>2</sub>, a<sub>3</sub> are three alleles of a gene in *Neurospora*. Crosses between different a mutants gave the following results

$a_1 \times a_2$	$a_1 \times a_3$	$a_2 \times a_3$
$\downarrow$	$\downarrow$	$\downarrow$
a <sub>l</sub> +	$a_1+$	a <sub>2</sub> +
++	$a_1+$	$a_2$ +
$+ a_{2}$	++	++
+ a <sub>2</sub>	+a <sub>3</sub>	+ a <sub>3</sub>

The process which best explains this result is

- gene conversion.
- forward mutation.
- non-homologous recombination.
- 4. in complete dominance.
- 93. You are provided with an E. coli strain are otrophic for the amino acids Trp, Leu and Val. To select for double reversants in Trp and Leu, which amino acid(s) would you include in the growth medium?
  - 1. Trp
  - Leu
  - Val
  - 4. Trp, Leu, Way
- 94. UAG and U. A are both nonsense codons. What kind of single point mutation would car we resion of UAG to a meaningful codon?
  - 1. Washsition
  - 2. Sansversion
  - Frameshift
  - Inversion
- 95. E. coli cells were simultaneously infected by two rII bacteriophage mutants. From the progeny obtained after lysis of the E. coli cells it was observed that some of the bacteriophages showed a wild type phenotype. These were obtained at extremely low frequency. This is due to
  - complementation of the two mutations.
  - recombination between the two mutant chromosomes.
  - transposition of the mutation.
  - incomplete penetrance.

96.		of the following hormones stimulates the reabsorption of $\mathrm{Na}^+$ and the on of $\mathrm{K}^+$ in the kidney?
	1. 2. 3. 4.	Vasopressin Thyroxine Prolactin Aldosterone
97.	The correct sequence in vertebrate embryonic development is	
98.	1. 2. 3. 4. A perso	gastrocoel - blastocoel - notochord - neural crest.  blastocoel - gastrocoel - neural crest - notochord.  gastrocoel - blastocoel - neural crest - notochord.  blastocoel - neural crest - gastrocoel - notochord.  on has a vision problem caused by the image of an object at infinity getting d in front of retina. The error can be corrected by the use of
	1. 2. 3. 4.	biconvex lens. cylindrical lens. plano convex lens. biconcave lens.
99.	During  1. 2. 3. 4.	electrical stimulation-induced depelarization of neuron, voltage-gated  Na <sup>+</sup> channels will close  K <sup>+</sup> channels will open.  Na <sup>+</sup> channels will open.
100.	1. 2.	mary function of diaphragm is to  continuous diaphragm is to  reputable respiration.  port the heart.  the port the rib cage dilated.
101.	In neph 1. 2. 3. 4.	reduce urine pH. reduce water loss. release urea. release uric acid.


102.	During inspiration, the air that we breathe moves through different regions of the associated organs in the sequence of
	<ol> <li>larynx &gt; nasopharynx &gt; trachea &gt; glottis.</li> <li>nasopharynx &gt; glottis &gt; larynx &gt; trachea.</li> <li>glottis &gt; nasopharynx &gt; larynx &gt; trachea.</li> <li>larynx &gt; glottis &gt; nasopharynx &gt; trachea.</li> </ol>
103.	Knee jerk reaction is an example of what type of reflex?
104.	<ol> <li>Monosynaptic</li> <li>Multisynaptic</li> <li>Conditioned</li> <li>Conscious</li> <li>Which of the following processes is a major problem in interpreting molecular phylogeny?</li> </ol>
	1. Horizontal transfer of genes 2. Gene duplication 3. Synonymous mutations 4. Non-synonymous mutations
105.	In a tissue, a cell that markedly differs in form, size and content from other cells of the same tissue is called  1. intermediary cell. 2. isotropic cell. 3. idioblast cell. 4. myrosin cell
106.	Among the extant reptiles which group is phylogenetically closely related to Aves?  1. Curtles 2. Lizards 3. Snakes 4. Crocodiles
107.	The group of organisms that is now separated from the other groups of fungi based on their motile spores and cellulose-rich cell wall is

Myxomycetes. Zygomycetes. Deuteromycetes.

Oomycetes.

1. 2. 3. 4.

- 108. Some floristic elements common to both India and China are in the genus
  - Ginkgo.
  - Rhododendron.
  - Poeciloneuron.
  - 4. Erinocarpus.
- 109. The difference between Indian and African wild herbivore fauna is that there are no
  - 1. antelopes in India.
  - deer in Africa.
  - odd-toed animals in India.
  - even-toed hoofed animals in Africa.
- 110. Which of the following bird species is endangered?
  - Hill myna
  - Great Indian bustard
  - Crow-pheasant
  - Grey hombill
- 111. In spite of the prevalence of herbivory the earth continues to be largely green because
  - 1. the number of herbivore species is low.
  - herbivores are very in efficient feeders.
  - herbivore number mark kept low by their predators.
  - herbivory promote plant growth.
- 112. Which of the following curves represents the general relationship between body size (S) and influence rate of population growth (r)?



- 113. Which of the following reproductive strategies is characteristic of marine invertebrates?
  - Long generation time, small clutch size
  - Short generation time, small clutch size
  - Long generation time, large clutch size
  - 4. Short generation time, large clutch size
- 114. Bergmann's Rule refers to a general tendency of mammals to be
  - 1. larger in size in colder areas of their distribution.
  - smaller in size in areas of their distribution.
  - darker-pigmented in warmer areas of their distribution.
  - lighter-pigmented in warmer areas of their distribution.
- 115. When removal of a species from an ecosystem affects persistence of many other species and the impact of that species removal is disproportionale to its abundance, the species is known as
  - 1 indicator species.
  - keystone species.
  - flagship species.
  - umbrella species.
- 116. Biomass turnover time is the ratio between biomass and productivity of an ecosystem. Which of the following forests should have highest biomass turnover time?
  - Tropical dry forestess
  - 2. Tropical wet foresh
  - 3. Temperate decidinals forests
  - Boreal foresh;
- 117. Conversion of Nitrite to nitrate in soil is done by the bacteria of genus
  - Waltobacter.
  - Strosomonas.
  - Mitrobacter.
  - Pseudomonas.
- 118. In a population with two alleles 'a' and 'b' of a genotype in a ratio of 0.2 and 0.8 in Hardy-Weinberg equilibrium, how many individuals in a sample of 300 can be expected to be homozygous for allele 'a'?
  - 1. 192
  - 2. 12
  - 3. 64
  - 4. 96

119. Defective alleles are eliminated rapidly from a population if they are 1. recessive. 2. dominant. 3. codominant. in multiple copies. 120. The correct expression of Hamilton rule for the evaluation of altruism is [C = the cost of a behavioral act to the actor, b = the benefit of that act to a beneficiary, and r =the genetic relatedness between the actor and the beneficiary] 1. c < b.r 2. c < b 3. c-r < b 4.  $r \le b \cdot c$ The evolutionary basis of sexual dimorphism is 121. 1. differential investment in offspring. 2. difference in aggression. 3. difference in sex chromosomes. 4. difference in autosomes. The evolutionary appearance of the first mammals was 122. after the extinction of a so aurs and before the appearance of birds. 1. 2. before the extinction chimical constants and after the appearance of birds. before the exting two of dinosaurs and before the appearance of birds. 3. after the extinction of dinosaurs and after the appearance of birds. 4. Both the Luria Respirack experiment and the Lederberg and Lederberg experiment 123. demonstrate 🖔 șelection mutations. st-selection mutations. 3. directed mutations. 4. adaptive mutations. 124. Vampire bats regurgitate food in order to feed a starving member of their group. This is an example of 1. group selection. 2. reciprocal altruism. 3. selfish behavior. 4. K-selection.

125.	The first appearance of amphibians on earth was during the period
	1. Silurian.
	2. Carboniferous.
	3. Triassic.
	4. Jurassic.
126.	In a transgenic mouse experiment a founder male produces 100 pups and only 20
	of these are transgenic. This result leads to the conclusion that the
	founder transgenic animal is chimaeric.
	2. founder animal is mosaic.
	3. transgene is integrated on Y-chromosome.
	4. transgene is integrated on X-chromosome.
127.	An inbred mouse is cloned using another mouse strain as an esg wonor. The
	genetic relationship between the original inbred mouse and its coone will be
	1. 100%.
	2. 99–100%.
	3. 95–99%.
	4/ 90–95%.
100	Which and of the control of the cont
128.	Which of the following elements can be sed to immobilize a reduced protein?
	1. Calcium
	2. Potassium
	3. Gold
	4. Sodium
129.	D-amino acids 🐞 🔻
	1. cannot be produced by fermentation.
	2. cart produced by E. coli.
	3. Par be produced by yeast.
	4. In be produced by microbes provided with chiral precursors of D-amino
	aci ds.
130.	Which of the following viruses is used for biocontrol of insect pests of plants?
	···
	1. Cauliflower mosaic virus
	2. Cucumber mosaic virus
	<ol> <li>Rice tungro virus</li> </ol>
	4. Nuclear polyhedrosis virus

- 131. What ratio do you expect for codominant loci in F2 populations?
  1. 1:1
  2. 3:1
  3. 1:2:1
  4. 9:3:3:1
  132. The copy number of a transgene in plants can be deciphered by
  1. Southern blotting.
  - northern blotting.
  - 3. south western blotting.
  - far western blotting.
- 133. It is hypothesized that the mean (μ) longevity of a Drosophila sit sin is 18 days, with a variance (σ) of 3 days. What values of longevity in a campled population will lead to rejection of the null hypothesis at 95% confidence level?
  - 1. Only values less than 15.
  - Values less than 15 and more than 18
  - Only values more than 21
  - 4. Values less than 12 and more that? 2
- 134. Normalized Differential Vegetation Index (NDVI) in remote sensing refers to the following spectral band derivative:
  - 1. Near IR Red
  - Red/Near Red
  - (Near IR − Red) Medear IR + Red)
  - 4. (Near IR wkked)₩ Red
- 135. Function of conformator in a spectrophotometer is
  - 1. sing a straight beam of light.
  - Wording a light beam into its component wavelengths.
  - selecting a desired wavelength.
  - creating a light source.
- 136. Which of the following methods is the most appropriate for estimating the population density of burrowing animals?
  - Quadrat sampling
  - Line transect sampling
  - Tag-recapture method
  - Nearest neighbour distance method

137.	A radioactive sample was counted in a scintillation counter and a value of 194930 cpm was obtained. The counting efficiency was found to be 95.7%. What is the actual amount of radioactivity present in the sample?
	1. 181171 dpm 2. 190411 dpm 3. 203600 dpm 4. 211326 dpm
138.	Which one of the following treatments does NOT enhance the response of a film to radioisotopes in autoradiography?
	<ol> <li>Staining the gel with Coomassie blue before drying</li> <li>Use of intensifying screens</li> </ol>

What is the length of oligonucleotide required to give consistent specific

A protein is poorly expressed in a seased tissue. To determine whether the defect is at the level of translation or translation, which of the following

Exposure at low temperature

hybridization signal in a microarray?

blotting methods would you use?

Southern, and northern Northern and western

Southern 💩

Preflashing the film with a light flash

3.

4.

1. 2.

3.

5

7

9 20

139.

140.