

Seat No.: _____

Enrolment No. _____

(EM-8)

GUJARAT TECHNOLOGICAL UNIVERSITY

B.E. all Sem-I Examination December 08/January 09

Maths-I (110008)**DATE: 22-12-2008, Monday TIME: 12.00 to 3.00 p.m. MAX. MARKS: 70****Instructions:**

- 1. Attempt all questions.**
- 2. Make suitable assumptions wherever necessary.**
- 3. Figures to the right indicate full marks.**

Q.1(a) (i) Let the function f be defined by**02**

$$f(x) = \begin{cases} |x|/x & ; x > -2, x \neq 0 \\ x+1 & ; x < -2 \\ 3 & ; x = -2 \end{cases}$$

Evaluate each of the following limits if it exists.

$$\lim_{x \rightarrow -2} f(x), \lim_{x \rightarrow 0} f(x), \lim_{x \rightarrow 2/5} f(x).$$

(ii) Explain various types of discontinuities with an example.**02****(b) (i) Using L'Hospital rule, evaluate $\lim_{x \rightarrow 0} \frac{1}{x}(1-x \cot x)$.****02****(ii) State Sandwich theorem on sequences and using it show that, if $x \in \mathbb{R}$ with $|x| < 1$, then $x^n \rightarrow 0$ as $n \rightarrow \infty$.****02****(c) (i) Using mean value theorem, show that $3 + \frac{1}{28} < \sqrt[3]{28} < 3 + \frac{1}{27}$.****03****(ii) Prove that $\tan^{-1} \left[\frac{x(3-4x^2)}{\sqrt{1-x^2}(1-4x^2)} \right] = 3 \left(x + \frac{x^3}{6} + \frac{3x^5}{40} + \dots \right)$.****03****Q.2(a) Attempt any two of the following.****04****(i) For $f(x) = x + 2$, $x \in [0,5]$, find $U(f,P)$ and $L(f,P)$ for $P = \{0,1,2,3,4,5\}$.****(ii) State Fundamental theorem of Integral calculus and give one example on which it is not applicable.****(iii) Evaluate the improper integral $\int_0^\infty \frac{1}{x^2} dx$.****(b) Evaluate $\int_3^7 4\sqrt{(x-3)(7-x)} dx$ using the concept of Gamma and Beta function.****04****(c) (i) Test the convergence and divergence of the following series (any two).****04**

$$[A] \sum_{n=1}^{\infty} \tan^{-1} \left(\frac{1}{n^2 + n + 1} \right) [B] 5 - \frac{10}{3} + \frac{20}{9} - \frac{40}{27} + \dots [C] \sum_{n=1}^{\infty} \frac{2n^2 + 3n}{5 + n^5} .$$

(ii) Evaluate $\iint\limits_{0,0}^{1,x} (x^2 + y^2) dA$, where dA indicates small area in xy -plane. **02**

Q.3(a) Suppose that $u = f(x, y, z)$ and $x = g_1(t), y = g_2(t), z = g_3(t)$.

Then write the chain rule for derivative of u w.r.t. t .

Express $\partial w/\partial r$ and $\partial w/\partial s$ in terms of r and s if $w = x + 2y + z^2$,
 $x = r/s, y = r^2 + \log(s), z = 2r$. **05**

(b) (i) Evaluate $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^4 + y^2}$. **02**

(ii) Find the absolute maximum and minimum values of
 $f(x, y) = 2 + 2x + 2y - x^2 - y^2$ on the triangular plate in the first quadrant bounded by the lines $x = 0, y = 0, y = 9 - x$. **03**

(c) Find the length of the arc of the curve $x = a \cos\theta$ and $y = a \sin\theta ; a > 0$. **04**

OR

Q.3(a) State Euler's theorem on homogeneous function. Using it show that **05**

$$(i) x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \sin 2u \quad (ii) x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = 2 \sin u \cos 3u$$

if $u = \tan^{-1}(x^2 + 2y^2)$.

Explain whether we can apply Euler's theorem for the function

$$u = f(x, y) = \frac{x^2 + y^2 + 1}{x + y}.$$

(b) (i) If $u = f(x, y)$ then what geometrically $\partial u/\partial x$ indicates? **01**

(ii) If $x = \sqrt{vw}, y = \sqrt{wu}, z = \sqrt{uv}$ and $u = r \sin\theta \cos\phi$,

$v = r \sin\theta \sin\phi, w = r \cos\theta$. Calculate $\frac{\partial(x, y, z)}{\partial(r, \theta, \phi)}$. **02**

(iii) The pressure P at any point (x, y, z) in space is $P = 400xyz^2$.

Find the highest pressure on the surface of the unit sphere

$$x^2 + y^2 + z^2 = 1.$$

(c) Find the volume generated by revolving the parabola $y^2 = 4ax ; a > 0$ about the latus rectum. **04**

Q.4(a) Evaluate $\iint\limits_R (x + y) dy dx$; R is the region bounded by $x = 0, x = 2$,

$y = x, y = x + 2$. **05**

(b) Find the volume of the solid that lies under the plane $3x + 2y + z = 12$ and above the rectangle $R = \{(x, y) \mid 0 \leq x \leq 1, -2 \leq y \leq 3\}$. **05**

(c) For the vector field $\vec{A} = k \vec{i}$ and $\vec{A} = k \vec{r}$. Find $\nabla \bullet \vec{A}$ and $\nabla \times \vec{A}$.

Draw the sketch in each case. **04**

OR

Q.4 (a) Change the order of integration and evaluate $\int\limits_0^{a/\sqrt{2}} \int\limits_x^{\sqrt{a^2 - x^2}} y^2 dA$. **05**

(b) Find the area common to $r = a$ and $r = 2a\cos\theta$; $a > 0$. 05

(c) Find the scalar potential function f for $\vec{A} = y^2\vec{i} + 2xy\vec{j} - z^2\vec{k}$. 04

Q.5(a) Evaluate $\oint_C -\frac{y}{x^2+y^2}dx + \frac{x}{x^2+y^2}dy$, where $C = C_1 \cup C_2$ with
 $C_1 : x^2 + y^2 = 1$ and $C_2 : x = \pm 2, y = \pm 2$. 05

(b) Evaluate $\iint_S 6xydS$, where S is the portion of the plane $x + y + z = 1$
that lies in front of the yz -plane. 05
(c) Trace the curve $y = x + (1/x)$. 04

OR

Q.5 (a) Find the surface area of a sphere of radius a . 05

(b) Evaluate $\int_C \vec{F} \bullet d\vec{r}$, where $\vec{F}(x, y, z) = -y^2\vec{i} + x\vec{j} + z^2\vec{k}$ and C is the
curve of intersection of the plane $y + z = 2$ and the cylinder $x^2 + y^2 = 1$. 05
(c) Trace the curve $9a y^2 = x(x - 3a)^2$, $a > 0$. 04
