



**First Semester B.E. Degree Examination, December 2011**  
**Engineering Mathematics - I**

Time: 3 hrs.

Max. Marks: 100

- Note: 1. Answer any **FIVE** full questions, choosing at least two from each part.  
 2. Answer all objective type questions only on OMR sheet page 5 of the answer booklet.  
 3. Answer to objective type questions on sheets other than OMR will not be valued.

**PART - A**

- 1 a. Choose your answers for the following : (04 Marks)

i) If  $y = \frac{x}{x-1}$ , then  $y_n$  is

A)  $\frac{(-1)^{n-1} n!}{(x-1)^{n+1}}$       B)  $\frac{(-1)^n n!}{(x-1)^{n+1}}$       C)  $\frac{(-1)^n (n+1)!}{(x-1)^{n+1}}$       D)  $\frac{(-1)^n n!}{(x-1)^n}$

ii) If  $y = \log(ax+b)$ , then  $y_n$  is

A)  $\frac{(-1)^n n! a^n}{(ax+b)^n}$       B)  $\frac{(-1)^{n-1} n! a^n}{(ax+b)^{n+1}}$       C)  $\frac{(-1)^{n-1} (n-1)! a^n}{(ax+b)^n}$       D)  $\frac{(-1)^n (n-1)! a^n}{(ax+b)^{n+1}}$

iii) If  $f(x) = \sin x$ ,  $x \in (0, \pi)$ , then by Rolle's theorem the value of 'x', where the Tangent is parallel to x-axis.

A) 0      B)  $\frac{\pi}{2}$       C)  $\frac{\pi}{3}$       D)  $\frac{\pi}{4}$

iv) Expansion of  $\log(1+x)$  in powers of x is

A)  $x + \frac{x^2}{2} + \frac{x^3}{3} + \dots$       B)  $x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$   
 C)  $1 - \frac{x}{1!} + \frac{x^2}{2!} - \frac{x^3}{3!} + \dots$       D)  $\frac{x}{1!} - \frac{x^2}{2!} + \frac{x^3}{3!} - \frac{x^4}{4!} + \dots$

a. If  $x = \tan(\log y)$ , show that  $(1+x^2)y_{n+1} + (2nx-1)y_n + n(n-1)y_{n-1} = 0$ . (04 Marks)

b. State and prove Cauchy's mean value theorem. (06 Marks)

c. Expand  $f(x) = \sin(e^x - 1)$  in power's of 'x' upto the terms containing  $x^4$ . (06 Marks)

- 2 a. Choose your answers for the following : (04 Marks)

i) The indeterminate form of  $\lim_{x \rightarrow 1} \left( \frac{x}{x-1} - \frac{(x-1)}{\log x} \right)$  is

A)  $\infty - \infty$       B)  $\frac{0}{0}$       C)  $\frac{\infty}{\infty}$       D) None of these

ii) The angle between the radius vector and the tangent to the curve  $r = k e^{\theta \cot \alpha}$ , where K and  $\alpha$  are constants, is :

A) K      B)  $\theta$       C)  $\alpha$       D) O

iii) The Pedal equation of the curve  $r = a\theta$  is.

A)  $p^2 = ar$       B)  $\frac{1}{p^2} = \frac{a}{r^2}$       C)  $\frac{1}{p^2} = \frac{1}{r^2} + a^2$       D)  $\frac{1}{p^2} = \frac{1}{r^2} + \frac{a^2}{r^4}$

iv) The radius of curvature at any point 't' on the curve defined by  $x = f(t)$ ,  $y = \phi(t)$  is given by

A)  $\frac{[(x')^2 + (y')^2]^{\frac{3}{2}}}{x'y'' - y'x''}$       B)  $\frac{x'y'' - y'x''}{[(x')^2 + (y')^2]^{\frac{3}{2}}}$       C)  $\frac{(x')^2 + (y')^2}{(x'y'' - y'x'')^{\frac{3}{2}}}$       D)  $\frac{(x'y'' - y'x'')^{\frac{3}{2}}}{(x')^2 + (y')^2}$

- b. Find the angle of intersection between the curves  $r^n \cos(n\theta) = a^n$  and  $r^n \sin(n\theta) = b^n$ . (04 Marks)
- c. Show that the radius of curvature at any point ' $\theta$ ' to the curve  $x = a(\theta + \sin \theta)$ ,  $y = a(1 - \cos \theta)$ , is  $4a \cos(\frac{\theta}{2})$ . (06 Marks)
- d. Evaluate  $\lim_{x \rightarrow 0} \left( \frac{a^x + b^x + c^x}{3} \right)^{\frac{1}{x}}$ . (06 Marks)
- 3 a. Choose your answers for the following : (04 Marks)
- If  $u = x^{y-1}$ , then  $\frac{\partial u}{\partial y}$  is  
A)  $x^{y-1} \log x$     B)  $(y-1)x^{y-2}$     C)  $x^{y-1} \log y$     D)  $x^y \log x$
  - If  $Z = f(u, v)$ , where  $u = x + ct$  and  $v = x - ct$ , then  $\frac{\partial Z}{\partial t}$  is given by  
A)  $\frac{\partial z}{\partial u} - \frac{\partial z}{\partial v}$     B)  $\frac{\partial z}{\partial u} + \frac{\partial z}{\partial v}$     C)  $c \left( \frac{\partial z}{\partial u} - \frac{\partial z}{\partial v} \right)$     D)  $c \left( \frac{\partial z}{\partial v} - \frac{\partial z}{\partial u} \right)$
  - If  $x = u(1-v)$ ,  $y = uv$ , then  $J\left(\frac{x, y}{u, v}\right)$  is equal to  
A)  $u$     B)  $\frac{1}{u}$     C)  $uv$     D)  $\frac{u}{v}$
  - The necessary condition for the function  $f(x, y)$  to possess extreme values is  
A)  $f_x = f_y = 0$     B)  $f_{xx} - f_{yy} = 0$     C)  $(f_{xx})(f_{yy}) - f_{xy}^2 = 0$     D)  $f_x > 0, f_y > 0$
- b. If  $u = f\left(\frac{y-x}{xy}, \frac{z-x}{xz}\right)$ , find  $x^2 \frac{\partial u}{\partial x}$ . (04 Marks)
- c. If  $x + y + z = u$ ,  $y + z = v$  and  $z = uvw$ , show that  $J\left(\frac{x, y, z}{u, v, w}\right) = uv$ . (06 Marks)
- d. The Horse power required to propel a steamer is proportional to the square of the distance and cube of the velocity. If the distance is increased by 4% and velocity increased by 3%, find the percentage of increase in the Horse power. (06 Marks)
- 4 a. Choose your answers for the following : (04 Marks)
- If  $\vec{R} = xi + yj + zk$ ,  $|\vec{R}| = r$ , then  $\nabla r^2$  is equal to  
A)  $\frac{\vec{R}}{r^2}$     B)  $\frac{-\vec{R}}{2}$     C)  $\frac{\vec{R}}{r}$     D)  $2\vec{R}$
  - If  $\vec{F} = 3x^2i - xyj + (a-3)xz k$  is solenoidal, then 'a' is equal to  
A) 0    B) -2    C) 2    D) 3
  - If  $\vec{A} = x^2i + y^2j + z^2k$ , then  $\text{curl } \vec{A}$  is given by  
A)  $2xi + 2yj + 2zk$     B) 0    C)  $\frac{xi + yj + zk}{2}$     D)  $2x + 2y + 2z$
  - The scale factors for cylindrical coordinate system  $(\rho \phi z)$  are given by  
A)  $(\rho, 1, 1)$     B)  $(1, \rho, 1)$     C)  $(1, 1, \rho)$     D) None of these
- b. Prove that  $\nabla \cdot \vec{F} = \nabla \phi \cdot \vec{F} + \phi (\nabla \cdot \vec{F})$ . (04 Marks)
- c. If  $\vec{F} = 2xy^3z^4i + 3x^2y^2z^4j + 4x^2y^3z^3k$ , find i)  $(\nabla \cdot \vec{F})$  ii)  $\nabla \times \vec{F}$ . (06 Marks)
- d. Obtain the expression for  $\nabla \cdot \vec{F}$  in orthogonal curvilinear coordinate system  $(u_1 u_2 u_3)$ . (06 Marks)

PART - B

5. a. Choose your answers for the following : (04 Marks)
- Given  $\int_0^1 x^n dx = \frac{1}{n+1}$ , then  $\frac{d^2}{dx^2} \int_0^1 x^n dx$  gives
 

A)  $\int_0^1 (\log x)^2 x^n dx = \frac{2}{(1+n)^2}$   
     B)  $\int_0^1 (\log x)^2 x^n dx = \frac{2}{(1+n)^3}$   
     C)  $\int_0^1 (\log x)^n x^n dx = \frac{2}{(1+n)^2}$   
     D)  $\int_0^1 (\log x)^2 x^n dx = \frac{-2}{(1+n)^3}$
  - The value of the integral  $\int_0^{\pi} \sin^6 x \cos^5 x dx$  is
 

A) 0  
     B)  $\frac{8}{693}$   
     C)  $\frac{8\pi}{693}$   
     D) None of these
  - The volume of the solid generated by revolving the curve  $r = a(1 + \cos\theta)$  about the line  $\theta = 0$  is given by
 

A)  $\frac{2\pi}{3} a^3 \int_0^{\pi} (1 + \cos\theta)^3 \sin\theta d\theta$   
     B)  $\frac{2\pi}{3} a^3 \int_0^{\pi} (1 + \cos\theta)^3 \cos\theta d\theta$   
     C)  $\frac{2\pi}{3} a^3 \int_0^{2\pi} (1 + \cos\theta)^3 \sin\theta d\theta$   
     D)  $\frac{4\pi a^3}{3}$
  - The entire length of the asteroid  $x^{2/3} + y^{2/3} = a^{2/3}$  is
 

A)  $4a$   
     B)  $8a$   
     C)  $6a$   
     D)  $3a$
- b. Obtain the reduction formula of the integral  $\int \cos^n x dx$ . (04 Marks)
- c. Using Leibnitz rule under differentiation under integral sign, evaluate  $\int_0^{\pi} \frac{\log(1 + 2\cos x)}{\cos x} dx$ . (06 Marks)
- d. Find the surface generated by revolving the cycloid  $x = a(\theta - \sin\theta)$ ,  $y = a(1 - \cos\theta)$  about its base, (consider one arc in the 1<sup>st</sup> quadrant). (06 Marks)
6. a. Choose your answers for the following : (04 Marks)
- The general solution of the differential equation  $\frac{dy}{dx} = \sec\left(\frac{y}{x}\right) + \frac{y}{x}$  is
 

A)  $\tan\left(\frac{y}{x}\right) - \log x = c$   
     B)  $\sin\left(\frac{y}{x}\right) - \log x = c$   
     C)  $\operatorname{cosec}\left(\frac{y}{x}\right) - \log x = c$   
     D)  $\cos\left(\frac{y}{x}\right) - \log x = c$
  - Integrating factor for the differential equation  $\frac{dx}{dy} + \frac{2x}{y} = y^2$  is  $\mu = \frac{e^{-2y}}{y^2}$ 

A)  $y^2$   
     B)  $e^{x^2}$   
     C)  $e^{2y}$   
     D)  $e^{y^2}$
  - The general solution of the differential equation  $(x - y) dx + (y - x) dy = 0$  is
 

A)  $\frac{x^2}{2} - y - \frac{y^2}{2} = c$   
     B)  $\frac{x^2}{2} - y + \frac{y^2}{2} = c$   
     C)  $\frac{x^2}{2} - yx + \frac{y^2}{2} = c$   
     D) None of these
  - Given the differential equation of  $f(r, \theta, c) = 0$ , we get differential equation of orthogonal trajectories by changing  $r \frac{d\theta}{dr}$  by
 

A)  $\frac{1}{r} \frac{dr}{d\theta}$   
     B)  $-r^2 \frac{dr}{d\theta}$   
     C)  $\frac{-1}{r} \frac{dr}{d\theta}$   
     D)  $r \frac{dr}{d\theta}$
- b. Solve  $(x^2 - 4xy - 2y^2) dx + (y^2 - 4xy - 2x^2) dy = 0$ . (04 Marks)
- c. Solve  $(x + 2y^3) \frac{dy}{dx} = y$ . (06 Marks)
- d. Find the orthogonal trajectories of the family of curves  $\frac{x^2}{a^2} + \frac{y^2}{b^2 + \lambda} = 1$  (' $\lambda$ ' being the parameter). (06 Marks)

7. a. Choose your answers for the following : (04 Marks)

- i) The rank of the matrix  $\begin{pmatrix} 6 & 1 & 3 & 8 \\ 4 & 2 & 6 & -1 \\ 10 & 3 & 9 & 7 \\ 16 & 4 & 12 & 15 \end{pmatrix}$  is equal to  
 A) 2      B) 3      C) 4      D) 1
- ii) The exact solution of the system of equations  $10x + y + z = 12$ ,  $x + 10y + z = 12$ ,  $x + y + 10z = 12$  by inspection is equal to  
 A)  $[0 \ 0 \ 0]^T$       B)  $[1 \ 1 \ 1]^T$       C)  $[1 \ 1 \ -1]^T$       D)  $[-1 \ -1 \ -1]^T$
- iii) If the given system of linear equations in 'n' variables is consistent then the number of linearly independent solution is given by  
 A) n      B) n - 1      C) r - n      D) n - r  
 (Where 'r' stands for rank of co-efficient matrix).
- iv) The trivial solution for the given system of equations  
 $qx - y + 4z = 0$ ,  $4x - 2y + 3z = 0$ ,  $5x + y - 6z = 0$  is  
 A) (1, 2, 0)      B) (0 4 1)      C) (0 0 0)      D) (1 -5 0)
- b. Using elementary row transformations find the rank of the matrix  $\begin{pmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 1 & 1 \\ 3 & 1 & 0 & 2 \\ 1 & 1 & -2 & 0 \end{pmatrix}$ . (04 Marks)
- c. Test for consistency and solve the system of equations  $x + 4 + 3z = 0$ ,  $x - y + z = 0$ ,  $2x - y + 3z = 0$ . (06 Marks)
- d. Applying Gauss Jordan method solve  $2x + 3y - z = 5$ ,  $4x + 4y - 3z = 3$ ,  $2x - 3y + 2z = 2$ . (06 Marks)

8. a. Choose your answers for the following : (04 Marks)

- i) The linear transformation  $y = Ax$  is regular if  
 A)  $|A| = 0$       B)  $|A| = 1$       C)  $|A| = -1$       D)  $|A| \neq 0$
- ii) The transformation  $\xi = x \cos\alpha - y \sin\alpha$ ,  $\eta = x \sin\alpha + y \cos\alpha$  is orthogonal then the inverse of the transformation matrix is given by  
 A)  $\begin{pmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{pmatrix}$       B)  $\begin{pmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{pmatrix}$       C)  $\begin{pmatrix} \sin\alpha & \cos\alpha \\ \cos\alpha & -\sin\alpha \end{pmatrix}$       D)  $\begin{pmatrix} -\sin\alpha & \cos\alpha \\ \cos\alpha & \sin\alpha \end{pmatrix}$
- iii) The eigen vector 'x' of the matrix 'A' corresponding to eigen value ' $\lambda$ ' satisfy the equation  
 A)  $AX = \lambda X$       B)  $\lambda(A - X) = 0$       C)  $XA - \lambda A = 0$       D)  $|A - \lambda I|X = 0$
- iv) Two square matrices A and B are similar if  
 A)  $A = B$       B)  $B = P^{-1}AP$       C)  $A^T = B^T$       D)  $A^{-1} = B^{-1}$
- b. Show that the transformation given below  $y_1 = 2x_1 + x_2 + x_3$ ,  $y_2 = x_1 + x_2 + 2x_3$ ,  $y_3 = x_1 - 2x_3$  is regular and find the inverse transformation. (04 Marks)
- c. Find the matrix P which diagonalizes the matrix  $A = \begin{bmatrix} -1 & 1 & 2 \\ 0 & -2 & -1 \\ 0 & 0 & -3 \end{bmatrix}$ . (06 Marks)
- d. Reduce the quadratic form  $x_1^2 + 3x_2^2 + 3x_3^2 - 2x_2x_3$  in to canonical form by an appropriate orthogonal transformation which transforms  $x_1$   $x_2$   $x_3$  in terms of new variables  $y_1$   $y_2$   $y_3$ . (06 Marks)

\* \* \* \*